



**EUROPEAN COMMISSION
Directorate General
Joint Research Centre**

Directorate F – Health, Consumers and Reference Materials

CERTIFIED REFERENCE MATERIALS 2018

Reference Materials Unit
Retieseweg 111
B - 2440 Geel, Belgium
Fax: +32-(0)14-590 406
Tel.: +32-(0)14-571 705
e-mail: jrc-rm-distribution@ec.europa.eu
Online catalogue: <https://crm.jrc.ec.europa.eu>
Information on CRMs in general: <https://ec.europa.eu/jrc/en/reference-materials>

TABLE OF CONTENTS

INTRODUCTION	1
1 MATERIALS RELATED TO ENVIRONMENTAL ANALYSIS	4
1.1 PURE MATERIALS AND SYNTHETIC MIXTURES	4
1.2 MATRIX MATERIALS	13
1.2.1 CERTIFIED FOR THE TOTAL ELEMENT CONTENT	13
1.2.2 CERTIFIED FOR THE EXTRACTABLE ELEMENT CONTENT AND SPECIES	212
1.2.3 CERTIFIED FOR ORGANIC POLLUTANTS.....	24
1.2.4 OTHERS.....	28
2 MATERIALS RELATED TO THE ANALYSIS OF FOOD AND FEEDING STUFF ...	29
2.1 PURE MATERIALS AND SYNTHETIC MIXTURES	29
2.2 MATRIX MATERIALS	31
2.2.1 CERTIFIED FOR GMO CONTENT.....	31
2.2.2 CERTIFIED FOR NATURAL TOXINS AND XENOBIOTICS.....	40
2.2.3 CERTIFIED FOR THE TOTAL ELEMENT CONTENT	45
2.2.4 CERTIFIED FOR PROXIMATES AND CONVENTIONAL PROPERTIES.....	47
2.2.5 CERTIFIED FOR MICROBIOLOGICAL PROPERTIES	51
2.2.6 CERTIFIED FOR VETERINARY DRUGS.....	54
2.2.7 CERTIFIED FOR IDENTITY.....	56
2.2.8 OTHERS.....	57
3 MATERIALS RELATED TO CLINICAL CHEMISTRY	57
3.1 PURE STANDARDS AND SYNTHETIC MATERIALS	57
3.2 MATRIX MATERIALS	58
3.2.1 CERTIFIED FOR THE HORMONE CONTENT.....	58
3.2.2 CERTIFIED FOR THE TOTAL ELEMENT CONTENT AND OTHER PROPERTIES.....	59
3.2.3 CERTIFIED FOR PROTEIN CONTENT.....	60
3.2.4 CERTIFIED FOR CATALYTIC ACTIVITY	62
3.2.5 CERTIFIED FOR DNA SEQUENCE AND MASS CONCENTRATION	63
3.2.6 OTHERS.....	64
4 MATERIALS CERTIFIED FOR PHYSICAL PROPERTIES	64
4.1 CERTIFIED FOR THERMAL PROPERTIES.....	64
4.2 CERTIFIED FOR MECHANICAL PROPERTIES	66
4.3 CERTIFIED FOR MORPHOLOGICAL PROPERTIES	68

5 MATERIALS RELATED TO INDUSTRIAL APPLICATIONS	71
5.1 CERTIFIED FOR COMPOSITION.....	71
5.2 CERTIFIED FOR TRACE ELEMENT CONTENT	73
5.3 OTHERS.....	79
6 MATERIALS RELATED TO ISOTOPIC MEASUREMENTS.....	80
6.1 CERTIFIED FOR ISOTOPE ABUNDANCE RATIO (AMOUNT RATIO).....	80
6.2 CERTIFIED FOR ISOTOPE AMOUNT CONTENT	82
INDEX.....	84
NUMERICAL LIST	844
ALPHABETICAL LIST.....	95

INTRODUCTION

Public confidence in measurement results is important in many aspects of modern society, including consumer protection in food consumption, health-care, environmental protection, and fair trade. Certified Reference Materials (CRMs) are cornerstones of modern analytical quality assurance because they allow calibration of instruments, validation of methods, and quality control of methods and laboratories based on traceability and comparability of measurement results.

The JRC-Geel provides

IRMM certified reference materials, produced by the EC-JRC

BCR® certified reference materials (BCR® is a registered trademark of the EC), for which production was supported by research funding of the European Commission, DG Research

and

ERM® certified reference materials (ERM® is a registered trademark of the EC), a brand launched in 2003 for CRMs of the highest quality.

These CRMs are produced according to specific Guidelines of the European Commission which are in agreement with the relevant ISO Guides 34 and 35. The JRC-Geel has been accredited for the production of reference materials since 2004 (accreditation certificate BELAC 268-RM)

Certificates

Certificates carry a certified value with its uncertainty which is traceable either to a SI unit or an internationally accepted reference. The intended use for each CRM is stated on the certificate.

CRMs are stored under controlled conditions which ensure their stability. Monitoring programmes have been set up to control CRM stability during the whole shelf-life.

At present JRC-Geel's Reference Materials Unit offers about 700 different CRMs. A complete list of these CRMs, including description and price as well as the CRM catalogue and all certificates and certification reports can be accessed directly via the online catalogue:

<https://crm.jrc.ec.europa.eu>

To order a free hardcopy of the catalogue simply write to:

jrc-rm-distribution@ec.europa.eu

Availability

As CRMs should become regularly used items in any measurement laboratory, JRC-Geel's Reference Materials Unit continues its effort to supply CRMs in sufficient amount to cover market needs. However, in exceptional cases where only limited amounts are available, it reserves the right to restrict and/or refuse orders. The reference materials contained in this catalogue are made available world-wide through the JRC-Geel's Reference Materials Unit and its authorised distributors.

They can be purchased from the

European Commission
Directorate General Joint Research Centre
Directorate F – Health, Consumers and Reference Materials
Reference Materials Unit
Retieseweg 111, B-2440 Geel, Belgium
Tel. +32-(0)14-571 705, Fax +32-(0)14-590 406
jrc-rm-distribution@ec.europa.eu

or from

AUTORISED DISTRIBUTORS OF CRMs

LGC STANDARDS GmbH
Louis-Pasteur-Strasse 30
D-14943 Luckenwalde
Germany
Tel. +49 3371 68 16 80
Fax +49 3371 68 16 89
<http://www.lgcstandards.com>

SIGMA-ALDRICH CHEMIE GmbH
Industriestrasse 25
CH-9471 Buchs
Switzerland
Tel. +41 81 755 2828
Fax +41 81 755 2815
<http://www.sigmaaldrich.com>

Sigma-Aldrich RTC Inc.

P.O. Box 1346, 2931 Soldier Springs Road
Laramie, WY 82070
USA
Tel. +1 307 742 5452
Fax +1 307 745 7936
<http://www.sigmaaldrich.com>

LGC STANDARDS US - ARMI

1 Perimeter Road, Suite 200
Manchester, NH 03103
USA
Tel. +1 603 935 4100
Fax +1 603 935 4101
<https://us.lgcstandards.com/us/en/armi>

Prices and sales conditions are freely determined by each distributor and can be obtained from them on request. They can provide advice and help in ordering and importing reference materials.

If samples are purchased through **UNAUTHORISED COMPANIES**, the JRC-Geel cannot be held responsible for the integrity of the materials (especially in case of potentially unstable materials) nor, for the accuracy and/or completeness of the accompanying information (certificates, reports, etc.).

1 MATERIALS RELATED TO ENVIRONMENTAL ANALYSIS

1.1 PURE MATERIALS AND SYNTHETIC MIXTURES

	Substance	Purity (g/g)		
	Polycyclic aromatic compounds			
BCR-046	BENZO[b]CHRYSENE	0.994	+	0.006
		-		0.008
BCR-047	BENZO[b]FLUORANTHENE	0.997	4	0.002 6
BCR-048R	BENZO[k]FLUORANTHENE (unit size 10 mg)	0.997	+	0.003
		-		0.004
BCR-049	BENZO[j]FLUORANTHENE	0.997	±	0.003
		±		0.006
BCR-050	BENZO[e]PYRENE	0.991	+	0.009
		-		0.010
BCR-052	BENZO[ghi]PERYLENE	0.992	3	0.002 1
BCR-077R	1-METHYLCHRYSENE (unit size 10 mg)	0.991	±	0.007
BCR-078R	2-METHYLCHRYSENE (unit size 10 mg)	0.993	±	0.005
BCR-079R	3-METHYLCHRYSENE (unit size 10 mg)	0.993	±	0.005
BCR-080R	4-METHYLCHRYSENE (unit size 10 mg)	0.994	±	0.004
BCR-081R	5-METHYLCHRYSENE (unit size 10 mg)	0.997	3	0.001 3
BCR-091	ANTHANTHRENE	0.996	±	0.004
BCR-092	10-AZABENZO[a]PYRENE	0.996	±	0.006
BCR-093R	1-METHYLBENZ[a]ANTHRACENE (unit size 10 mg)	0.996	±	0.005
BCR-094	DIBENZ[a,c]ANTHRACENE	0.996	±	0.004
BCR-095	DIBENZ[a,j]ANTHRACENE	0.997	8	0.002 5
BCR-096	DIBENZO[a,l]PYRENE	0.997	2	0.002 5
BCR-097	BENZO[a]FLUORANTHENE	0.996	±	0.004
BCR-133	DIBENZO[a,e]PYRENE	0.996	+	0.004
		-		0.005
BCR-134	BENZO[c]PHENANTHRENE	0.996	8	0.001 4
BCR-136R	BENZO[b]NAPHTHO[2,3-d]THIOPHENE (unit size 10 mg)	0.994	±	0.006
BCR-137R	BENZO[b]NAPHTHO[1,2-d]THIOPHENE (unit size 10 mg)	0.996	6	0.002 9
BCR-138	DIBENZ[a,h]ANTHRACENE	0.990	±	0.007
BCR-139	BENZO[ghi]FLUORANTHENE	0.995	±	0.004
BCR-140	BENZO[c]CHRYSENE	0.996	+	0.004
		-		0.005
BCR-153R	DIBENZ[a,h]ACRIDINE (unit size 10 mg)	0.999	2	0.000 6
BCR-154	DIBENZ[a,j]ACRIDINE	0.999	0	0.000 7
		-		0.001 0
BCR-155	DIBENZ[a,c]ACRIDINE	0.999	1	0.000 7
		-		0.000 8
BCR-156R	DIBENZ[c,h]ACRIDINE (unit size 10 mg)	0.993	6	0.002 1
BCR-157	BENZ[a]ACRIDINE	0.998	2	0.001 8
BCR-158	BENZ[c]ACRIDINE	0.998	7	0.001 3
		-		0.001 8
BCR-159	DIBENZO[a,h]PYRENE	0.993	±	0.007
BCR-160R	FLUORANTHENE (unit size 10 mg)	0.996	+	0.004
		-		0.005
BCR-168	PICENE (unit size 10 mg)	0.998	+	0.001 3
		-		0.004
BCR-177R	PYRENE (unit size 10 mg)	0.998	0	0.000 4

Availability: Amber vials containing about 100 mg of powdered material unless otherwise specified.

	Substance	Purity (g/g)		
	Polycyclic aromatic compounds			
BCR-152	DIBENZ[a,i]ACRIDINE	0.998	5	0.001 0
		-		0.000 8
BCR-265	DIBENZO[a,e]FLUORANTHENE	0.998	5	0.001 6
		-		0.001 0
BCR-266	7H-DIBENZO[c,g]CARBAZOLE	0.997	1	0.001 6
BCR-267	INDENO[1,2,3-cd]FLUORANTHENE	0.998	6	0.000 9
		-		0.000 8
BCR-269	CHRYSENE	0.992	8	0.002 8
BCR-270	TRIPHENYLENE	0.998	4	0.001 0
		-		0.000 6
BCR-271	BENZ[a]ANTHRACENE	0.998	4	0.000 9
BCR-272	CORONENE	0.998	9	0.000 6
		-		0.000 4

Availability: Amber vials containing about 20 mg of powdered material.

Nitro-polycyclic aromatic hydrocarbons						
BCR-305	1-NITROPYRENE		0.997 6	±	0.000 7	
BCR-306	1-NITRONAPHTALENE		0.996 9	±	0.001 0	
BCR-307	2-NITRONAPHTALENE		0.997 7	+	0.000 9	- 0.001 1
BCR-308	9-NITROANTHRACENE		0.997 5	±	0.001 0	
BCR-309	6-NITROCHRYSENE		0.989	±	0.004	
BCR-310	3-NITROFLUORANTHENE		0.996 8	+	0.001 2	- 0.002 1
BCR-311	6-NITROBENZO[a]PYRENE		0.997 8	+	0.000 8	- 0.001 0
BCR-312	2-NITRO-7-METHOXYNAPHTHO[2.1-b]FURAN		0.998 4	±	0.000 7	

Availability: Amber vials containing about 10 mg of powdered material.

Oxygenated polycyclic aromatic hydrocarbons						
BCR-337	DIBENZO[b,d]FURAN		0.987	±	0.007	
BCR-339	6H-BENZO[c,d]PYREN-6-ONE		0.988	±	0.009	
BCR-340	BENZO[b]NAPHTHO[1,2-d]FURAN		0.997	+	0.003	- 0.005
BCR-341	BENZO[b]NAPHTHO[2,1-d]FURAN		0.996	+	0.004	- 0.005
BCR-342	BENZO[a]FLUORENONE		0.997 9	+	0.002 1	- 0.002 2

Availability: Amber vials containing about 10 mg of powdered material.

Polychlorinated biphenyls						
IUPAC No.						
BCR-289	8	2,4' - DICHLOROBIPHENYL		0.996 3	+	0.005
					-	0.001 8
BCR-290	20	2,3,3' - TRICHLOROBIPHENYL		0.998 5	±	0.001 3
BCR-291	28	2,4,4' - TRICHLOROBIPHENYL		0.997 9	±	0.001 3
BCR-293	52	2,2',5,5' - TETRACHLOROBIPHENYL		0.995 9	±	0.002 5
BCR-296	138	2,2',3,4,4',5' - HEXACHLOROBIPHENYL		0.999 2	±	0.000 7
BCR-297	153	2,2',4,4',5,5' - HEXACHLOROBIPHENYL		0.999 4	+	0.000 9
					-	0.000 5
BCR-298	180	2,2',3,4,4',5,5' - HEPTACHLOROBIPHENYL		0.995 7	±	0.001 4

Availability: Amber vials containing about 25 mg of powdered material.

Polychlorinated biphenyls in iso-octane (BCR-365)

IUPAC No.	Content in mg/kg	Concentration in g/m ³ at 25 °C ¹⁾
8	11.4 ± 0.4	(7.8 ± 0.2)
20	15.2 ± 0.9	(10.5 ± 0.7)
28	24.8 ± 1.1	(17.1 ± 0.8)
35	14.3 ± 0.8	(9.8 ± 0.5)
52	14.8 ± 0.6	(10.2 ± 0.4)
101	14.4 ± 0.6	(9.9 ± 0.4)
118	14.9 ± 0.8	(10.3 ± 0.6)
138	8.6 ± 0.6	(5.9 ± 0.5)
153	14.2 ± 0.6	(9.8 ± 0.4)
180	15.2 ± 0.6	(10.4 ± 0.3)

1) Not certified concentrations (g/m³) were calculated from the certified content assuming a density of iso-octane of 687.77 kg/m³ at 25 °C.

Availability: Unit consisting of a pair of dark glass ampoules, each containing 2 cm³ of 2,2,4-Trimethylpentane (iso-octane) sealed under nitrogen.

The pair of ampoules is supplied in a metal can which is packed with absorbent material.

Polychlorodibenzo-p-dioxins (PCDDs) and polychlorodibenzofurans (PCDFs)

BCR-614 SOLUTION – S0

Congener	Certified mass fraction ¹⁾ ($\mu\text{g/kg}$)	Uncertainty ²⁾ ($\mu\text{g/kg}$)	Mass fraction expressed in concentration units ³⁾ ($\mu\text{g/L}$)	Uncertainty expressed in concentration units ³⁾ ($\mu\text{g/L}$)
2,3,7,8-T ₄ CDD	0.137	0.004	0.098 3	0.002 9
1,2,3,7,8-P ₅ CDD	0.698	0.014	0.501	0.010
1,2,3,4,7,8-HCDD	0.688	0.021	0.494	0.015
1,2,3,6,7,8-HCDD	0.696	0.006	0.500	0.004
1,2,3,7,8,9-HCDD	0.705	0.008	0.506	0.006
1,2,3,4,6,7,8-HCDD	1.400	0.020	1.005	0.014
1,2,3,4,6,7,8,9-O ₈ CDD	1.396	0.007	1.001	0.005
2,3,7,8-T ₄ CDF	0.139 7	0.001 1	0.100 2	0.000 8
1,2,3,7,8-P ₅ CDF	0.707	0.013	0.507	0.009
2,3,4,7,8-P ₅ CDF	0.698	0.005	0.501	0.004
1,2,3,4,7,8-HCDF	0.700	0.006	0.502	0.005
1,2,3,6,7,8-HCDF	0.698	0.005	0.501	0.004
1,2,3,7,8,9-HCDF	0.699	0.009	0.502	0.007
2,3,4,6,7,8-HCDF	0.694	0.007	0.498	0.005
1,2,3,4,6,7,8-HCDF	1.396	0.008	1.001	0.006
1,2,3,4,7,8,9-HCDF	1.394	0.030	1.001	0.022
1,2,3,4,6,7,8,9-O ₈ CDF	1.397	0.024	1.002	0.017

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

BCR-614 SOLUTION – S0

Congener	Certified mass fraction ¹⁾ ($\mu\text{g/kg}$)	Uncertainty ²⁾ ($\mu\text{g/kg}$)	Mass fraction expressed in concentration units ³⁾ ($\mu\text{g/L}$)	Uncertainty expressed in concentration units ³⁾ ($\mu\text{g/L}$)
¹³ C-2,3,7,8-T ₄ CDD	13.95	0.06	10.01	0.05
¹³ C-1,2,3,7,8-P ₅ CDD	13.9	0.4	10.00	0.23
¹³ C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
¹³ C-1,2,3,6,7,8-HCDD	13.94	0.24	10.00	0.17
¹³ C-1,2,3,7,8,9-HCDD	13.95	0.10	10.01	0.07
¹³ C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDD	27.87	0.16	20.00	0.12
¹³ C-2,3,7,8-T ₄ CDF	13.96	0.09	10.02	0.07
¹³ C-1,2,3,7,8-P ₅ CDF	13.94	0.24	10.00	0.17
¹³ C-2,3,4,7,8-P ₅ CDF	13.95	0.06	10.01	0.05
¹³ C-1,2,3,4,7,8-HCDF	13.90	0.07	9.97	0.05
¹³ C-1,2,3,6,7,8-HCDF	13.93	0.10	10.00	0.08
¹³ C-1,2,3,7,8,9-HCDF	13.93	0.10	10.00	0.07
¹³ C-2,3,4,6,7,8-HCDF	13.93	0.09	10.00	0.06
¹³ C-1,2,3,4,6,7,8-HCDF	27.92	0.20	20.03	0.15
¹³ C-1,2,3,4,7,8,9-HCDF	27.87	0.24	20.00	0.17
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDF	27.88	0.25	20.01	0.18
¹³ C-1,2,3,4-T ₄ CDD	13.94	0.08	10.00	0.06

BCR-614 SOLUTION – S1

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
2,3,7,8-T ₄ CDD	0.273	0.008	0.196	0.006
1,2,3,7,8-P ₅ CDD	1.394	0.027	1.000	0.020
1,2,3,4,7,8-HCDD	1.37	0.05	0.986	0.030
1,2,3,6,7,8-HCDD	1.391	0.010	0.998	0.007
1,2,3,7,8,9-HCDD	1.408	0.015	1.011	0.011
1,2,3,4,6,7,8-HCDD	2.80	0.04	2.006	0.028
1,2,3,4,6,7,8,9-O ₈ CDD	2.787	0.010	2.000	0.007
2,3,7,8-T ₄ CDF	0.279 0	0.002 1	0.200 2	0.001 5
1,2,3,7,8-P ₅ CDF	1.412	0.025	1.013	0.018
2,3,4,7,8-P ₅ CDF	1.395	0.008	1.001	0.006
1,2,3,4,7,8-HCDF	1.398	0.011	1.003	0.008
1,2,3,6,7,8-HCDF	1.393	0.009	1.000	0.006
1,2,3,7,8,9-HCDF	1.397	0.017	1.002	0.012
2,3,4,6,7,8-HCDF	1.387	0.012	0.995	0.009
1,2,3,4,6,7,8-HCDF	2.787	0.012	2.000	0.009
1,2,3,4,7,8,9-HCDF	2.78	0.06	2.00	0.05
1,2,3,4,6,7,8,9-O ₈ CDF	2.79	0.05	2.00	0.04
¹³ C-2,3,7,8-T ₄ CDD	13.95	0.06	10.01	0.05
¹³ C-1,2,3,7,8-P ₅ CDD	13.9	0.4	10.00	0.23
¹³ C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
¹³ C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
¹³ C-1,2,3,7,8,9-HCDD	13.94	0.10	10.01	0.07
¹³ C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDD	27.86	0.16	19.99	0.11
¹³ C-2,3,7,8-T ₄ CDF	13.96	0.09	10.01	0.07
¹³ C-1,2,3,7,8-P ₅ CDF	13.93	0.24	10.00	0.17
¹³ C-2,3,4,7,8-P ₅ CDF	13.94	0.06	10.00	0.05
¹³ C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
¹³ C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
¹³ C-1,2,3,7,8,9-HCDF	13.92	0.10	9.99	0.07
¹³ C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
¹³ C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.14
¹³ C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDF	27.87	0.25	20.00	0.18
¹³ C-1,2,3,4-T ₄ CDD	13.93	0.07	10.00	0.05

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

BCR-614 SOLUTION – S2

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
2,3,7,8-T ₄ CDD	1.09	0.04	0.785	0.023
1,2,3,7,8-P ₅ CDD	5.57	0.11	4.00	0.08
1,2,3,4,7,8-HCDD	5.49	0.17	3.94	0.12
1,2,3,6,7,8-HCDD	5.56	0.04	3.992	0.027
1,2,3,7,8,9-HCDD	5.63	0.06	4.04	0.05
1,2,3,4,6,7,8-HCDD	11.18	0.16	8.02	0.11
1,2,3,4,6,7,8,9-O ₈ CDD	11.15	0.04	8.000	0.027
2,3,7,8-T ₄ CDF	1.116	0.008	0.801	0.006
1,2,3,7,8-P ₅ CDF	5.65	0.10	4.05	0.07
2,3,4,7,8-P ₅ CDF	5.58	0.03	4.004	0.022
1,2,3,4,7,8-HCDF	5.59	0.05	4.01	0.04
1,2,3,6,7,8-HCDF	5.57	0.04	3.999	0.024
1,2,3,7,8,9-HCDF	5.59	0.07	4.01	0.05
2,3,4,6,7,8-HCDF	5.55	0.05	3.98	0.04
1,2,3,4,6,7,8-HCDF	11.15	0.05	8.00	0.04
1,2,3,4,7,8,9-HCDF	11.14	0.24	7.99	0.17
1,2,3,4,6,7,8,9-O ₈ CDF	11.16	0.19	8.01	0.14
¹³ C-2,3,7,8-T ₄ CDD	13.95	0.06	10.01	0.05
¹³ C-1,2,3,7,8-P ₅ CDD	13.9	0.4	10.00	0.23
¹³ C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
¹³ C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
¹³ C-1,2,3,7,8,9-HCDD	13.94	0.10	10.01	0.07
¹³ C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDD	27.86	0.16	19.99	0.11
¹³ C-2,3,7,8-T ₄ CDF	13.96	0.09	10.01	0.07
¹³ C-1,2,3,7,8-P ₅ CDF	13.93	0.24	10.00	0.17
¹³ C-2,3,4,7,8-P ₅ CDF	13.94	0.06	10.00	0.05
¹³ C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
¹³ C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
¹³ C-1,2,3,7,8,9-HCDF	13.93	0.10	9.99	0.07
¹³ C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
¹³ C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.15
¹³ C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDF	27.87	0.25	20.00	0.18
¹³ C-1,2,3,4-T ₄ CDD	13.93	0.08	10.00	0.06

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

BCR-614 SOLUTION – S3

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
2,3,7,8-T ₄ CDD	5.47	0.16	3.92	0.12
1,2,3,7,8-P ₅ CDD	27.9	0.6	20.0	0.4
1,2,3,4,7,8-HCDD	27.5	0.9	19.7	0.6
1,2,3,6,7,8-HCDD	27.81	0.19	19.96	0.14
1,2,3,7,8,9-HCDD	28.17	0.30	20.21	0.21
1,2,3,4,6,7,8-HCDD	55.9	0.8	40.1	0.6
1,2,3,4,6,7,8,9-O ₈ CDD	55.74	0.19	40.00	0.14
2,3,7,8-T ₄ CDF	5.58	0.04	4.003	0.029
1,2,3,7,8-P ₅ CDF	28.2	0.5	20.3	0.4
2,3,4,7,8-P ₅ CDF	27.90	0.16	20.02	0.11
1,2,3,4,7,8-HCDF	27.96	0.22	20.06	0.16
1,2,3,6,7,8-HCDF	27.87	0.17	20.00	0.12
1,2,3,7,8,9-HCDF	27.9	0.4	20.04	0.24
2,3,4,6,7,8-HCDF	27.73	0.23	19.90	0.17
1,2,3,4,6,7,8-HCDF	55.74	0.24	40.00	0.17
1,2,3,4,7,8,9-HCDF	55.7	1.2	40.0	0.9
1,2,3,4,6,7,8,9-O ₈ CDF	55.8	1.0	40.0	0.7
¹³ C-2,3,7,8-T ₄ CDD	13.95	0.06	10.01	0.05
¹³ C-1,2,3,7,8-P ₅ CDD	13.9	0.4	10.00	0.23
¹³ C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
¹³ C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
¹³ C-1,2,3,7,8,9-HCDD	13.95	0.10	10.01	0.07
¹³ C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDD	27.87	0.16	20.00	0.11
¹³ C-2,3,7,8-T ₄ CDF	13.96	0.09	10.02	0.07
¹³ C-1,2,3,7,8-P ₅ CDF	13.93	0.24	10.00	0.17
¹³ C-2,3,4,7,8-P ₅ CDF	13.94	0.06	10.01	0.05
¹³ C-1,2,3,4,7,8-HCDF	13.90	0.07	9.97	0.05
¹³ C-1,2,3,6,7,8-HCDF	13.93	0.11	10.00	0.08
¹³ C-1,2,3,7,8,9-HCDF	13.93	0.10	10.00	0.07
¹³ C-2,3,4,6,7,8-HCDF	13.93	0.09	10.00	0.06
¹³ C-1,2,3,4,6,7,8-HCDF	27.91	0.20	20.03	0.14
¹³ C-1,2,3,4,7,8,9-HCDF	27.87	0.24	20.00	0.17
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDF	27.88	0.25	20.00	0.18
¹³ C-1,2,3,4-T ₄ CDD	13.93	0.07	10.00	0.05

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

BCR-614 SOLUTION – S4

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
2,3,7,8-T ₄ CDD	27.3	0.8	9.6	0.6
1,2,3,7,8-P ₅ CDD	139.3	2.7	100.0	2.0
1,2,3,4,7,8-HCDD	137	5	98.6	3.0
1,2,3,6,7,8-HCDD	139.1	1.0	99.8	0.7
1,2,3,7,8,9-HCDD	140.8	1.5	101.1	1.1
1,2,3,4,6,7,8-HCDD	280	4	200.6	2.8
1,2,3,4,6,7,8,9-O ₈ CDD	278.7	1.0	200.0	0.7
2,3,7,8-T ₄ CDF	27.89	0.21	20.02	0.15
1,2,3,7,8-P ₅ CDF	141.2	2.5	101.3	1.8
2,3,4,7,8-P ₅ CDF	139.5	0.8	100.1	0.6
1,2,3,4,7,8-HCDF	139.8	1.1	100.3	0.8
1,2,3,6,7,8-HCDF	139.3	0.9	100.0	0.6
1,2,3,7,8,9-HCDF	139.6	1.7	100.2	1.2
2,3,4,6,7,8-HCDF	138.7	1.2	99.5	0.9
1,2,3,4,6,7,8-HCDF	278.7	1.2	200.0	0.9
1,2,3,4,7,8,9-HCDF	278	6	200	5
1,2,3,4,6,7,8,9-O ₈ CDF	279	5	200	4
¹³ C-2,3,7,8-T ₄ CDD	13.95	0.06	10.01	0.05
¹³ C-1,2,3,7,8-P ₅ CDD	13.99	0.4	10.00	0.23
¹³ C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
¹³ C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
¹³ C-1,2,3,7,8,9-HCDD	13.94	0.10	10.01	0.07
¹³ C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDD	27.86	0.16	19.99	0.11
¹³ C-2,3,7,8-T ₄ CDF	13.96	0.09	10.01	0.07
¹³ C-1,2,3,7,8-P ₅ CDF	13.93	0.24	10.00	0.17
¹³ C-2,3,4,7,8-P ₅ CDF	13.94	0.06	10.00	0.05
¹³ C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
¹³ C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
¹³ C-1,2,3,7,8,9-HCDF	13.92	0.10	9.99	0.07
¹³ C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
¹³ C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.15
¹³ C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDF	27.87	0.25	20.00	0.18
¹³ C-1,2,3,4-T ₄ CDD	13.93	0.08	10.00	0.06

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

BCR-614 SOLUTION – S5

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
2,3,7,8-T ₄ CDD	109	4	78.5	2.3
1,2,3,7,8-P ₅ CDD	557	11	400	8
1,2,3,4,7,8-HCDD	549	17	394	12
1,2,3,6,7,8-HCDD	556	4	399.1	2.7
1,2,3,7,8,9-HCDD	563	6	404	5
1,2,3,4,6,7,8-HCDD	1118	16	802	11
1,2,3,4,6,7,8,9-O ₈ CDD	1115	4	799.9	2.7
2,3,7,8-T ₄ CDF	1116	0.8	80.1	0.6
1,2,3,7,8-P ₅ CDF	565	0	405	7
2,3,4,7,8-P ₅ CDF	558	3	400.4	2.2
1,2,3,4,7,8-HCDF	559	5	401	4
1,2,3,6,7,8-HCDF	557	4	399.9	2.4
1,2,3,7,8,9-HCDF	559	7	401	5
2,3,4,6,7,8-HCDF	555	5	398	4
1,2,3,4,6,7,8-HCDF	1115	5	800	4
1,2,3,4,7,8,9-HCDF	1114	24	799	17
1,2,3,4,6,7,8,9-O ₈ CDF	1116	19	801	14
¹³ C-2,3,7,8-T ₄ CDD	13.95	0.06	10.01	0.05
¹³ C-1,2,3,7,8-P ₅ CDD	13.9	0.4	10.00	0.23
¹³ C-1,2,3,4,7,8-HCDD	13.98	0.7	10.03	0.05
¹³ C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
¹³ C-1,2,3,7,8,9-HCDD	13.95	0.10	10.01	0.07
¹³ C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDD	27.86	0.16	19.99	0.11
¹³ C-2,3,7,8-T ₄ CDF	13.96	0.09	10.02	0.07
¹³ C-1,2,3,7,8-P ₅ CDF	13.93	0.24	10.00	0.17
¹³ C-2,3,4,7,8-P ₅ CDF	13.94	0.06	10.00	0.05
¹³ C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
¹³ C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
¹³ C-1,2,3,7,8,9-HCDF	13.93	0.10	9.99	0.07
¹³ C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
¹³ C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.15
¹³ C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDF	27.87	0.25	20.00	0.18
¹³ C-1,2,3,4-T ₄ CDD	13.93	0.08	10.00	0.06

BCR-614 SOLUTION – S6

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
¹³ C-1,2,3,7,8-P ₅ CDF	139.3	2.3	100.0	1.7
¹³ C-1,2,3,7,8,9-HCDF	139.4	0.9	100.0	0.7
¹³ C-1,2,3,4,7,8,9-HCDF	278.7	2.4	200.0	1.7

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

BCR-614 SOLUTION – S7

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
¹³ C-2,3,7,8-T ₄ CDD	139.5	0.6	100.1	0.4
¹³ C-1,2,3,7,8-P ₅ CDD	139	4	99.9	2.4
¹³ C-1,2,3,4,7,8-HCDD	139.8	0.7	100.3	0.5
¹³ C-1,2,3,6,7,8-HCDD	139.3	2.4	100.0	1.7
¹³ C-1,2,3,4,6,7,8-HCDD	279	6	200	4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDD	278.7	1.6	200.0	1.1
¹³ C-2,3,7,8-T ₄ CDF	139.5	0.9	100.1	0.6
¹³ C-2,3,4,7,8-P ₅ CDF	139.2	0.6	99.9	0.4
¹³ C-1,2,3,4,7,8-HCDF	138.9	0.6	99.7	0.5
¹³ C-1,2,3,6,7,8-HCDF	139.4	1.1	100.0	0.8
¹³ C-2,3,4,6,7,8-HCDF	139.4	0.8	100.0	0.6
¹³ C-1,2,3,4,6,7,8-HCDF	278.7	2.0	200.0	1.4
¹³ C-1,2,3,4,6,7,8,9-O ₈ CDF	278.7	2.5	200.0	1.8

BCR-614 SOLUTION – S8

Congener	Certified mass fraction ¹⁾ (µg/kg)	Uncertainty ²⁾ (µg/kg)	Mass fraction expressed in concentration units ³⁾ (µg/L)	Uncertainty expressed in concentration units ³⁾ (µg/L)
¹³ C-1,2,3,7,8,9-HCDD	558	4	400.5	2.7
¹³ C-1,2,3,4-T ₄ CDD	557.4	2.7	400.0	2.0

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

Availability: BCR-614 Solutions S1-S7 contain about 1 mL solution and BCR-614 Solution S8 about 0.5 mL. The solutions are available individually or as set. A set consists of 11 ampoules, one of S2, S4-S8 and two of S1 and S3. In addition, the set contains an additional solution S9, which is not certified. This solution is intended as complementary tool for QA/QC purposes.

Substance	ERM-AC213		
	PAHs in Acetonitrile / Toluene (µg/g)		
Benz[a]anthracene	3.09	±	0.04
Chrysene	3.06	±	0.05
5-methylchrysene	3.08	±	0.07
Benzo[b]fluoranthene	3.05	±	0.05
Benzo[k]fluoranthene	3.06	±	0.08
Benzo[j]fluoranthene	3.05	±	0.10
Dibenz[a,h]anthracene	2.76	±	0.05
Benzo[gh]perylene	3.07	±	0.05
Dibenzo[a,l]pyrene	2.85	±	0.10
Dibenzo[a,e]pyrene	2.97	±	0.10
Benzo[c]fluorene	(2.13	±	0.11)
Cyclopental[cd]pyrene	(2.96	±	0.12)
Dibenzo[a,i]pyrene	(2.37	±	0.15)

Values in brackets are not certified.

Availability: ERM-AC213 consists of 2 mL toluene containing 15 PAHs in an ampoule.

1.2 MATRIX MATERIALS

1.2.1 CERTIFIED FOR THE TOTAL ELEMENT CONTENT

Substance	BCR-142R Light sandy soil (mg/kg)	ERM-CC141 Loam soil (mg/kg)	BCR-143R Sewage sludge amended soil (mg/kg)	ERM-CC144 Sewage sludge
As		9.9 ± 1.5		7.7 ± 0.7 mg/kg
Cd	0.34 ± 0.04	0.35 ± 0.05	71.8 ± 1.2	14.5 ± 1.4 mg/kg
Co	12.1 ± 0.7	8.5 ± 0.5	12.3 ± 0.3	6.5 ± 0.4 mg/kg
Cr		86 ± 8		168 ± 14 mg/kg
Cu	69.7 ± 1.3	14.4 ± 1.4	130.6 ± 1.5	348 ± 18 mg/kg
Fe				32.9 ± 1.6 g/kg
Hg	0.067 ± 0.011	0.083 ± 0.017	1.10 ± 0.07	5.9 ± 0.6 mg/kg
Mn	970 ± 16	464 ± 18	904 ± 13	352 ± 14 mg/kg
Ni	64.5 ± 2.5	26.4 ± 2.4	299 ± 5	91 ± 7 mg/kg
Pb	40.2 ± 1.9	41 ± 4	179.7 ± 2.1	157 ± 9 mg/kg
Zn	(101 ± 6)	57 ± 4	1055 ± 14	0.98 ± 0.04 g/kg
Aqua regia soluble ¹⁾				
As	0.249 ± 0.010	7.5 ± 1.4	72.0 ± 1.8	7.7 ± 0.7 mg/kg
Cd	(10.2 ± 0.6)	0.25 ± 0.04	(11.8 ± 1.0)	13.3 ± 0.9 mg/kg
Co		7.9 ± 0.9	426 ± 12	5.9 ± 0.4 mg/kg
Cr	(69.8 ± 1.0)	31 ± 4	(128 ± 7)	150 ± 11 mg/kg
Cu		12.4 ± 0.9	(1.10 ± 0.06)	346 ± 15 mg/kg
Fe				32.8 ± 1.5 g/kg
Hg	(800 ± 50)	0.080 ± 0.008	858 ± 11	6.1 ± 0.7 mg/kg
Mn	61.1 ± 1.5	387 ± 17	296 ± 4	340 ± 13 mg/kg
Ni	25.7 ± 1.6	21.9 ± 1.6	174 ± 5	86 ± 6 mg/kg
Pb	93.3 ± 2.7	32.2 ± 1.4	1063 ± 16	156 ± 7 mg/kg
Zn		50 ± 4		0.97 ± 0.04 g/kg

Values in brackets are not certified.

Availability: Glass bottles containing about 50 g of powdered material; ERM-CC141 contains minimum 24 g; ERM-CC144 minimum 30 g.

¹⁾ Details of the analytical procedure to obtain the aqua regia soluble content of the elements are given in the certification report.

Substance	BCR-667 Estuarine sediment (mg/kg)	ERM-CC690 Calcareous soil (mg/kg)
Br	(99.7 ± 2.5)	
Cd	(0.67 ± 0.11)	
Ce	56.7 ± 2.5	
Co	(23.0 ± 1.3)	
Cr	(178 ± 16)	
Cs	(7.8 ± 0.7)	
Cu	(60 ± 9)	
Dy	4.01 ± 0.14	2.90 ± 0.28
Er	2.35 ± 0.15	
Eu	1.00 ± 0.05	
Fe	(44800 ± 1)	
Gd	4.41 ± 0.12	3.2 ± 0.4
Ho	0.80 ± 0.06	
La	27.8 ± 1.0	24.4 ± 1.7
Lu	0.325 ± 0.020	
Mn	(920 ± 40)	
Nd	25.0 ± 1.4	19.1 ± 2.2
Ni	(128 ± 9)	
Pb	(31.9 ± 1.1)	
Pr	6.1 ± 0.5	
Sb	(0.96 ± 0.05)	
Sc	13.7 ± 0.7	7.9 ± 0.9
Se	(1.59 ± 0.08)	
Sm	4.66 ± 0.20	3.5 ± 0.4
Ta	(0.876 ± 0.017)	
Tb	0.682 ± 0.017	0.50 ± 0.07
Th	10.0 ± 0.5	7.6 ± 0.8
Tm	0.326 ± 0.025	0.232 ± 0.026
U	2.26 ± 0.15	1.90 ± 0.23
Yb	2.20 ± 0.09	1.57 ± 0.19
Zn	(175 ± 13)	

Values in brackets are not certified.

Availability: BCR-667: Glass bottles containing about 40 g of powdered material.

ERM-CC690: Glass bottles containing about 70 g of powdered material. The report gives additional indicative values for As, Au, Co, Cr, Cs, Cu, Er, Eu, Fe, Hf, Ho, Lu, Ni, Pb, Pr, Sb, Ta, W, Y and Zn.

Substance	BCR-277R			BCR-280R			BCR-320R		
	Estuarine sediment (mg/kg)			Lake sediment (mg/kg)			Channel sediment (mg/kg)		
As	18.3	±	1.8	33.4	±	2.9	21.7	±	2.0
Cd	0.61	±	0.07	0.85	±	0.10	2.64	±	0.18
Co	22.5	±	1.4	16.8	±	0.9	9.7	±	0.6
Cr	188	±	14	126	±	7	59	±	4
Cu	63	±	7	53	±	6	46.3	±	2.9
Fe							25700	±	1300
Hg	0.128	±	0.017	1.46	±	0.20	0.85	±	0.09
Mn							910	±	50
Ni	130	±	8	69	±	5	27.1	±	2.2
Pb							85	±	5
Sc							5.2	±	0.4
Se	(0.58	±	0.11)	(0.46	±	0.09)	(0.96	±	0.18)
Sn	(6.5	±	1.8)	(9.5	±	1.7)	(9.4	±	1.7)
Th							5.3	±	0.4
Tl							0.65	±	0.08
U							1.56	±	0.20
V							46.5	±	2.8
Zn	178	+	20	224	+	25	318	+	20

Values in brackets are not certified.

Availability: Amber glass bottles containing 40 g of powder for BCR-277R and BCR-320R and 30 g for BCR-280R.

Substance	BCR-145R			BCR-146R		
	Sewage sludge (mixed origin) (mg/kg)			Sewage sludge (industrial origin) (mg/kg)		
Cd	3.50	±	0.15	18.8	±	0.5
Co	5.6	±	0.4	7.39	±	0.27
Cr				196	±	7
Cu	696	±	12	838	±	16
Hg	2.01	±	0.22	8.6	±	0.4
Mn	156	±	4	323	±	7
Ni	247	±	7	70	±	5
Pb	286	±	5	609	±	14
Zn	2122	±	23	3060	±	60
Aqua regia soluble ¹⁾						
Cd	(3.43	±	0.17)	18.4	±	0.4
Co	(5.3	±	0.7)	6.5	±	0.4
Cr	307	±	13	174	±	7
Cu	707	±	9	831	±	16
Hg	(1.99	±	0.08)	8.39	±	0.25
Mn	(145	±	7)	298	±	9
Ni	251	±	6	65.0	±	3.0
Pb	282	±	9	583	±	17
Zn	2140	±	50	3040	±	60

Values in brackets are not certified.

Availability: Glass bottles containing about 50 g of powdered material for BCR-146R, 40 g for BCR-145R.

¹⁾ Details of the analytical procedure to obtain the aqua regia soluble content of the elements are given in the certification report.

Substance	ERM-CC580		
	Estuarine sediment (mg/kg)		
Total Hg	132	±	3
CH ₃ Hg ⁺	0.075	±	0.004

Availability: Glass bottles containing about 40 g powder.

Substance	BCR-038 Fly ash from pulverised coal (mg/kg)		
As	48.0	±	2.3
Cd	4.6	±	0.3
Co	53.8	±	1.9
Cr	192	±	10
Cu	176	±	9
Fe	33.8×10^3	±	0.7×10^3
Hg	2.10	±	0.15
Mn	479	±	16
Na	3.74×10^3	±	0.15×10^3
Pb	262	±	11
Zn	581	±	29

Availability: BCR-038 in ampoules containing about 5 g.

Substance	BCR-176R Fly ash (mg/kg)		
As	54	±	5
Cd	226	±	19
Co	26.7	±	1.6
Cr	810	±	70
Cu	1050	±	70
Fe	13100	±	500
Hg	(1.60	±	0.23)
Mn	(730	±	50)
Ni	117	±	6
Pb	5000	±	500
Sb	850	±	50
Se	18.3	±	1.9
Tl	1.32	±	0.21
V	(35	±	6)
Zn	16800	±	400

Values in brackets are not certified.

The report gives additional indicative values for Ag, Au, Ba, Br, Ce, Cs, Eu, Hf, La, Na, Rb, Sc, Ta, Th and W.

Availability: Amber glass bottles containing about 40 g of powdered material.

Substance	BCR-723 Trace elements in road dust (µg/kg)		
Pd	6.1	±	1.9
Pt	81.3	±	2.5
Rh	12.8	±	1.3

Availability: Brown glass bottles with screw cap containing approximately 25 g of powder.

Substance	ERM-CZ120 Elements in fine dust (PM ₁₀ -like) (mg/kg)		
As	7.1	±	0.7
Cd	0.90	±	0.22
Pb	113	±	17
Ni	58	±	7

Availability: Vial containing approximately 0.5 g of fine dust.

Substance	BCR-596 Trapa natans (Aquatic plant) (mg/kg)		
Al Cd Cr Cu Hg Mn Pb Zn	36.3	±	1.7

Availability: CRMs are provided in units of 25 g.

Substance	BCR-129 Hay powder (g/kg)			BCR-402 White clover (mg/kg)		
As	6.40	±	0.10	0.093	±	0.010
Ca				0.178	±	0.008
Co	0.167×10^{-3}	±	0.024×10^{-3}			
I	33.8	±	0.8			
K	1.45	±	0.04			
Mg	37.2	±	0.5	6.93	±	0.19
Mo	2.36	±	0.07			
N	3.16	±	0.04			
P	32.1×10^{-3}	±	1.7×10^{-3}	6.70	±	0.25
S	34.2	±	0.4			
Se						
Zn						
Kjeldahl-N						

Availability: CRMs are provided in powder form in bottles containing approximately for BCR-129 30 g, BCR-402 25 g.

Note: BCR-402 was produced from white clover grown on a ground specially rich in selenium. This explains the high content of this element.

Substance	ERM-CD281 Rye grass (mg/kg)		
As	0.042	±	0.010
B	5.5	±	0.5
Ca		(6.3 g/kg)	
Cd	0.120	±	0.007
Cr	24.8	±	1.3
Cu	10.2	±	0.5
Fe		(0.18 g/kg)	
Hg	0.0164	±	0.0022
K		(34 g/kg)	
Mg		(1.6 g/kg)	
Mn	82	±	4
Mo	2.22	±	0.12
Na		(4.0 g/kg)	
Ni	15.2	±	0.6
P		(2.8 g/kg)	
Pb	1.67	±	0.11
S		(3.4 g/kg)	
Sb	0.042	±	0.007
Se	0.023	±	0.004
Si		(1.3 g/kg)	
Sn	0.062	±	0.011
Zn	30.5	±	1.1

Values in brackets are not certified.

Availability: Amber glass vial containing approximately 10 g.

Substance	BCR-414 Plankton (mg/kg)		
As	6.82	±	0.28
Cd	0.383	±	0.014
Co	(1.43	±	0.06)
Cr	23.8	±	1.2
Cu	29.5	±	1.3
Fe	(1.85	±	0.19 g/kg)
Hg	0.276	±	0.018
K	(7.55	±	0.17 g/kg)
Mn	299	±	13
Mo	(1.35	±	0.20)
Ni	18.8	±	0.8
Pb	3.97	±	0.19
Sc	(0.54	±	0.02)
Se	1.75	±	0.10
Sr	(261	±	25)
V	8.10	±	0.18
Zn	111.6	±	2.5

Values in brackets are not certified.

Availability: CRM is provided in powder form in bottles containing approximately 5 g.

Substance	BCR-482 Lichen (mg/kg)			ERM-CD200 Bladderwrack (<i>Fucus vesiculosus</i>) (mg/kg)		
Al	1103	±	24			
As	0.85	±	0.07	55	±	4
Cd	0.56	±	0.02	0.95	±	0.06
Cr	4.12	±	0.15			
Cu	7.03	±	0.19	1.71	±	0.18
Hg	0.48	±	0.02	0.0186	±	0.0016
Ni	2.47	±	0.07			
Pb	40.9	±	1.4	0.51	±	0.06
Se				0.088	±	0.010
Zn	100.6	±	2.2	25.3	±	1.7

Availability: BCR-482 is provided in powder form in bottles containing approximately 15 g.

ERM-CD200 is provided in powder form in bottles containing approximately 5 g.

Substance	BCR-670 Lemna minor (Aquatic plant) (duck weed) (µg/kg)		
As	(1980	±	190)
Cd	(75.5	±	2.5)
Ce	990	±	40
Cr	(2050	±	100
Cs	(77	±	10)
Cu	(1820	±	300)
Dy	79	±	7
Er	44.0	±	2.8
Eu	23.2	±	1.5
Gd	98	±	8
Ho	15.8	±	1.8
La	487	±	20
Lu	6.3	±	0.5
Mo	(560	±	70)
Nd	473	±	15
Pb	(2060	±	120)
Pr	121	±	6
Sc	191	±	11
Sm	94	±	7
Tb	14.0	±	1.1
Th	159	±	18
Tm	5.7	±	0.7
U	82	±	8
Y	460	±	60
Yb	40	±	4
Zn	(24000	±	2100)

Values in brackets are not certified.

Availability: Glass bottles containing about 10 g of powdered material.

Substance	ERM-CE278k Mussel tissue (mg/kg)		
As	6.7	±	0.4
Ag	(0.044	±	0.016)
Cd	0.336	±	0.025
Cr	0.73	±	0.22
Cu	5.98	±	0.27
Fe	161	±	8
Hg	0.071	±	0.007
Mn	4.88	±	0.24
Ni	0.69	±	0.15
Pb	2.18	±	0.18
Rb	2.46	±	0.16
Se	1.62	±	0.12
Sr	19.0	±	1.2
Zn	71	±	4

Values in brackets are not certified.

Availability: CRMs are provided in powder form in bottles containing approximately 8 g.

Substance	BCR-668 Mussel tissue (µg/kg)		
As	(7100	±	500)
Cd	(275	±	11)
Ce	89	±	7
Cr	(370	±	60)
Cs	(13.8	±	1.5)
Dy	8.9	±	0.6
Er	4.5	±	0.5
Eu	2.79	±	0.16
Gd	13.0	±	0.6
Ho	(1.8	±	0.6)
La	80	±	6
Lu	0.389	±	0.024
Mo	(1990	±	150)
Nd	54	±	4
Pr	12.3	±	1.1
Sc	(8.5	±	1.8)
Sm	11.2	±	0.8
Tb	1.62	±	0.12
Th	10.7	±	1.2
Tm	0.48	±	0.08
U	56	±	5
Y	59	±	5
Yb	(2.8	±	0.5)
Zn	(70700	±	400)

Values in brackets are not certified.

Availability: Glass bottles containing about 10 g of powdered material.

Substance	ERM-CE464 Tuna fish (mg/kg)		
Total Hg	5.24	±	0.10
CH ₃ Hg ⁺	5.50	±	0.17

Availability: Glass bottles containing about 15 g.

Substance	BCR-505 Trace elements in estuarine water (nmol/kg)			BCR-579 Coastal sea-water (ng/kg)		
Cd	0.80	±	0.04			
Co	(0.99	±	0.26)			
Cu	29.4	±	1.5			
Fe	(19	±	4)			
Hg						
Ni	24.1	±	2.0	1.9	±	0.5
Pb	(0.24	±	0.14)			
Zn	172	±	11			

Values in brackets are not certified.

Availability: BCR-505 is provided in 1 L polyethylene bottles and BCR-579 in 1 L glass bottles.

Substance	ERM-CA408 Simulated rainwater (low contents) (mg/L)		
Mass concentration:			
Ammonium	0.910	±	0.028
Cl	1.96	±	0.07
Fluoride	0.194	±	0.008
Mg	0.145	±	0.022
NO ₃	2.01	±	0.09
Ortho-phosphate	1.00	±	0.05
SO ₄	1.46	±	0.04
Electrochemical property:			
Conductivity (20 °C)	18.7	±	1.8 µS/cm
pH (20 °C)	6.3	±	0.6

Availability: ERM-CA408 is provided in units of about 95 mL in flame-sealed ampoules.

Substance	BCR-479 Freshwater (low contents)	BCR-480 Freshwater (high contents)
Nitrate		
As amount of substance content	214 ± 4 µmol/kg	885 ± 13 µmol/kg
As mass fraction	13.3 ± 0.3 mg/kg	54.9 ± 0.8 mg/kg

Availability: Units of about 100 mL in white glass ampoules.

Substance	BCR-611 Bromide in ground water based on IC-measurements (low contents) (µg/kg)	BCR-612 Bromide in ground water based on IC-measurements (high contents) (µg/kg)
Br	93 ± 4	252 ± 10

Availability: Set of four brown glass ampoules of 25 mL.

Substance	ERM-CA616 Ground water (high carbonate content) (mg/L)
Mass concentration:	
Ca	42.6 ± 1.4
Cl	44.6 ± 10.9
K	5.79 ± 0.15
Mg	10.1 ± 0.3
Na	27.9 ± 0.8
Ortho-phosphate	2.24 ± 0.10
Electrochemical property:	
Conductivity (20 °C)	426 ± 5 µS/cm
pH (20 °C)	7.12 ± 0.18

Availability: ERM-CA616 consists of about 95 mL natural groundwater in a flame-sealed ampoule

Substance	ERM-CA615 Groundwater			
As	9.9	±	0.7	µg/L
Cd	0.106	±	0.011	µg/L
Fe	5.11	±	0.26	mg/L
Hg	0.037	±	0.004	µg/L
Mn	107	±	5	µg/L
Ni	25.3	±	1.1	µg/L
Pb	7.1	±	0.6	µg/L

Availability: One unit consists of about 95 mL natural groundwater in a flame-sealed ampoule.

Substance	ERM-CA713 Wastewater µg/L		
As	10.8	±	0.3
Cd	5.09	±	0.20
Cr	20.9	±	1.3
Cu	101	±	7
Fe	445	±	27
Hg	1.84	±	0.11
Mn	95	±	4
Ni	50.3	±	1.4
Pb	49.7	±	1.7
Se	4.9	±	1.1

Availability: ampoule containing approximately 100 mL of wastewater effluent acidified with HNO₃ to about pH 2.

Substance	ERM-CA400 Seawater		
Hg mass concentration	16.8	±	1.1
Hg mass fraction	16.4	±	1.0
True density at 20 °C	1.0226	±	0.0003

Availability: One unit consists of 3 vials, each containing 100 mL acidified coastal surface seawater.

1.2.2 CERTIFIED FOR THE EXTRACTABLE ELEMENT CONTENT AND SPECIES

Substance		BCR-483 Sewage sludge amended soil (mg/kg)			BCR-484 Sewage sludge amended (terra rossa) soil (mg/kg)			BCR-700 Organic-rich soil (mg/kg)		
<u>EDTA:</u>	Cd	24.3	±	1.3	0.509	±	0.030	65.2	±	3.5
	Cr	28.6	±	2.6				10.1	±	0.9
	Cu	215	±	11	88	±	4	89.4	±	2.8
	Ni	28.7	±	1.7	1.39	±	0.11	53.2	±	2.8
	Pb	229	±	8	47.9	±	2.6	103	±	5
	Zn	612	±	20	152	±	7	510	±	17
<u>Acetic acid:</u>	Cd	18.3	±	0.6	0.48	±	0.04	67.5	±	2.8
	Cr	18.7	±	1.0				19.0	±	1.1
	Cu	33.5	±	1.6	33.9	±	1.4	36.3	±	1.6
	Ni	25.8	±	1.0	1.69	±	0.16	99.0	±	5.1
	Pb	2.10	±	0.25	1.17	±	0.16	4.85	±	0.38
	Zn	620	±	24	193	±	7	719	±	24
<u>Calcium chloride extractable content</u>	Cd	(0.45	±	0.05)			(< 0.08)			
	Cr	(0.35	±	0.09)			(< 0.09)			
	Cu	(1.2	±	0.4)			(0.67	±	0.29)	
	Ni	(1.4	±	0.2)			(< 0.05)			
	Pb	(< 0.06)					(< 0.06)			
	Zn	(8.3	±	0.7)			(0.31	±	0.17)	
<u>Sodium nitrate extractable content</u>	Cd	(0.08	±	0.03)			(< 0.05)			
	Cr	(0.30	±	0.07)			(< 0.03)			
	Cu	(0.89	±	0.22)			(0.48	±	0.15)	
	Ni	(0.65	±	0.07)			(0.023	±	0.005)	
	Pb	(< 0.03)					(< 0.06)			
	Zn	(2.7	±	0.8)			(0.09	±	0.04)	
<u>Ammonium nitrate extractable content</u>	Cd	(0.26	±	0.05)			(0.003	±	0.002)	
	Cr	(0.27	±	0.10)			(< 0.06)			
	Cu	(1.2	±	0.3)			(1.1	±	0.4)	
	Ni	(1.1	±	0.3)			(0.033	±	0.017)	
	Pb	(0.020	±	0.013)			(< 0.06)			
	Zn	(6.5	±	0.9)			(0.17	±	0.05)	

Availability: BCR-483 and -484 are provided in glass bottles containing about 70 g of powder. BCR-700 is provided in glass bottles containing about 40 g of powder.

Substance		BCR-684 River sediment (mg/kg)		
NaOH-extractable P		550	±	21
HCl-extractable P		536	±	28
Inorganic P		1113	±	24
Organic P		209	±	9
Conc. HCl-extract. P		1373	±	35

Availability: Glass bottles containing about 35 g of powdered material.

Substance	BCR-701 Lake sediment (mg/kg)		
	<u>Extractable mass fraction based on dry mass</u>		
<u>Step 1:</u>	Cd 7.3 ± 0.4 Cr 2.26 ± 0.16 Cu 49.3 ± 1.7 Ni 15.4 ± 0.9 Pb 3.18 ± 0.21 Zn 205 ± 6		
<u>Step 2:</u>	Cd 3.77 ± 0.28 Cr 45.7 ± 2.0 Cu 124 ± 3 Ni 26.6 ± 1.3 Pb 126 ± 3 Zn 114 ± 5		
<u>Step 3:</u>	Cd 0.27 ± 0.06 Cr 143 ± 7 Cu 55 ± 4 Ni 15.3 ± 0.9 Pb 9.3 ± 2.0 Zn 46 ± 4		
	<u>Mass fraction based on dry mass</u>		
Cd	(0.13 ± 0.08) Cr (62.5 ± 7.4) Cu (38.5 ± 11.2) Ni (41.4 ± 4.0) Pb (11.0 ± 5.2) Zn (95 ± 13)		

Values in brackets are not certified.

Availability: Glass bottles containing about 20 g of powdered material.

Substance	BCR-462 Coastal sediment (µg/kg)			BCR-646 Freshwater sediment (µg/kg)		
	54	±	15	480	±	80
Tributyltin (TBT)	68	±	12	770	±	90
Dibutyltin (DBT)				610	±	120
Monobutyltin (MBT)				29	±	11
Triphenyltin (TPhT)				36	±	8
Diphenyltin (DPhT)				69	±	18
Monophenyltin (MPhT)						

Availability: Glass bottle containing about 25 g of powder for BCR-462 and 40 g of powder for BCR-646.

Substance	ERM-CC580 Estuarine sediment (mg/kg)		
Total Hg CH_3Hg^+	132 ± 3 0.075 ± 0.004		

Availability: Glass bottles containing about 40 g powder.

	BCR-545 Welding dust loaded on a filter (g/kg)		
Cr (VI) total leachable Cr	40.2 39.5	±	0.6 1.3

Availability: Glass fibre filter loaded with welding dust containing about 100 µg Cr (VI).

Substance	ERM-CE477 Mussel tissue (mg/kg)		
Tributyltin (TBT)	2.20	±	0.19
Dibutyltin (DBT)	1.54	±	0.12
Monobutyltin (MBT)	1.50	±	0.28

Availability: Glass bottle containing about 14 g of powder.

Substance	ERM-CE464 Tuna fish (mg/kg)		
Total Hg	5.24	±	0.10
CH ₃ Hg ⁺	5.50	±	0.17

Availability: Glass bottles containing about 15 g powder.

Substance	BCR-627 Tuna fish tissue			ERM-BC211 Rice				
Total As	4.8	±	0.3	mg/kg	260	±	13	µg/kg
Dimethylarsinic acid	2.0	±	0.3	µmol/kg	119	±	13	µg/kg
Sum of arsenite and arsenate					124	±	11	µg/kg
Arsenobetaine	52	±	3	µmol/kg				

Availability: BCR-627: Glass bottles containing about 10 g powder

ERM-BC211: Vial containing about 10 g of powder

1.2.3 CERTIFIED FOR ORGANIC POLLUTANTS

Substance	ERM-CZ100 Fine dust (PM ₁₀ -like) (mg/kg)		
Benz[a]anthracene	0.91	±	0.07
Benzo[a]pyrene	0.72	±	0.05
Benzo[b]fluoranthene	1.42	±	0.14
Benzo[j]fluoranthene	0.75	±	0.14
Benzo[k]fluoranthene	0.67	±	0.06
Dibenzo[a,h]anthracene	0.18	±	0.04
Indeno[1,2,3-c,d]pyrene	1.07	±	0.10
Sum of Benzo[b]fluoranthene, benzo[k]fluoranthene and benzo[j]fluoranthene	2.84	±	0.21

Availability: Vial containing about 0.5 g of fine dust.

Substance	BCR-524 Contaminated industrial soil (mg/kg)		
Pyrene	173	±	11
Benz[a]anthracene	22.5	±	1.8
Benzo[a]pyrene	8.6	±	0.5
Benzo[e]pyrene	10.6	±	1.4
Benzo[b]fluoranthene	13.5	±	1.6
Benzo[k]fluoranthene	6.2	±	0.6
Benzo[b]naphtho[2,1-d]-thiophene	3.8	±	0.6
Indeno[1,2,3-cd]pyrene	5.1	±	0.4
Pentachlorophenol	0.034	±	0.005

Availability: Glass bottle containing about 40 g of powder.

Substance	BCR-535 Freshwater harbour sediment (mg/kg)		
Pyrene	2.52	±	0.18
Benz[a]anthracene	1.54	±	0.10
Benzo[a]pyrene	1.16	±	0.10
Benzo[e]pyrene	1.86	±	0.13
Benzo[b]fluoranthene	2.29	±	0.15
Benzo[k]fluoranthene	1.09	±	0.15
Indeno[1,2,3-cd]pyrene	1.56	±	0.14

Availability: Glass bottle containing about 40 g of powder.

Polychlorinated biphenyls (IUPAC No.)	BCR-481 Industrial soil (mg/kg)			BCR-536 Freshwater harbour sediment (µg/kg)		
28				44	±	5
52				38	±	4
101	37	±	3	44	±	4
105				3.5	±	0.6
118	9.4	±	0.7	28	±	3
128	9.1	±	0.8	5.4	±	1.2
138				27	±	4
149	97	±	7	49	±	4
153	137	±	7	50	±	4
156	7.0	±	0.5	3.0	±	0.4
163				17	±	3
170	52	±	4	13.4	±	1.4
180	124	±	6	22	±	2

Availability: BCR-481 is provided in brown glass bottles with a polyethylene insert containing approximately 25 g of soil. BCR-536 is provided in a glass bottle containing about 40 g of powder.

Substance	BCR-529 Industrial (sandy) soil				BCR-530 Industrial (clay) soil			
3,4-dichlorophenol	0.23	±	0.04	mg/kg	6.0	±	0.5	mg/kg
2,4,5-trichlorophenol	1.51	±	0.10	mg/kg	40	±	7	mg/kg
Pentachlorophenol	0.23	±	0.04	mg/kg	0.47	±	0.09	mg/kg
2,3,7,8 - TCDD (D48)	4.5	±	0.6	µg/kg				
1,2,3,7,8 - PeCDD (D54)	0.44	±	0.05	µg/kg				
1,2,3,4,7,8 - HxCDD (D66)	1.22	±	0.21	µg/kg				
1,2,3,6,7,8 - HxCDD (D67)	5.4	±	0.9	µg/kg	0.061	±	0.011	µg/kg
1,2,3,7,8,9 - HxCDD (D70)	3.0	±	0.4	µg/kg	0.0218	±	0.0029	µg/kg
2,3,7,8 - TCDF (F83)	0.078	±	0.013	µg/kg				
1,2,3,7,8 - PeCDF (F94)	0.145	±	0.028	µg/kg	0.24	±	0.04	µg/kg
2,3,4,7,8 - PeCDF (F114)	0.36	±	0.07	µg/kg	0.62	±	0.07	µg/kg
1,2,3,4,7,8 - HxCDF (F118)	3.4	±	0.5	µg/kg	0.321	±	0.016	µg/kg
1,2,3,6,7,8 - HxCDF (F121)	1.09	±	0.15	µg/kg	0.186	±	0.023	µg/kg
1,2,3,7,8,9 - HxCDF (F124)	0.022	±	0.010	µg/kg				
2,3,4,6,7,8 - HxCDF (F130)	0.37	±	0.05	µg/kg	0.126	±	0.012	µg/kg

Availability: Amber glass bottles containing about 50 g of dried soil.

Substance	BCR-677 Sewage sludge (ng/kg)		
2,3,7,8 - T ₄ CDD (D48)	1.51	±	0.16
1,2,3,7,8 - P ₅ CDD (D54)	4.1	±	0.9
1,2,3,6,7,8 - H ₆ CDD (D67)	235	±	16
1,2,3,7,8,9 - H ₆ CDD (D70)	79	±	7
1,2,3,4,6,7,8 - H ₇ CDD (D73)	3.5×10^3	±	0.4×10^3
O ₈ CDD (D75)	12.7×10^3	±	0.8×10^3
2,3,7,8 - T ₄ CDF (F83)	45	±	4
1,2,3,7,8 - P ₅ CDF (F94)	24.8	±	1.6
2,3,4,7,8 - P ₅ CDF (F114)	16.9	±	1.5
1,2,3,4,7,8 - H ₆ CDF (F118)	14.5	±	1.6
1,2,3,6,7,8 - H ₆ CDF (F121)	6.1	±	0.8
1,2,3,7,8,9 - H ₆ CDF (F124)	0.84	±	0.29
2,3,4,6,7,8 - H ₆ CDF (F130)	5.6	±	0.6
1,2,3,4,6,7,8 - H ₇ CDF (F131)	62	±	3
1,2,3,4,7,8,9 - H ₇ CDF (F134)	6.3	±	0.8
O ₈ CDF (F135)	177	±	7

Availability: BCR-677 consists of approximately 40 g of dried sewage sludge in amber glass bottles.

Substance	BCR-490 Fly ash (µg/kg)		
2,3,7,8 - T ₄ CDD (D48)	0.169	±	0.012
1,2,3,7,8 - P ₅ CDD (D54)	0.67	±	0.04
1,2,3,4,7,8 - H ₆ CDD (D66)	0.95	±	0.11
1,2,3,6,7,8 - H ₆ CDD (D67)	4.8	±	0.4
1,2,3,7,8,9 - H ₆ CDD (D70)	2.84	±	0.17
2,3,7,8 - T ₄ CDF (F83)	0.90	±	0.05
1,2,3,7,8 - P ₅ CDF (F94)	1.71	±	0.12
2,3,4,7,8 - P ₅ CDF (F114)	1.85	±	0.11
1,2,3,4,7,8 - H ₆ CDF (F118)	2.37	±	0.12
1,2,3,6,7,8 - H ₆ CDF (F121)	2.64	±	0.14
1,2,3,7,8,9 - H ₆ CDF (F124)	0.34	±	0.05
2,3,4,6,7,8 - H ₆ CDF (F130)	2.47	±	0.17

Availability: BCR-490 consists of approximately 30 g of fly ash in amber glass bottles.

Substance	BCR-615 Fly ash (low level) (ng/kg)		
2,3,7,8 - T ₄ CDD (D48)	27	±	5
1,2,3,7,8 - P ₅ CDD (D54)	92	±	12
1,2,3,4,7,8 - H ₆ CDD (D66)	74	±	12
1,2,3,6,7,8 - H ₆ CDD (D67)	103	±	13
1,2,3,7,8,9 - H ₆ CDD (D70)	108	±	16
1,2,3,4,6,7,8 - H ₇ CDD (D73)	0.87×10^3	±	0.13×10^3
O ₈ CDD (D75)	1.75×10^3	±	0.20×10^3
2,3,7,8 - T ₄ CDF (F83)	86	±	28
1,2,3,7,8 - P ₅ CDF (F94)	176	±	26
2,3,4,7,8 - P ₅ CDF (F114)	125	±	20
1,2,3,4,7,8 - H ₆ CDF (F118)	203	±	21
1,2,3,6,7,8 - H ₆ CDF (F121)	204	±	23
1,2,3,7,8,9 - H ₆ CDF (F124)	13.3	±	2.0
2,3,4,6,7,8 - H ₆ CDF (F130)	130	±	15
1,2,3,4,6,7,8 - H ₇ CDF (F131)	0.75×10^3	±	0.09×10^3
1,2,3,4,7,8,9 - H ₇ CDF (F134)	61	±	6
O ₈ CDF (F135)	0.29×10^3	±	0.04×10^3

Availability: BCR-615 consists of approximately 50 g of dried fly ash in amber glass bottles.

Substance	BCR-683 Beech wood (mg/kg)		
Benz[a]anthracene	6.5	±	0.7
Benzo[a]pyrene	3.4	±	0.4
Benzo[e]pyrene	9.3	±	1.0
Benzo[b]fluoranthene	5.8	±	0.6
Benzo[k]fluoranthene	2.58	±	0.29
Pentachlorophenol	3.6	±	0.5

Availability: Glass bottle containing about 60 g of powder.

Polychlorinated biphenyls (IUPAC No.)	BCR-682 Mussel tissue (µg/kg)			BCR-718 Canned fresh herring (µg/kg)		
	28	0.30	±	0.07	0.41	±
52	0.78	±		1.00	±	0.04
101				2.12	±	0.06
105				0.63	±	0.06
118	2.6	±	0.3	1.78	±	0.07
128				0.62	±	0.101
138	4.6	±	0.8	2.97	±	0.11
149	5.7	±	0.9	2.58	±	0.11
153	9.2	±	0.8	4.62	±	0.10
170	0.17	±	0.05	0.350	±	0.026
180	0.77	±	0.07	0.795	±	0.027

Availability: BCR-682 and BCR-718 are provided in sealed tin cans containing approximately 70 g fresh mussel tissue.

Polychlorinated biphenyls (IUPAC No.)	BCR-420 Waste mineral oil (low level) (mg/kg)			BCR-449 Waste mineral oil (high level) (mg/kg)		
	28	0.61	±	0.06	0.80	±
52				31.4	±	1.8
101	1.45	±		57.2	±	1.9
105				17.4	±	1.0
118	1.69	±	0.14	46.6	±	2.4
128				12.5	±	0.7
153	0.92	±	0.06	39.0	±	1.7
156				6.9	±	0.5
170				6.6	±	0.6
180	0.195	±	0.017	10.4	±	0.4

Availability: BCR-420 is provided in units of about 7.5 g in glass ampoules and BCR-449 in units of about 50 g in glass ampoules.

Substance	IRMM-427 Fish tissue (Pike-Perch) (ng/g)		
Linear perfluorooctane sulfonate (L-PFOS)	16.0	±	1.7
Perfluorodecanoic acid (PFDA)	1.28	±	0.17
Perfluoroundecanoic acid (PFUnDA)	0.74	±	0.20
Perfluorododecanoic acid (PFDoDA)	0.97	±	0.21

Availability: Glass jars containing 35 g of fish paste.

Substance	IRMM-428 Water (ng/L)		
Perfluorobutane sulfonate (PFBS)	5.5	±	1.4
Perfluorohexane sulfonate (PFHxS)	3.6	±	1.0
Linear perfluoroctane sulfonate (L-PFOS)	9.6	±	1.7
Perfluoropentanoic acid (PFPeA)	4.0	±	1.0
Perfluorohexanoic acid (PFHxA)	7.4	±	1.0
Perfluoroheptanoic acid (PFHpA)	3.7	±	0.7

Availability: High density polyethylene bottles containing 410 mL of water.

Substance	ERM-CA100 Surface water			
Naphthalene	1.21	±	0.13	µg/L
Anthracene	91	±	11	ng/L
Fluoranthene	104	±	11	ng/L
Benzo[b]fluoranthene	32	±	9	ng/L
Benzo[k]fluoranthene	38	±	9	ng/L
Benzo[a]pyrene	42	±	8	ng/L
Indeno[1,2,3-cd]pyrene	29	±	7	ng/L

Availability: The CRM is available as unit kit that includes a plastic container with at least 1000 mL of surface water, an amber glass ampoule with at least 24 mL humic acids solution in water and an amber glass ampoule with at least 2 mL PAH solution in acetonitrile.

Substance	ERM-CE100 Fish tissue (µg/kg)		
Hexachlorobenzene	120	±	8
Hexachlorobutadiene	36	±	4

Availability: The CRM is available in glass jars with twist-off lids containing approximately 40 g of fish paste.

1.2.4 OTHERS

Parameter	IRMM443-1 (EUROSOIL 1)	IRMM443-2 (EUROSOIL 2)	IRMM443-3 (EUROSOIL 3)
K _f of atrazine ⁽¹⁾	7.0 ± 1.5	2.7 ± 0.7	2.4 ± 0.7
1/n of atrazine ⁽¹⁾	0.91 ± 0.11	0.93 ± 0.12	0.91 ± 0.13
K _f of 2,4-D ⁽¹⁾	2.5 ± 1.0	0.99 ± 0.30	1.31 ± 0.28
1/n of 2,4-D ⁽¹⁾	0.9 ± 0.4	0.96 ± 0.15	0.93 ± 0.15
K _f of lindane ⁽¹⁾		48 ± 11	
1/n of lindane ⁽¹⁾		0.98 ± 0.15	
pH in water ⁽²⁾	6.21 ± 0.30	8.1 ± 0.9	6.2 ± 0.4
pH in 0.01M CaCl ₂ ⁽²⁾	5.65 ± 0.24	7.5 ± 0.8	5.5 ± 0.4

Parameter	IRMM443-4 (EUROSOIL 4)	IRMM443-5 (EUROSOIL 5)	IRMM443-7 (EUROSOIL 7)
K _f of atrazine ⁽¹⁾	0.7 ± 0.4	13 ± 6	4.8 ± 1.1
1/n of atrazine ⁽¹⁾	0.87 ± 0.22	0.9 ± 0.4	0.92 ± 0.15
K _f of 2,4-D ⁽¹⁾	0.39 ± 0.21	18 ± 7	8.2 ± 1.8
1/n of 2,4-D ⁽¹⁾	0.9 ± 0.4	0.9 ± 0.4	0.88 ± 0.15
K _f of lindane ⁽¹⁾	8.3 ± 2.2		
1/n of lindane ⁽¹⁾	0.96 ± 0.12		
pH in water ⁽²⁾	7.5 ± 0.7	4.1 ± 1.5	5.1 ± 0.8
pH in 0.01M CaCl ₂ ⁽²⁾	6.8 ± 0.6	3.1 ± 1.1	4.3 ± 0.7

(1) Determination according OECD Testguideline 106.

(2) Measurement based on ISO Standard 10390.

Uncertainty express as estimated expanded uncertainty as defined in the Guide to the Expression of Uncertainty in Measurement (GUM).
Availability: Brown glass bottles with 200 g of air-dried fine soil (< 2 mm).

2 MATERIALS RELATED TO THE ANALYSIS OF FOOD AND FEEDING STUFF

2.1 PURE MATERIALS AND SYNTHETIC MIXTURES

BCR-123 Ethanol						
Parameter	Ethanol H		Ethanol M		Ethanol L	
(D/H) _I	109.65×10^{-6}	$\pm 0.20 \times 10^{-6}$	101.69×10^{-6}	$\pm 0.17 \times 10^{-6}$	90.30×10^{-6}	$\pm 0.18 \times 10^{-6}$
(D/H) _{II}	119.76×10^{-6}	$\pm 0.25 \times 10^{-6}$	130.94×10^{-6}	$\pm 0.21 \times 10^{-6}$	122.20×10^{-6}	$\pm 0.4 \times 10^{-6}$
R	2.184	± 0.005	2.575	± 0.006	2.708	± 0.009

Availability: Units of 3 sealed NMR tubes containing respectively H-, M-, and L-ethanols, to which the tetramethylurea internal standard and the C₆F₆ lock substance are added. 10 mm (BCR-123A) or 15 mm (BCR-123B) O.D. NRM tubes can be supplied.

Parameter	Unit	BCR-656 (96% ethanol)		
(D/H) _I by ² H-NMR	ppm	102.84	\pm	0.20
(D/H) _{II} by ² H-NMR	ppm	132.07	\pm	0.30
R by ² H-NMR		2.570	\pm	0.005
$\delta^{13}\text{C}_{\text{VPDB}}$ by IRMS	\textperthousand	-26.91	\pm	0.07
Alcoholic grade t _D	w/w %	(94)		

Value in brackets is not certified.

Availability: BCR-656: Units of 25 mL of 96 % vol. neutral ethanol from wine in glass bottle.

Parameter	Unit	BCR-657 (Sugar)	BCR-660 (Ethanol in water)
(D/H) _I by ² H-NMR	ppm		102.90 \pm 0.16
(D/H) _{II} by ² H-NMR	ppm		131.95 \pm 0.23
R by ² H-NMR			2.567 \pm 0.005
$\delta^{13}\text{C}_{\text{VPDB}}$ by IRMS	\textperthousand	-10.76 \pm 0.04	-26.72 \pm 0.09
(D/H) _w of water (IRMS)	ppm		148.68 \pm 0.14
Alcoholic grade t _D	w/w %		11.96 \pm 0.06 ¹⁾

1) in v/v %

Availability: BCR-657: Units of approx. 1 g of dry glucose in a sealed amber vial;
BCR-660: Units of 370 mL of aqueous ethanol solution in glass bottle.

Tetramethylurea (STA-003m)

Tetramethylurea (TMU) which is used as Internal Standard in routine SNIF-NMR analysis is available in 500 mL quantities. The D/H nominal value of tetramethylurea batches is determined by multiple calibration at 61.45/400 MHz, 61.4/400 MHz and 45.05/400 MHz for deuterium and given in an accompanying analytical report.

Substance	BCR-423 (RM) Aflatoxin M ₁ in chloroform ($\mu\text{g/mL}$)
Aflatoxin M ₁	(9.93)

Value in brackets is not certified.

Availability: Sealed ampoules containing about 2.5 mL.

Compound	ERM-AC626 Arsenobetaine in water		
Arsenobetaine	250.0	±	2.5

Availability: ERM-AC626 is available in ampoules containing 1 mL.

Compound	BCR-663 Saxitoxin in acetic acid		
Saxitoxin·2HCl	9.8	±	1.2

Availability: BCR-663 is available in ampoules containing 1 mL.

Substance	ERM-AC699 Zearalenone in acetonitrile		
ZON	9.95	±	0.30

Availability: ERM-AC699 is supplied in ampoules filled and sealed under nitrogen in amounts of 4 mL.

Substance	ERM-AC057 Aflatoxin B1 in acetonitrile		
Aflatoxin B1	Mass fraction (µg/g) 3.79	± 0.11	Mass concentration at 20 °C (µg/mL) (2.97 ± 0.09)

Values in brackets are not certified.

Availability: ERM-AC057 is supplied in amber glass ampoules filled with 4 mL.

Substance	ERM-AC058 Aflatoxin B2 in acetonitrile		
Aflatoxin B2	Mass fraction (µg/g) 3.80	± 0.08	Mass concentration at 20 °C (µg/mL) (2.98 ± 0.06)

Values in brackets are not certified.

Availability: ERM-AC058 is supplied in amber glass ampoules filled with 4 mL.

Substance	ERM-AC059 Aflatoxin G1 in acetonitrile		
Aflatoxin G1	Mass fraction (µg/g) 3.78	± 0.13	Mass concentration at 20 °C (µg/mL) (2.96 ± 0.10)

Values in brackets are not certified.

Availability: ERM-AC059 is supplied in amber glass ampoules filled with 4 mL.

Substance	ERM-AC060 Aflatoxin G2 in acetonitrile		
Aflatoxin G2	Mass fraction (µg/g) 3.80	± 0.07	Mass concentration at 20 °C (µg/mL) (2.98 ± 0.06)

Values in brackets are not certified.

Availability: ERM-AC060 is supplied in amber glass ampoules filled with 4 mL.

Substance	IRMM-315 4-Deoxynivalenol in acetonitrile		
4-Deoxynivalenol	25.1	<u>Mass fraction</u> ($\mu\text{g/g}$) \pm	1.2 (19.7 \pm 0.9)

Values in brackets are not certified.

Availability: IRMM-315 is supplied in amber glass ampoules filled with 4 mL.

Substance	IRMM-316 Nivalenol in acetonitrile		
Nivalenol	24.0	<u>Mass fraction</u> ($\mu\text{g/g}$) \pm	1.1 (18.8 \pm 0.9)

Values in brackets are not certified.

Availability: IRMM-316 is supplied in amber glass ampoules filled with 4 mL.

2.2 MATRIX MATERIALS

2.2.1 CERTIFIED FOR GMO CONTENT

The materials were prepared by quantitative mixing of non-genetically modified powder and genetically modified powder, produced from ground seed with the help of a dry-mixing technique, and are intended for the calibration of methods for the detection of genetically modified food.

CRMs for genetically modified GTS 40-3-2 soya beans (ERM-BF410p)

Five CRMs of dried soya bean powder with different mass fractions of genetically modified (GTS 40-3-2) soya beans were produced.

	Certified value GTS 40-3-2 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF410ap	< 0.09	-
ERM-BF410bp	> 985	-
ERM-BF410cp	1.00	0.10
ERM-BF410dp	10.0	0.6
ERM-BF410ep	100	5

Availability: Vials containing about 1 g of soya bean powder.

CRMs for genetically modified Bt-176 maize (ERM-BF411)

Six CRMs of dried maize powder with different mass fractions of genetically modified (Bt-176) maize were produced.

	Certified value Bt-176 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF411a	< 0.14	-
ERM-BF411b	1.00	0.29
ERM-BF411c	5.0	0.6
ERM-BF411d	10.0	0.8
ERM-BF411e	20.0	1.1
ERM-BF411f	50.0	1.8

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified Bt-11 maize (ERM-BF412k)

Five CRMs of dried maize powder with different mass fractions of genetically modified (Bt-11) maize were produced.

	Certified value Bt-11 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF412ak	< 0.12	-
ERM-BF412bk	> 970	-
ERM-BF412ck	0.99	0.13
ERM-BF412dk	9.9	0.7
ERM-BF412ek	99	4

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified MON 810 maize (ERM-BF413k)

Four CRMs of dried maize powder with different mass fractions of genetically modified (MON 810) maize were produced.

	Certified value MON 810 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF413ak	< 0.9	-
ERM-BF413ck	4.9	1.0
ERM-BF413ek *	19.8	1.5
ERM-BF413gk	99	5

Availability: Vials containing about 1 g of maize powder.

* ERM-BF413ek is also certified for the DNA copy number ratio.

	Certified value MON 810 DNA copy number ratio (%)	Uncertainty (%)
ERM-BF413ek	0.77	0.08

ERM-AD413 DNA fragments per plasmid	
	Number
Fragment of 5' plant-P35S junction DNA/plasmid	1
Fragment of hmg DNA/plasmid	1
	(negligible uncertainty) (negligible uncertainty)
Ratio between the number of 5' plant-P35S junction and hmg fragments in the plasmid by duplex rt-PCR ⁽¹⁾ and simplex rt-PCR ⁽²⁾	<u>Number ratio</u> (1.00 ¹ ± 0.06) (1.04 ² ± 0.06)

Values in brackets are not certified.

Availability: ERM-AD413 is available in vials.

CRMs for genetically modified GA21 maize (ERM-BF414)

Six CRMs of dried maize powder with different mass fractions of genetically modified (GA21) maize were produced.

	Certified value GA21 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF414a	< 0.8	-
ERM-BF414b	1.0	0.8
ERM-BF414c	4.9	1.0
ERM-BF414d	9.9	1.1
ERM-BF414e	17.2	1.2
ERM-BF414f	42.9	1.7

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified NK603 maize (ERM-BF415)

Six CRMs of dried maize powder with different mass fractions of genetically modified (NK603) maize were produced.

	Certified value NK603 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF415a	< 0.4	-
ERM-BF415b	1.0	0.4
ERM-BF415c	4.9	0.5
ERM-BF415d	9.8	0.7
ERM-BF415e *	19.6	0.9
ERM-BF415f	49.1	1.3

Availability: Vials containing about 1 g of maize powder.

* ERM-BF415e is also certified for the DNA copy number ratio.

	Certified value NK603 maize DNA copy number ratio (%)	Uncertainty (%)
ERM-BF415e	0.95	0.11

	ERM-AD415	
	DNA fragments per plasmid	
Fragment of 3' insertion-specific DNA / pIRMM-0086	1	(negligible uncertainty)
Fragment of hmg DNA / pIRMM-0086	1	(negligible uncertainty)

Availability: ERM-AD415 is available in vials.

CRMs for genetically modified MON 863 maize (ERM-BF416)

Four CRMs of dried maize powder with different mass fractions of genetically modified (MON 863) maize were produced.

	Certified value MON 863 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF416a	< 1.0	-
ERM-BF416b	1.0	- 0.3 ; + 1.0
ERM-BF416c	9.8	- 0.7 ; + 1.2
ERM-BF416d	98.5	- 2.2 ; + 2.5

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified MON 863 x MON 810 maize (ERM-BF417)

Four CRMs of dried maize powder with different mass fractions of genetically modified (MON 863 x MON 810) maize were produced.

	Certified value MON 863 x MON 810 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF417a	< 1.0	-
ERM-BF417b	1.0	- 0.2 ; + 1.0
ERM-BF417c	9.8	- 0.7 ; + 1.2
ERM-BF417d	98.5	- 2.0 ; + 2.4

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified 1507 maize (ERM-BF418)

Four CRMs of dried maize powder with different mass fractions of genetically modified (1507) maize were produced.

	Certified value 1507 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF418a	< 0.5	-
ERM-BF418b	1.0	- 0.2 ; + 0.6
ERM-BF418c	9.9	- 0.6 ; + 0.8
ERM-BF418d	98.6	- 1.7 ; + 2.0

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified H7-1 sugar beet (ERM-BF419)

Two CRMs of dried sugar beet powder with different mass fractions of genetically modified (H7-1) sugar beet were produced.

	Certified value H7-1 sugar beet mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF419a	0	0
ERM-BF419b	1000	0

Availability: Vials containing about 1 g of sugar beet powder.

CRMs for genetically modified 3272 maize (ERM-BF420)

Three CRMs of dried maize powder with different mass fractions of genetically modified (3272) maize were produced.

	Certified value 3272 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF420a	< 1.3	-
ERM-BF420b	9.8	1.2
ERM-BF420c	98	8

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified EH92-527-1 potato (ERM-BF421)

Two CRMs of dried potato powder with different mass fractions of genetically modified (EH92-527-1) potato were produced.

	Certified value Number fraction of EH92-527-1 potato / total number of potatoes [%]	Certified property Identity	Uncertainty
ERM-BF421a	0	potato without the EH92-527-1 event	
ERM-BF421b	100	EH92-527-1 potato	Not applicable

Availability: ERM-BF421a: vials containing about 1 g of potato powder, ERM-BF421b: vials containing about 0.5 g of potato powder.

CRMs for genetically modified 281-24-236 x 3006-210-23 cotton seed (ERM-BF422)

Four CRMs of dried cotton seed powder with different mass fractions of genetically modified (281-24-236 x 3006-210-23) cotton seed were produced.

	Certified value 281-24-236 x 3006-210-23 cotton seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF422a	< 0.5	-
ERM-BF422b	> 979	-
ERM-BF422c	10.0	1.7
ERM-BF422d	100	16

Availability: Vials containing about 1 g of cotton seed powder.

CRMs for genetically modified MIR604 maize (ERM-BF423)

Four CRMs of dried maize powder with different mass fractions of genetically modified (MIR604) maize were produced.

	Certified value MIR604 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF423a	< 0.9	-
ERM-BF423b	1.0	- 0.3 ; + 1.0
ERM-BF423c	9.8	- 0.9 ; + 1.3
ERM-BF423d	98.5	- 2.6 ; + 2.9

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified 59122 maize (ERM-BF424)

Four CRMs of dried maize powder with different mass fractions of genetically modified (59122) maize were produced.

	Certified value 59122 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF424a	< 1.2	-
ERM-BF424b	1.0	- 0.2 ; + 1.2
ERM-BF424c	9.9	- 0.8 ; + 1.4
ERM-BF424d	98.7	- 5.8 ; + 5.9

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified 356043 soya seed (ERM-BF425)

Four CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed were produced.

	Certified value 356043 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF425a	< 0.5	-
ERM-BF425b	1.0	0.4
ERM-BF425c *	10.0	1.1
ERM-BF425d	100	9

Availability: Vials containing about 1 g of soya seed powder.

* ERM-BF425c is also certified for the DNA copy number ratio.

	Certified value 356043 soya DNA copy number ratio (%)	Uncertainty (%)
ERM-BF425c	0.85	0.11

Substance	ERM-AD425 DNA fragments per plasmid	
	<u>Number</u>	
Fragment of 5' insert-to-plant junction DNA / pIRMM-0073	1	(negligible uncertainty)
Fragment of <i>le1</i> DNA / pIRMM-0073	1	(negligible uncertainty)

Availability: ERM-AD425 is available in vials.

CRMs for genetically modified 305423 soya seed (ERM-BF426)

Four CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed were produced.

	Certified value 305423 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF426a	< 0.8	-
ERM-BF426b	5.0	0.8
ERM-BF426c	10.0	1.0
ERM-BF426d	100	7

Availability: Vials containing about 1 g of soya seed powder.

CRMs for genetically modified 98140 maize (ERM-BF427)

Four CRMs of dried maize seed powder with different mass fractions of genetically modified (98140) maize were produced.

	Certified value 98140 maize seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF427a	< 0.4	-
ERM-BF427b	5.0	0.6
ERM-BF427c *	20.0	0.8
ERM-BF427d	100	4

Availability: Vials containing about 1 g of maize seed powder.

* ERM-BF427c is also certified for the DNA copy number ratio.

	Certified value 98140 maize DNA copy number ratio (%)	Uncertainty (%)
ERM-BF427c	1.75	0.13

	ERM-AD427 DNA fragments per plasmid
Fragment of 5' insert-to-plant junction DNA / pIRMM-0090 Fragment of <i>hmg</i> DNA / pIRMM-0090	1 1 <u>Number</u> (negligible uncertainty) (negligible uncertainty)

Availability: ERM-AD427 is available in vials.

CRMs for genetically modified GHB119 cotton (ERM-BF428)

Three CRMs of dried cotton seed powder with different mass fractions of genetically modified (GHB119) cotton were produced.

	Certified value GHB119 cotton seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF428a	< 0.2	-
ERM-BF428b	10	4
ERM-BF428c	100	11

Availability: Vials containing about 1 g of cotton seed powder.

CRMs for genetically modified T304-40 cotton (ERM-BF429)

Three CRMs of dried cotton seed powder with different mass fractions of genetically modified (T304-40) cotton were produced.

	Certified value T304-40 cotton seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF429a	< 0.4	-
ERM-BF429b	10	1.3
ERM-BF429c	100	11

Availability: Vials containing about 1 g of cotton seed powder.

CRMs for genetically modified AM04-1020 potato (ERM-BF430)

Two CRMs for the detection of genetically modified AM04-1020 potato were produced.

	Certified value AM04-1020 potato mass fraction (g/kg)
ERM-BF430a	0

	Certified identity
ERM-BF430b	Positive for event AM04-1020

Availability: Vials containing about 1 g of potato powder.

CRMs for genetically modified AV43-6-G7 potato (ERM-BF431)

Two CRMs for the detection of genetically modified AV43-6-G7 potato were produced.

Certified value AV43-6-G7 potato mass fraction (g/kg)	
ERM-BF431a	0
Certified identity	
ERM-BF431b	Positive for event AV43-6-G7

Availability: Vials containing about 1 g of potato powder.

CRMs for genetically modified DAS-68416-4 soya seed (ERM-BF432)

Four CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed DAS-68416-4 were produced.

	Certified value DAS-68416-4 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF432a	< 0.3	-
ERM-BF432b	5.0	0.6
ERM-BF432c	10.0	1.7
ERM-BF432d	100	13

Availability: Vials containing about 1 g of soya seed powder.

CRMs for genetically modified DAS-40278-9 maize (ERM-BF433)

Four CRMs of dried maize powder with different mass fractions of genetically modified DAS-40278-9 maize were produced.

	Certified value DAS-40278-9 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF433a	< 0.3	-
ERM-BF433b	5.0	0.6
ERM-BF433c	10.0	0.9
ERM-BF433d	100	8

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified 73496 rapeseed (ERM-BF434)

Five CRMs of dried rapeseed powder with different mass fractions of genetically modified rapeseed 73496 were produced.

	Certified value 73496 rapeseed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF434a	< 0.04	-
ERM-BF434b	>988	-
ERM-BF434c	1.00	0.15
ERM-BF434d	10.0	1.4
ERM-BF434e	100	12

Availability: Vials containing about 1 g of rapeseed powder.

CRMs for genetically modified PH05-026-0048 potato (ERM-BF435)

Two CRMs for the detection of genetically modified PH05-026-0048 potato were produced.

	Certified value PH05-026-0048 potato mass fraction (g/kg)
ERM-BF435a	< 0.4

	Certified identity
ERM-BF435b	Positive for event PH05-026-0048

Availability: Vials containing about 1 g of potato powder

CRMs for genetically modified DAS-44406-6 soya seed (ERM-BF436)

Five CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed DAS-44406-6 were produced.

	Certified value DAS-44406-6 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF436a	< 0.06	-
ERM-BF436b	>986	-
ERM-BF436c	1.00	0.14
ERM-BF436d	10.0	1.0
ERM-BF436e	100	9

Availability: Vials containing about 1 g of soya seed powder.

CRMs for genetically modified DAS-81419-2 soya seed (ERM-BF437)

Five CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed DAS-81419-2 were produced.

	Certified value DAS-81419-2 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF437a	< 0.07	-
ERM-BF437b	>986	-
ERM-BF437c	0.99	0.12
ERM-BF437d	9.9	1.5
ERM-BF437e	100	9

Availability: Vials containing about 1 g of soya seed powder.

CRMs for genetically modified VCO-Ø1981-5 maize (ERM-BF438)

Five CRMs of dried maize powder with different mass fractions of genetically modified VCO-Ø1981-5 maize were produced.

	Certified value VCO-Ø1981-5 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF438a	< 0.06	-
ERM-BF438b	>986	-
ERM-BF438c	1.00	0.13
ERM-BF438d	10.0	0.8
ERM-BF438e	100	5

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified DP-ØØ4114-3 maize (ERM-BF439)

Five CRMs of dried maize powder with different mass fractions of genetically modified DP-ØØ4114-3 maize were produced.

	Certified value DP-ØØ4114-3 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF439a	< 0.06	-
ERM-BF439b	>986	-
ERM-BF439c	1.00	0.13
ERM-BF439d	10.0	0.8
ERM-BF439e	100	5

Availability: Vials containing about 1 g of maize powder.

CRMs for genetically modified DAS-81910-7 cotton (ERM-BF440)

Five CRMs of dried cotton seed powder with different mass fractions of genetically modified (DAS-81910-7) cotton were produced.

	Certified value DAS-81910-7 cotton seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF440a	< 0.07	-
ERM-BF440b	>986	-
ERM-BF440c	1.00	0.08
ERM-BF440d	10.0	0.8
ERM-BF440e	100	7

Availability: Vials containing about 1 g of cotton seed powder.

2.2.2 CERTIFIED FOR NATURAL TOXINS AND XENOBIOTICS

Substance	BCR-459 Coconut oil (µg/kg)
Pyrene	< 0.9
Chrysene	< 0.6
Benzo[k]fluoranthene	< 0.2
Benzo[a]pyrene	< 0.3
Benz[ghi]perylene	< 0.2
Indeno[1,2,3-cd]pyrene	< 0.2

Availability: BCR-459 is provided in ampoules containing approximately 45 g.

Polychlorinated biphenyls (IUPAC No.)	BCR-450 Natural milk powder ($\mu\text{g}/\text{kg}$)	ERM-BB444 Natural pork fat (blank) ($\mu\text{g}/\text{kg}$)	ERM-BB445 Spiked pork fat (very low level) ($\mu\text{g}/\text{kg}$)	ERM-BB446 Spiked pork fat (low level) ($\mu\text{g}/\text{kg}$)
28		< 2	14.8 \pm 1.3	29.6 \pm 2.1
52	1.16 \pm 0.17	< 2	12.9 \pm 0.9	25.5 \pm 1.8
101		< 2	12.5 \pm 1.2	30 \pm 4
118	3.3 \pm 0.4	< 2	12.7 \pm 1.3	30.2 \pm 2.7
138		< 2	14.6 \pm 1.6	32 \pm 4
153	19.0 \pm 0.7	< 2	13.1 \pm 1.1	30.8 \pm 2.4
156	1.62 \pm 0.20			
170	4.8 \pm 0.6			
180	11.0 \pm 0.7	< 2	12.6 \pm 0.9	29.8 \pm 2.5
sum (28, 52, 101, 118, 138, 153, 180)		< 14	93 \pm 7	207 \pm 11
*PDBE 47		(3.7)	(3.9)	(6.1)
γ -HCH (lindane)		(5.7)	(5.6)	(4.6)

* 2,2'4,4'-Tetrabromo-diphenylether. / Values in brackets are not certified.

Availability: BCR-450 in brown glass bottles of about 20 g. ERM-BB444 to 446 in glass ampoules of about 5 g.

Polychlorinated biphenyls (IUPAC No.)	BCR-349 Cod liver oil ($\mu\text{g}/\text{kg}$)	ERM-BB350 Fish oil (ng/g)
28	68 \pm 8	21.3 \pm 1.1
52	149 \pm 21	37.4 \pm 2.2
74		23.0 \pm 1.9
95		(38) \pm 47)
99		62 \pm 6
101	372 \pm 18	111 \pm 5
105		25.8 \pm 2.1
110		54.1 \pm 2.8
118	460 \pm 40	84 \pm 4
138		137 \pm 10
149		88 \pm 9
153	940 \pm 40	220 \pm 11
156		20.1 \pm 1.3
163		(43) \pm 73)
167		(17) \pm 27)
177		25.8 \pm 2.0
180	282 \pm 23	67 \pm 4
183		22.5 \pm 1.8
187		67 \pm 5
194		23.4 \pm 1.5
196		41 \pm 7

Values in brackets are not certified.

Availability: BCR-349 and ERM-BB350 are provided in sealed glass ampoules containing approximately 2 g fish oil.

Substance	BCR-598 Cod liver oil ($\mu\text{g}/\text{kg}$)		
HCB	55.7	\pm	2.0
α -HCH	42	\pm	3
β -HCH	16	\pm	3
γ -HCH	23	\pm	4
γ -Chlordane	6.9	\pm	1.6
α -Chlordane	24.4	\pm	1.8
Oxychlordane	11.0	\pm	1.8
Transnonachlor	39	\pm	4
Dieldrin	59	\pm	4
p,p'-DDE	0.61×10^3	\pm	0.04×10^3
o,p'-DDD	30	\pm	4
p,p'-DDD	0.40×10^3	\pm	0.03×10^3
p,p'-DDT	0.179×10^3	\pm	0.018×10^3

Availability: BCR-598 is provided in sealed glass ampoules containing approximately 5 g under dry argon.

Substance	ERM-BB430		
	Pork fat	(mg/kg)	
HCB	0.193	±	0.017
α-HCH	0.25	±	0.04
β-HCH	0.109	±	0.010
γ-HCH	(1.87	±	0.31)
β-HEPO	0.213	±	0.016
Dieldrin	(0.21	±	0.05)
Endrin	(0.055	±	0.016)
p,p'-DDT	0.48	±	0.07
p,p'-DDD	0.222	±	0.022
p,p'-DDE	0.38	±	0.09

Values in brackets are not certified.

Availability: ERM-BB430 is provided in sealed amber glass ampoules containing approximately 5 g of material bottled under argon.

Substance	BCR-115		
	Animal feed	(mg/kg)	
HCB	0.019 4	±	0.001 4
β-HCH	0.023 4	±	0.002 6
γ-HCH	0.021 8	±	0.002 0
Heptachlor	0.019 0	±	0.001 5
γ-Chlordane	0.048	±	0.006
α-Endosulfan	0.046	±	0.004
Dieldrin	0.018 1	±	0.002 3
Endrin	0.046	±	0.006
o,p'-DDT	0.046	±	0.005
p,p'-DDE	0.047	±	0.004

Availability: BCR-115 is provided in sealed hard glass ampoules containing approximately 30 g under dry N₂. The sample is a homogeneous animal feed obtained from commonly used ingredients and enriched with organochlorine pesticides.

Substance	BCR-187			BCR-188		
	Milk powder	(μ g/kg)		Milk powder (spiked)	(μ g/kg)	
HCB	1.45	±	0.21	37.4	±	2.7
β-HCH			(> 3.6)	12.0	±	1.2
γ-HCH						(> 25)
β-HEPO				32.0	±	1.9
p,p'-DDE	6.6	±	0.6	51	±	4
Dieldrin				36.1	±	2.5
Endrin				6.2	±	0.9
p,p'-DDT				69	±	5

Values in brackets are not certified.

Availability: The samples are provided in sealed hard glass ampoules containing about 20 g (under Argon).

Substance	BCR-607		
	Natural spray dried milk powder	(ng/kg)	
2,3,7,8 - T ₄ CDD (D48)	0.25	±	0.03
1,2,3,7,8 - P ₅ CDD (D54)	0.79	±	0.04
1,2,3,4,7,8 - H ₆ CDD (D66)	0.42	±	0.07
1,2,3,6,7,8 - H ₆ CDD (D67)	0.98	±	0.11
1,2,3,7,8,9 - H ₆ CDD (D70)	0.34	±	0.05
2,3,7,8 - T ₄ CDF (F83)	0.05	±	0.03
1,2,3,7,8 - P ₅ CDF (F94)	0.054	±	0.013
2,3,4,7,8 - P ₅ CDF (F114)	1.81	±	0.13
1,2,3,4,7,8 - H ₆ CDF (F118)	0.94	±	0.04
1,2,3,6,7,8 - H ₆ CDF (F121)	1.01	±	0.09
2,3,4,6,7,8 - H ₆ CDF (F130)	1.07	±	0.05

Availability: Amber glass bottles containing approximately 100 g.

Substance	ERM-BC190 Rapeseed (colza)	ERM-BC366 Rapeseed (colza)	ERM-BC367 Rapeseed (colza)
	<u>Certified values</u>	<u>Certified values</u>	<u>Certified values</u>
Total glucosinolate content	23 \pm 4 mmol/kg	11.9 \pm 1.3 mmol/kg	99 \pm 9 mmol/kg
Sulphur content	4.72 \pm 0.22 g/kg	3.31 \pm 0.17 g/kg	10.3 \pm 0.5 g/kg

Availability: 20 g whole seed in an aluminium plastic laminated sachet sealed under nitrogen.

Substance	BCR-262R Defatted peanut meal (blank)	BCR-263R Defatted peanut meal (medium level)	BCR-264 Defatted peanut meal (high level)	BCR-375 Compound feed (very low level blank)	ERM-BE375 Compound feed (very low level)	ERM-BE376 Compound feed (high level)
	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)
Aflatoxin B1	< 3	17.1 \pm 2.4	206 \pm 13	< 1	2.6 \pm 0.4	12.9 \pm 1.8
Aflatoxin B2		3.0 \pm 0.4			0.20 \pm 0.04	0.68 \pm 0.10
Aflatoxin G1		3.0 \pm 0.5			0.40 \pm 0.10	5.2 \pm 0.8
Aflatoxin G2		(0.62 \pm 0.21)			< 0.2	
Sum of aflatoxin B1, B2, G1 and G2		(23.7 \pm 2.5)				

Values in brackets are not certified.

Availability: Sachets sealed under vacuum containing about 100 g (BCR-263R) and about 150 g (BCR-264) of finely ground defatted peanut meal.

BCR-262R is available in 280 mL amber glass bottles containing about 100 g of the peanut meal, additionally sealed in foil-laminate sachet.

BCR-375 is supplied in units of about 50 g of a finely ground compound feed, ERM-BE375 and ERM-BE376 consist of 2 bottles filled with about 75 g of compound feedingstuff each.

	Description	Aflatoxin M ₁ ($\mu\text{g}/\text{kg}$)
ERM-BD282	Whole milk powder (zero level)	< 0.02
ERM-BD283	Whole milk powder (low level)	0.111 \pm 0.018
ERM-BD284	Whole milk powder (high level)	0.44 \pm 0.06

Availability: The materials are provided in units of 30 g in amber glass bottles filled and sealed under nitrogen.

Substance	BCR-401R Peanut butter (very low level)	BCR-385R Peanut butter (low level)
	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)	<u>Mass fraction</u> ($\mu\text{g}/\text{kg}$)
Aflatoxin B1	< 0.2	1.77 \pm 0.30
Aflatoxin B2	< 0.2	0.48 \pm 0.08
Aflatoxin G1	< 0.2	0.9 \pm 0.4
Aflatoxin G2	< 0.2	0.30 \pm 0.12
Total		3.5 \pm 0.5

Availability: The samples are supplied in units of about 100 g in aluminium cans.

Substance	BCR-377 Maize Flour (very low level blank)	BCR-396 Wheat Flour (very low level blank)
	<u>Mass fraction</u> (mg/kg)	<u>Mass fraction</u> (mg/kg)
DON	< 0.05	< 0.05

Availability: Sachets sealed under vacuum containing about 150 g of sealed finely ground flour.

Substance	BCR-471 Wheat (blank)
Ochratoxin A	<u>Mass fraction</u> ($\mu\text{g/kg}$) < 0.6

Availability: Units of about 55 g in foil-laminate pouches sealed under vacuum.

Compound	BCR-543 Mussel
Saxitoxin·2HCl Saxitoxin·2HCl (if enriched with BCR-663) dc-saxitoxin	<u>Mass fraction</u> (mg/kg) < 0.07 0.48 \pm 0.06 < 0.04

Availability: BCR-543 is available in heat-sealed laminate sachets containing about 15 g of material each. BCR-663 is available in ampoules containing 1 mL.

Substance	ERM-BC716 Maize	ERM-BC717 Maize
DON NIV ZON	<u>Mass fraction</u> ($\mu\text{g/kg}$) < 5	<u>Mass fraction</u> ($\mu\text{g/kg}$) 673 \pm 87 53 \pm 10 83 \pm 9

Availability: ERM-BC716 and ERM-BC717 are supplied in sachets containing at least 60 g.

IRMM-359: *Staphylococcus aureus* enterotoxin A (SEA) in cheese

IRMM-359	Diagnostic specificity	
	Certified value [%]	One-sided lower confidence limit [%]
IRMM-359a	100	97.3
IRMM-359b	100	97.5
IRMM-359c	100	97.6

Availability: Provided in a set of IRMM-359a, IRMM-359b and IRMM-359c, in plastic sachets protected in an alumina pouch.

2.2.3 CERTIFIED FOR THE TOTAL ELEMENT CONTENT

Substance	ERM-BD150 Skimmed milk powder			ERM-BD151 Skimmed milk powder		
	(g/kg)			(g/kg)		
Ca	13.9	±	0.8	13.9	±	0.7
Cl	9.7	±	2.0	9.8	±	1.2
K	17.0	±	0.7	17.0	±	0.8
Mg	1.26	±	0.10	1.26	±	0.07
Na	4.18	±	0.19	4.19	±	0.23
	(mg/kg)			(mg/kg)		
Cd	0.0114	±	0.029	0.106	±	0.013
Cu	1.08	±	0.06	5.00	±	0.23
Fe	4.6	±	0.5	53	±	4
Hg	0.060	±	0.007	0.52	±	0.04
I	1.73	±	0.14	1.78	±	0.17
Mn	0.289	±	0.018	0.29	±	0.03
Pb	0.019	±	0.004	0.207	±	0.014
Se	0.188	±	0.014	0.19	±	0.04
Zn	44.8	±	2.0	44.9	±	2.3

Availability: Glass bottles containing 20 g of skimmed milk powder.

Substance	ERM-BB184 Bovine muscle (mg/kg)		ERM-BB186 Pig kidney (mg/kg)		ERM-BB422 Fish muscle (mg/kg)		BCR-185R Bovine liver (mg/kg)	
As	0.0234 ± 0.0026		(0.008 ± 0.006)		12.7 ± 0.7		0.0330 ± 0.0029	
Cd	0.0022 ± 0.0004		1.09 ± 0.05		0.0075 ± 0.0018		0.544 ± 0.017	
Cu	2.31 ± 0.09		36.5 ± 1.8		1.67 ± 0.16		277 ± 5	
Fe	75 ± 4		255 ± 13		9.4 ± 1.4		-	
Hg	(0.0018 ± 0.0010)		(0.023 ± 0.011)		0.601 ± 0.030		-	
I	-		-		1.4 ± 0.4		-	
Mn	0.276 ± 0.013		7.26 ± 0.25		0.368 ± 0.028		11.07 ± 0.29	
Pb	-		0.040 ± 0.005		-		0.172 ± 0.009	
Se	0.45 ± 0.04		10.3 ± 0.9		1.33 ± 0.13		1.68 ± 0.14	
Zn	146 ± 7		134 ± 5		16.0 ± 1.1		138.6 ± 2.1	

Values in brackets are not certified.

Availability: BCR-185R is provided in units of 15 g as lyophilised powders, in screw-capped glass bottles.

ERM-BB184 is provided in units of 7 g as lyophilised powder, in brown-glass vials.

ERM-BB186 and ERM-BB422 are provided in units of 10 g as lyophilised powders, in brown-glass vials.

Substance	BCR-191 Brown bread		
Cd µg/kg	28.4	±	1.4
Cu mg/kg	2.63	±	0.07
Fe mg/kg	40.7	±	2.3
Mn mg/kg	20.3	±	0.7
Pb µg/kg	187	±	14
Zn mg/kg	19.5	±	0.5

Availability: BCR-191 is provided in units of 25 g.

Substance	IRMM-804 Rice flour (mg/kg)		
As	0.049	±	0.004
Cd	1.61	±	0.07
Cu	2.74	±	0.24
Mn	34.2	±	2.3
Pb	0.42	±	0.07
Zn	23.1	±	1.9

Availability: Amber glass bottles containing about 15 g of powdered material.

Substance	BCR-273			BCR-274		
	Single cell protein (mg/g)			Single cell protein (µg/g)		
As				0.132	±	0.014
Ca	11.97	±	0.14			
Cd				0.030	±	0.002
Co				0.039	±	0.003
Cu				13.1	±	0.4
Fe	0.156	±	0.004			
K	2.22	±	0.05			
Mn				51.9	±	1.2
N	121.6	±	0.8			
P	26.8	±	0.4			
Pb				0.044	±	0.010
Se				1.03	±	0.05
Zn				42.7	±	1.0

Availability: BCR-273 and BCR-274 are provided in units of 10 g as a dry powder in sealed glass ampoules.

Substance	BCR-679		
	White cabbage		
B	mg/kg	(27.7	± 1.9)
Ba	mg/kg	(10.3	± 0.6)
Ca	mg/kg	(7768	± 655)
Cd	mg/kg	1.66	± 0.07
Cr	mg/kg	(0.6	± 0.1)
Cu	mg/kg	2.89	± 0.12
Fe	mg/kg	55.0	± 2.5
Hg	µg/kg	6.3	± 1.4
Mg	mg/kg	(1362	± 127)
Mn	mg/kg	13.3	± 0.5
Mo	mg/kg	14.8	± 0.5
Ni	mg/kg	27.0	± 0.8
P	mg/kg	(3307	± 241)
Sb	µg/kg	20.6	± 2.6
Sr	mg/kg	11.8	± 0.4
Tl	µg/kg	3.0	± 0.3
Zn	mg/kg	79.7	± 2.7

Values in brackets are not certified.

Availability: BCR-679 is provided in units of 15 g.

Substance	ERM-BC381			ERM-BC382			BCR-383			ERM-BB384		
	Rye Flour			Wheat Flour			Haricots Verts (Beans)			Lyophilised pork muscle		
<u>Major components (g / 100 g)</u>												
Glucose							(12.4)					
Fructose							(4.6)					
Sucrose							(1.0)					
N (Kjeldahl)	1.562	±	0.014	1.851	±	0.017	1.05	±	0.04	14.2	±	0.4
Fat	1.36	±	0.16	1.39	±	0.17				8.99	±	0.20
Starch ¹⁾	72.2	±	1.9	81.2	±	1.7						
Starch & Sugars ²⁾							(78.9)					
Dietary Fibre (Englyst)							(10.9)					
Dietary Fibre (AOAC 1985/1988) ³⁾							11.9	±	0.6			
Ash at 550 °C	1.08	±	0.11	0.60	±	0.10	2.39	±	0.10	4.51	±	0.19
<u>Essential elements (g/kg)</u>												
Ca	0.32	±	0.04	0.210	±	0.018	2.85	±	0.23	0.164	±	0.021
K	3.35	±	0.11	1.88	±	0.08	7.8	±	0.4			
Mg	0.567	±	0.013	0.247	±	0.010		(0.9)		1.03	±	0.04
Na							0.075	±	0.007	1.86	±	0.15
P	2.01	±	0.07	1.19	±	0.07		(1.8)		8.7	±	0.5

Values in brackets are not certified.

Availability: ERM-BC381 and ERM-BC382: 100 ml amber vial containing 37 g flour; ERM-BB384: 2 vials of 9 g lyophilised material; BCR-383: 100 g of powdered material in food grade laminated plastic/aluminium sachets sealed under nitrogen.

¹⁾ Mass fraction of polysaccharide in dry matter

²⁾ Mass fraction of monosaccharides in dry matter

³⁾ Prosky L. et al., J Assoc Off Anal Chem (1985) 68: 677-679, (1988) 71: 1017-1023

2.2.4 CERTIFIED FOR PROXIMATES AND CONVENTIONAL PROPERTIES

Substance		BCR-162R Soya-maize oil blend		
<u>Methyl ester of</u>		<u>Mass fraction fatty acid methyl ester /</u> <u>Total fatty acid methyl ester (g/100 g)</u>		
16:0	n-hexadecanoic acid	10.74	±	0.16
18:0	n-octadecanoic acid	2.82	±	0.04
18:1	n-octadecenoic acids		(26.5)	
18:2	n-octadecadienoic acids		(54.68)	
18:3	n-octadecatrienoic acids		(3.80)	
9c-18:1	n-octadecenoic acid	25.4	±	0.4
9c,12c-18:2	n-octadecadienoic acid	54.13	±	0.25
9c,12c,15c-18:3	n-octadecatrienoic acid	3.35	±	0.05

Values in brackets are not certified.

Availability: Each unit contains approximately 5.5 g soya-maize oil blend in 10 mL amber ampoule.

Substance		BCR-163 Beef-pork fat oil blend		
<u>Methyl ester of</u>		<u>Mass fraction fatty acid methyl ester /</u> <u>Total fatty acid methyl ester ¹⁾ (g/100 g)</u>		
14:0	n-tetradecanoic acids	2.29	±	0.04
16:0	n-hexadecanoic acids	25.96	±	0.30
16:1	n-hexadecenoic acids	2.58	±	0.16
18:0	n-octadecanoic acids	18.29	±	0.17
18:1	n-octadecenoic acids	38.3	±	0.4
18:2	n-octadecadienoic acids	7.05	±	0.17
18:3	n-octadecatrienoic acids	0.86	±	0.14
<u>Sterols</u> Cholesterol		<u>Mass fraction (mg/100 g) in fat</u>		
		134	±	5

Availability: In units of 2 × 5 mL in dark glass ampoules sealed under nitrogen.

- 1) Includes any geometric (i.e. cis/trans) and positional isomers, expressed as mass fraction of total fatty acid (methyl esters) derived from triglycerides.
 - 2) These components are included in the Certified Value for this group of fatty acids.
- The report gives additional indicative values: Fatty Acids and "Total" Sterol Mass Fraction.

Component		BCR-633 Tracers in anhydrous butter fat (mg/kg)		
β -Apo-8'-carotenic acid ethyl ester		26.5	±	1.4
β -Sitosterol		530	±	29
Stigmasterol		147	±	11
<i>n</i> -Heptanoic acid triglyceride		1.06 * 10 ³	±	0.04 * 10 ³

Availability: BCR-633 is supplied in units of about 5 g each in amber glass ampoules, which were filled under inert gas conditions (nitrogen).

Component		IRMM-801 Cocoa Butter (%)		
1,3-dipalmitoyl-2-oleyl-glycerol		18.14	±	0.26
1-palmitoyl-2-oleoyl-3-stearoyl-glycerol		44.68	±	0.30
1,2-dioleoyl-3-palmitoyl-glycerol		2.26	±	0.16
1,3-distearoyl-2-oleoyl-glycerol		31.63	±	0.29
1,2-dioleoyl-3-stearoyl-glycerol		3.29	±	0.17

Availability: IRMM-801 is supplied in units of 5 g in brown amber glass ampoules sealed under argon/helium.

Component	BCR-519 Anhydrous butter fat (%)		
Cholesterol	0.30	±	0.03
C24	0.05	±	0.02
C26	0.25	±	0.03
C28	0.59	±	0.04
C30	1.15	±	0.05
C32	2.43	±	0.12
C34	5.64	±	0.18
C36	10.47	±	0.19
C38	12.53	±	0.22
C40	10.03	±	0.16
C42	6.69	±	0.10
C44	6.11	±	0.08
C46	6.86	±	0.08
C48	8.69	±	0.15
C50	11.40	±	0.24
C52	10.96	±	0.25
C54	5.89	±	0.13

Availability: The material consists of anhydrous butter fat and is supplied in units of two glass ampoules sealed under nitrogen, each containing approximately 5 mL fat. At normal ambient temperature BCR-519 is solid.

	BCR-632 Butter fat					
Compound	BCR-632A Pure butter fat (g/100 g)			BCR-632B Adulterated butter fat (g/100 g)		
C24	0.07	±	0.04	0.08	±	0.04
Cholesterol	0.289	±	0.012	0.278	±	0.011
C26	0.33	±	0.06	0.34	±	0.06
C28	0.74	±	0.07	0.75	±	0.06
C30	1.37	±	0.08	1.46	±	0.07
C32	2.83	±	0.14	3.30	±	0.12
C34	6.09	±	0.29	6.57	±	0.25
C36	10.7	±	0.5	11.1	±	0.4
C38	12.5	±	0.4	12.7	±	0.4
C40	10.05	±	0.19	10.07	±	0.17
C42	7.07	±	0.13	7.10	±	0.10
C44	6.68	±	0.12	6.57	±	0.12
C46	7.36	±	0.17	7.12	±	0.17
C48	8.74	±	0.21	8.42	±	0.19
C50	10.74	±	0.24	10.28	±	0.19
C52	9.8	±	0.4	9.36	±	0.28
C54	4.7	±	0.5	4.5	±	0.4

Availability: BCR-632 is available as a set of BCR-632A and BCR-632B: 2 amber glass ampoules each with approximately 5 mL.

	BCR-121 Wholemeal flour (mg/kg)	BCR-122 Margarine (mg/kg)	BCR-431 Brussels sprouts (mg/kg)	BCR-485 Mixed vegetables (mg/kg)	BCR-487 Pig's liver (mg/kg)	ERM-BD600 Whole milk powder (mg/kg)
A (all-trans-retinol)						3.8 ± 0.6
A (all-trans-retinol and 13-cis-retinol)						4.1 ± 0.8
B ₁ (49pprox.49)	4.63 ± 0.39			3.07 ± 0.34	8.6 ± 1.1	4.5 ± 0.6
B ₂ (riboflavin)				4.8 ± 0.8	106.8 ± 5.6	16.7 ± 1.4
B ₆ (total pyridoxine)	4.10 ± 1.02			19.3 ± 2.9	19.3 ± 2.9	74 ± 11
B ₁₂ (cyanocobalamin)			4830 ± 240	1.12 ± 0.09	0.32 ± 0.07	
C (total ascorbate)					86 ± 15	
D ₃ (cholecalciferol)		0.125 ± 0.007				
E (tocopherol)		241 ± 12				
Folate (total)	0.50 ± 0.07			3.15 ± 0.28	13.3 ± 1.3	
Niacin			43 ± 3			
Trans-α-carotene				10.5 ± 0.6		
Trans-β-carotene				23.7 ± 1.5		
Total-α-carotene				9.8 ± 0.7		
Total-β-carotene				25.6 ± 1.2		
Lutein				12.5 ± 0.8		
Lutein + zeaxanthin				22.3 ± 1.3		
5-methyltetrahydro-folic acid (5-MTHF)				(2.14 ± 0.42)		

Values in brackets are not certified;

Availability: BCR-121: about 50 g unit size.

BCR-122: can filled with about 200 g.

BCR-431: about 20 g lyophilised and powdered material in food grade plastic/aluminium laminated sachets under nitrogen.

BCR-485: about 25 g unit size.

BCR-487: about 15 g unit size.

ERM-BD600: sachet with about 100 g.

	Description	Amylose mass fraction (g/100 g)
BCR-465	Rice flour (low level)	15.40 ± 0.30
BCR-466	Rice flour (medium level)	23.1 ± 0.5
BCR-467	Rice flour (high level)	27.7 ± 0.8

Availability: 10 g of ground rice flour in vacuum sealed laminated polyester/aluminium/polyethylene sachets.

Substance	BCR-644 Artificial foodstuff	BCR-645 Artificial foodstuff
Mass fraction on dry mass basis (g / 100 g)		
Fructose	16.2 ± 1.1	
Sucrose	10.81 ± 0.25	26.2 ± 0.8
Lactose	15.85 ± 0.29	27.8 ± 0.6
Starch / glucose	35.1 ± 1.2	25.2 ± 0.9

Availability: BCR-644 and BCR-645 are supplied in units of approximately 50 g in 125 mL amber glass bottles.

	BCR-651 Beer (% ethanol v/v)	BCR-652 Beer (% ethanol v/v)	BCR-653 Wine (% ethanol v/v)
Ethanol	0.505 ± 0.006	0.051 ± 0.002	0.539 ± 0.007

Availability: Amber glass ampoule, flushed with nitrogen, containing 10 mL of sample.

Substance	ERM-BC381 Rye Flour	ERM-BC382 Wheat Flour	BCR-383 Haricots Verts (Beans)	ERM-BB384 Lyophilised pork muscle
<u>Major components (g / 100 g)</u>				
Glucose			(12.4)	
Fructose			(4.6)	
Sucrose			(1.0)	
N (Kjeldahl)	1.562 ± 0.014	1.851 ± 0.017	1.05 ± 0.04	14.2 ± 0.4
Fat	1.36 ± 0.16	1.39 ± 0.17		8.99 ± 0.20
Starch ¹⁾	72.2 ± 1.9	81.2 ± 1.7		
Starch & Sugars ²⁾			(78.9) (10.9)	
Dietary Fibre (Englyst)			11.9 ± 0.6	
Dietary Fibre (AOAC 1985/1988) ³⁾				
Ash at 550 °C	1.08 ± 0.11	0.60 ± 0.10	2.39 ± 0.10	4.51 ± 0.19
<u>Essential elements (g/kg)</u>				
Ca	0.32 ± 0.04	0.210 ± 0.018	2.85 ± 0.23	0.164 ± 0.021
K	3.35 ± 0.11	1.88 ± 0.08	7.8 ± 0.4	
Mg	0.567 ± 0.013	0.247 ± 0.010	(0.9)	1.03 ± 0.04
Na			0.075 ± 0.007	1.86 ± 0.15
P	2.01 ± 0.07	1.19 ± 0.07	(1.8)	8.7 ± 0.5

Values in brackets are not certified.

Availability: ERM-BC381 and ERM-BC382: 100 ml amber vial containing 37 g flour; ERM-BB384: 2 vials of 9 g lyophilised material; BCR-383: 100 g of powdered material in food grade laminated plastic/aluminium sachets sealed under nitrogen.

1) Mass fraction of polysaccharide in dry matter

2) Mass fraction of monosaccharides in dry matter

3) Prosky L. et al., J Assoc Off Anal Chem (1985) 68: 677-679, (1988) 71: 1017-1023

Substance	BCR-685 Skim milk powder (g / 100 g)
<u>Mass fraction</u>	
Crude protein (Kjeldahl-N x 6.38)	38.2 ± 0.4
Fat	0.96 ± 0.12

Availability: BCR-685 is supplied in units of about 50 g, in amber glass bottles.

Quantity	BCR-446 Low oil content rapeseed	BCR-447 Medium oil content rapeseed
	Mass fraction (g/100 g)	Mass fraction (g/100 g)
'As is' Oil	39.49 ± 0.15	41.99 ± 0.15
Moisture and volatiles	7.01 ± 0.07	7.42 ± 0.07
'Dry basis' Oil	42.48 ± 0.15	45.36 ± 0.15

Availability: About 150 g of rapeseed in a specially laminated bag containing an oxygen absorber and sealed under dry argon.

Substance	BCR-708 Dairy feed			BCR-709 Pig feed			
<u>Mass fraction</u>							
Crude protein	240	±	12	199	±	5	g/kg
Crude oils and fats	65	±	8	51	±	14	g/kg
Crude fibre	93	±	14	56	±	12	g/kg
Crude ash	50.0	±	3.0	42	±	4	g/kg
Ca	4.8	±	0.5	1.05	±	0.16	g/kg
Cu	37	±	4	173	±	25	mg/kg
Mg	1.47	±	0.22	1.89	±	0.30	g/kg
P	4.7	±	0.4	5.4	±	0.7	g/kg

Availability: 100 mL amber glass bottle containing about 40 g of material.

Substance	ERM-BC514 Haricot beans (g/kg)	ERM-BC515 Carrot (g/kg)	ERM-BC516 Apple (g/kg)	ERM-BC517 Full fat soya flour (g/kg)	ERM-BD518 Bran breakfast cereal (g/kg)
<u>Dietary Fibres</u>					
Methods used:					
AOAC 1990	256 ± 5	311 ± 6	164 ± 4	126 ± 5	302 ± 8
Englyst (by gas chromatography)	198 ± 10	271 ± 6	137 ± 5	119 ± 7	241 ± 8
Uppsala	237 ± 15	298 ± 11	162 ± 8	128 ± 9	276 ± 18
AOAC 1992 MES-TRIS	259 ± 15	295 ± 4	149 ± 10	124 ± 21	305 ± 6
Englyst (by colorimetry)	201 ± 6	252 ± 12	134 ± 5	123 ± 8	250 ± 11

Availability: ERM-BC514 to BC517 and ERM-BD518 are supplied in units of approximately 25 g in food grade glass bottles sealed under vacuum.

	BCR-537 Plastic film A (mg/dm ²)	BCR-539 Plastic film C (mg/dm ²)
Overall migration by total immersion in olive oil 10 days at 40 °C	8.3 ± 1.0	
Overall migration by single sided cell in olive oil 10 days at 40 °C		
Overall migration by pouch in olive oil 10 days at 40 °C		6.1 ± 1.0

Availability: PET/foil/PE heat sealed pouches containing double thickness sheets of additive free linear low density polyethylene of (33 x 22.5) cm for BCR-537 and (45 x 25) cm for BCR-539.

2.2.5 CERTIFIED FOR MICROBIOLOGICAL PROPERTIES

BCR-528: Capsules filled with milk powder artificially contaminated by *Bacillus cereus* (ATCL 9139)

Procedure	Number of colony forming particles in one analytical portion ¹⁾		
	Certified value ²⁾	95 % confidence limits	
MEYP (ISO 7932) after 24 h incubation	53.4	51.7	- 55.2
MEYP (ISO 7932) after 48 h incubation	53.7	52.1	- 55.4
PEMBA (L 00.00 – 25) ³⁾ after 24 h incubation	55.0	52.8	- 57.4
PEMBA (L 00.00 – 25) ³⁾ after 48 h incubation	55.8	53.6	- 58.0

Availability: BCR-528 is provided in containers holding 10 gelatine capsules filled with artificially contaminated milk powder.

1) Number of colony forming particles of *Bacillus cereus* determined in one analytical portion. Analytical portion: A volume of (0.100 ± 0.002) mL from 10 mL peptone saline solution in which one capsule has been reconstituted.

2) This value is the geometric mean of 11 accepted sets of data, independently obtained by 11 laboratories.

3) German Federal Food Law method number.

IRMM-311: Genomic DNA (gDNA) of *Bacillus licheniformis* DSM 5749 in agarose inserts for Pulsed Field Gel Electrophoresis (PFGE)

SfiI digested DNA fragments in the size interval 50 kb – 90 kb		Fragment length (kb)		
Band no	1	89.6	±	4.7
	2	80.9	±	2.5
	3	75.3	±	2.7
	4	72.2	±	3.5
	5	66.9	±	1.9
	6	64.6	±	2.9
	7	60.3	±	1.3
	8	56.5	±	1.3
	9	53.9	±	1.3
	10	50.6	±	1.3

Availability: Each vial contains one agarose insert of undigested genomic DNA of *Bacillus licheniformis* DSM 5749 for PFGE.

IRMM-312: Genomic DNA (gDNA) of *Bacillus subtilis* DSM 5750 in agarose inserts for Pulsed Field Gel Electrophoresis (PFGE)

SfiI digested DNA fragments in the size interval 50 kb – 90 kb		Fragment length (kb)		
Band no	1	89.2	±	0.9
	2	81.4	±	0.8
	3	77.7	±	0.6
	4	62.5	±	1.8
	5	59.5	±	2.1
	6	44.0	±	2.4
	7	29.2	±	2.0
	8	23.6	±	1.3
	9	18.6	±	1.3

Availability: Each vial contains one agarose insert of undigested genomic DNA of *Bacillus subtilis* DSM 5750 for PFGE.

IRMM-313: Genomic DNA (gDNA) of *Campylobacter coli* (CNET068) and *Campylobacter jejuni* (CNET112) in agarose inserts for Pulsed Field Gel Electrophoresis (PFGE)

Smal digested DNA fragments		DNA fragment sizes (kb)		
Fragment no	2	458.8	±	2.0
	3	351.7	±	2.4
	4	303.0	±	2.3
	5	263.2	±	1.9
	6	188.2	±	1.2
	7	173.2	±	1.3
	8	131.1	±	1.5
	9	114.4	±	1.2
	10	95.5	±	1.4
	11	81.2	±	1.7
	12	54.9	±	2.2
	13	40.7	±	1.6
	14	(25.4	±	1.3)
	15	(17.6	±	0.3)
	16	(10.9	±	0.4)

Values in brackets are not certified.

Availability: Each vial contains one agarose plug for PFGE with undigested genomic DNA of *Campylobacter coli* CNET068 and *Campylobacter jejuni* CNET112 embedded.

IRMM-351: Escherichia coli 0157 in material spheres

	Number of colony forming unit (cfu)		
cfu per material sphere on nutrient agar	4	±	2
cfu per material sphere on enterohemolysin agar	4	±	2

Availability: Each vial contains one material sphere of Escherichia coli 0157 (NCTC 12900).

IRMM-352: Salmonella enteritidis in material spheres

	Number of colony forming unit (cfu)		
cfu per material sphere on nutrient agar	5	±	2
cfu per material sphere on xylose lysine deoxycholate agar	4	±	2

Availability: Each vial contains one material sphere of Salmonella enteritidis (NCTC 12694).

IRMM-354: Candida albicans in material spheres

	Number of colony forming unit (cfu)		
cfu per material sphere on nutrient agar	917	±	168
cfu per material sphere on Oxytetracyclin-Glucose-Yeast Extract agar (OGYE)	912	±	173

Availability: Each vial contains one material sphere of Candida albicans (NCPI 3179).

IRMM-355: Enterococcus faecalis in material spheres

	Number of colony forming unit (cfu)		
cfu per material sphere on horse blood agar	890	±	135
cfu per material sphere on Sianetz and Bartley agar	823	±	126

Availability: Each vial contains one material sphere of Enterococcus faecalis (CIP 106877).

IRMM-447: Genomic DNA (gDNA) of Listeria monocytogenes (strain 4B, NCTC 11994) with certified identity

	Indicative value (μ g)	
Mass of genomic DNA <i>Listeria monocytogenes</i> per vial	(1.1	± 0.7)

Values in brackets are not certified.

Availability: Vial containing approximately 1.1 μ g genomic DNA in lyophilised form and closed under argon atmosphere.

IRMM-448: Genomic DNA (gDNA) of Campylobacter Jejuni (NCTC 11351) with certified identity

Property		
Identity	Confirmed by ceuE gene	
Mass of genomic DNA per vial (ng)	(71	± 39)

Values in brackets are not certified.

Availability: Vial containing approximately 71 ng genomic DNA in lyophilised form and closed under argon atmosphere.

IRMM-449: Genomic DNA (gDNA) of *Escherichia coli* O157 (strain EDL 933) with certified identity

	Indicative value (μ g)
Mass of genomic DNA <i>Escherichia coli</i> per vial	(1.3 \pm 0.7)

Values in brackets are not certified.

Availability: Vial containing approximately 1.3 μ g genomic DNA in lyophilised form and closed under argon atmosphere.

2.2.6 CERTIFIED FOR VETERINARY DRUGS

	Description	Substance	Hormones in lyophilised bovine urine Mass concentration in reconstituted sample (μ g/L)		
			Content	Relevant below the certified value	Relevant above the certified value
BCR-502	Bovine urine	Clenbuterol	< 0.1		
		Salbutamol	< 0.2		
BCR-503	Bovine urine	Clenbuterol	2.5	0.4	0.4
		Salbutamol	2.3	0.6	0.9
BCR-504	Bovine urine	Clenbuterol	6.0	0.5	0.7
		Salbutamol	5.6	1.1	1.9

Value in brackets is not certified.

Availability: Units of lyophilised urine equivalent to about 5.0 mL in vials sealed under nitrogen.

	Description	Substance	Hormones in lyophilised bovine urine Mass concentration in reconstituted sample (μ g/kg)		
ERM-BB386 ⁽¹⁾	Bovine urine	Diethylstilboestrol (DES) Dienoestrol (DE) Hexoestrol (HEX)	< 0.6 < 0.6 < 0.4		
ERM-BB389 ⁽²⁾	Bovine urine	Diethylstilboestrol (DES) Dienoestrol (DE) Hexoestrol (HEX)	1.1 5.5 6.1	\pm \pm \pm	0.5 1.4 0.9

Availability: ⁽¹⁾ Vial containing approximately 0.36 g lyophilised bovine urine corresponding to 5.20 g of fresh bovine urine.

⁽²⁾ Vial containing approximately 0.31 g lyophilised bovine urine corresponding to 5.18 g of fresh bovine urine.

	Description	Substance	Mass fraction in reconstituted sample (μ g/kg)*		
BCR-648	Bovine liver	Clenbuterol		< 0.5	
BCR-649	Bovine liver	Clenbuterol	1.2	\pm	0.3
BCR-474	Bovine liver	17 α -trenbolone		< 0.5	
BCR-475	Bovine liver	17 α -trenbolone	7.6	\pm	2.2
BCR-412	Bovine muscle	Diethylstilbestrol		< 0.1	
BCR-673	Bovine eye	Clenbuterol		< 0.5	
BCR-674	Bovine eye	Clenbuterol	9.4	\pm	1.1

Availability: BCR-648 and -649 are provided in units of 10 g lyophilised bovine liver in vials sealed under argon. BCR-474 and -475 are sold as set and provided in brown glass vials in units of 2.8 g lyophilised liver corresponding to 10 g fresh liver. BCR-412 is provided in brown glass vials as lyophilised bovine muscle in units equivalent to about 5 g of fresh bovine tissue.

BCR-673 and BCR-674 are provided in brown glass vials containing about 0.1 g of material.

	BCR-444 Porcine muscle (blank) (μ g/kg)	ERM-BB130 Pork muscle (μ g/kg)
Chloramphenicol	< 0.2	0.230 \pm 0.021

Availability: BCR-444 is provided in brown glass vials, ERM-BB130 in an amber glass bottle, each containing about 7 g of lyophilised pork muscle tissue.

	Description	Substance	Mass fraction in reconstituted sample (mg/kg)
BCR-695	Pig liver	Chlortetracycline	< 0.004
BCR-696	Pig liver	Chlortetracycline	0.58 ± 0.11
BCR-697	Pig muscle	Chlortetracycline	< 0.006
BCR-706	Pig kidney	Chlortetracycline	< 0.005
BCR-707	Pig kidney	Chlortetracycline	1.30 ± 0.20

Availability: These CRMs are provided in sealed glass vials containing lyophilised tissue equivalent to 5 g of fresh tissue.

	Description	Substance	(µg/kg)
ERM-BB492	Milk powder	Sum of oxytetracycline and 4-epi-oxytetracycline	101 ± 11
ERM-BB493	Milk powder	Sum of oxytetracycline and 4-epi-oxytetracycline	< 5

Availability: ERM-BB492 and ERM-BB493: Units of approximately 5.5 g of spray-dried partially skimmed milk; sold as set of 1 unit ERM-BB492 and ERM-BB493 each.

Substance	BCR-725 Salmon tissue (µg/kg)	
Flumequine	1170	± 210
Oxolinic acid	600	± 100

Availability: BCR-725 is provided in amber glass vials containing 2.2 g of lyophilised salmon tissue material.

	ERM-BB124 Pork muscle
Nitroimidazoles in the reconstituted material	Mass fraction (µg/kg)
Ronidazole (RNZ)	2.09 ± 0.25
Metronidazole (MNZ)	1.93 ± 0.15
2-hydroxymethyl-1-methyl-5-nitroimidazole (HMMNI)	0.69 ± 0.09
Hydroxymetronidazole (MNZOH)	6.2 ± 0.9
Hydroxipronidazole (IPZOH)	1.67 ± 0.12
Dimetridazole (DMZ)	< 0.25

Availability: ERM-BB124 is provided in amber glass bottles containing 10 g of lyophilised pork muscle tissue.

2.2.7 CERTIFIED FOR IDENTITY

Ewes'/Goats' Curd (BCR-599)

BCR-599 consists of a set of two freeze dried curd materials made from a mixture of ewes' and goats' milk, intended to detect adulteration by cows' milk in cheeses made from ewes' milk, goats' milk and mixtures thereof, according to the reference method described in Commission Regulation (EC) No. 1081/96. The 0 % material is not adulterated, the 1 % material is adulterated milk 1 % of cows' milk.

Availability: BCR-599 is available as a set of two brown glass vials containing each about 15 g of lyophilised curd powder under Argon atmosphere.

Peanut Test Material Kit (IRMM-481)

The peanut test material is **not** a reference material, because it could not be tested for homogeneity and stability. The intention is to make a standard peanut matrix available to the research community who may wish to conduct wider studies on similar matrices pre-treated in different ways. At present only the nominal particle size and the pre-treatment applied to the peanuts before milling can be guaranteed as well as the tolerances of weighing of 1.000 ± 0.015 g of each variety in the peanut mixture provided in the kit.

	Peanut variety, origin	Peanut treatment	Nominal net weight of peanut powder	Colour code on cap
IRMM-481a	Runners, Argentina	blanched, strong air-roasting	2 g	blue
IRMM-481b	Common Natal, South-Africa	raw, mild air-roasting	2 g	green
IRMM-481c	Virginia, USA,	blanched, strong oil roasting	2 g	gold
IRMM-481d	Virginia, China	blanched, mild oil-roasting	2 g	red
IRMM-481e	Jumbo Runners, USA,	blanched only	2 g	brown
IRMM-481f	Mixture of 481a to 481e	all above	5 g (1 g of each variety 481a – 481e)	silver

Availability: IRMM-481 consists of a kit with six different vials containing non-salted peanut powders with a nominal particle size from 0.5 to 1.0 mm. Five of the vials are filled with approximately 2 g of each variety. The sixth vial contains a mixture of all five varieties provided in IRMM-481a to 481e. All vials have been filled with argon prior to capping to provide a protective atmosphere.

Calibration kit for the detection of ruminant material in feed by PCR

The calibration kit for ruminant detection by PCR is a certified reference material and should be used to construct calibration curves at low copy number concentration to determine a cut-off value.

	Copy number concentration of the plasmid [cp/ μ L]		
ERM-AD482a	123	\pm	30
ERM-AD482b	32	\pm	7
ERM-AD482c	8	\pm	3

Availability: ERM-AD482 consists of a kit of three plasmid solutions. Each of the three vials contains approximately 1000 μ L of plasmid solution.

Calibration kit for the detection of porcine material in feed by PCR

The calibration kit for porcine detection by PCR is a certified reference material and should be used to construct calibration curves at low copy number concentration to determine a cut-off value.

	Copy number concentration of the plasmid [cp/ μ L]		
ERM-AD483a	126	\pm	30
ERM-AD483b	34	\pm	6
ERM-AD483c	9	\pm	3

Availability: ERM-AD483 consists of a kit of three plasmid solutions. Each of the three vials contains approximately 1000 μ L of plasmid solution.

2.2.8 OTHERS

Substance	ERM-BD273 Toasted bread		
			<u>Mass fraction</u> (ng/g)
Acrylamide	425	±	29

Availability: ERM-BD273 is available in a brown glass vial containing about 30 g of toasted bread powder.

Radionuclide activity concentration at reference date 1 January 2009, 00h00 UTC	IRMM-426 Wild berries (Bq kg ⁻¹)		
¹³⁷ Cs	780	±	70
⁹⁰ Sr	153	±	29
⁴⁰ K	253	±	25

Availability: IRMM-426 is available in bottles containing about 100 g of material.

3 MATERIALS RELATED TO CLINICAL CHEMISTRY

3.1 PURE STANDARDS AND SYNTHETIC MATERIALS

	Description	Purity (%)
BCR-546	Formaldehyde 2,4-dinitrophenylhydrazone	> 99.3
BCR-547	Acetaldehyde 2,4-dinitrophenylhydrazone	98.3 ± 0.5
BCR-548	Acrolein 2,4-dinitrophenylhydrazone	> 97.9
BCR-549	Acetone 2,4-dinitrophenylhydrazone	> 99.6
BCR-550	Glutaraldehyde 2,4-dinitrophenylhydrazone	> 98.1

Availability: Approximately 10 mg of crystals in glass vials.

Compounds	BCR-551 Acetonitrile solution Mass concentration (µg/mL)	BCR-552 Acetonitrile solution (blank) (µg/mL)
Formaldehyde 2,4-Dinitrophenylhydrazone	2.94 ± 0.05	< 0.08
Acetaldehyde 2,4-dinitrophenylhydrazone	4.89 ± 0.07	< 0.05
Acrolein 2,4-dinitrophenylhydrazone	0.483 ± 0.011	< 0.04
Acetone 2,4-dinitrophenylhydrazone	4.96 ± 0.07	< 0.05

Availability: Set BCR-551-2 consists of 4 samples of BCR-551 and 1 sample of BCR-552.

	BCR-553 Glass fibre filters Spiked mass per filter (expressed as µg formaldehyde)	BCR-554 Glass fibre filters Mass per filter (blank) (expressed as µg formaldehyde)
Formaldehyde 2,4-dinitrophenylhydrazone on glass fibre filters	4.96 ± 0.06	< 0.1

Availability: Set BCR-553-4 consists of 2 samples of BCR-553 and 1 sample of BCR-554.

Substance	BCR-555 Chlorinated hydrocarbons on Tenax (ng)		
Dichloromethane	320	±	40
1,1,1-Trichloroethane	370	±	40
Trichloroethylene	390	±	40
Perchloroethylene	327	±	17
Toluene	57	±	7

Availability: Stainless steel tube of 9.0 cm length and 0.25 inches outer diameter containing a single section of 250 mg TENAX GR, charged with 4 chlorinated hydrocarbons and toluene at the levels shown above.

	Description	Latex spheres Parameters of the calibration line
BCR-165	Nominal 2 µm latex (0.02 % solids)	2.223 ± 0.013
BCR-166	Nominal 4.8 µm latex (0.2 % solids)	4.821 ± 0.019
BCR-167	Nominal 9.6 µm latex (1.4 % solids)	9.475 ± 0.018

Availability: Vials containing 2 mL of an aqueous suspension of latex spheres.

	Thyroxine (T ₄)	3,3',5-triiodothyronine (T ₃)
IRMM-468	98.6 ± 0.7	(0.51 ± 0.17)
IRMM-469	(1.50 ± 0.12)	97.1 ± 0.7

Values in brackets are not certified.

Availability: The material consists of an off-white crystalline powder in an amber glass vial sealed under N₂ atmosphere. Each vial contains about 100 mg of the powder.

3.2 MATRIX MATERIALS

3.2.1 CERTIFIED FOR THE HORMONE CONTENT

Cortisol reference panel of fresh frozen human sera ERM-DA451/IFCC

Serum No.	Certified value nmol/L	Uncertainty nmol/L	Serum No.	Certified value nmol/L	Uncertainty nmol/L
1	361	14	18	146	6
2	432	17	19	166	7
3	288	11	20	83	4
4	152	6	21	89	4
5	329	13	22	180	7
6	278	11	23	387	15
7	515	20	24	384	15
8	163	7	25	315	12
9	287	11	26	215	9
10	230	9	27	497	19
11	334	13	28	299	12
12	261	10	29	265	11
13	430	17	30	114	5
14	626	24	31	764	29
15	246	10	32	623	24
16	211	8	33	264	10
17	366	14	34	390	15

Availability: As panel of 34 x 1 mL serum in screw capped cryo-vials.

		Cortisol in human serum (concentration in the reconstituted material ¹⁾)			
		($\mu\text{g/L}$)		(nmol/L)	
ERM-DA192		98.8 277	\pm \pm	2.0 5	
ERM-DA193				273 763	\pm \pm
				6 14	

Availability: In units of lyophilised material of a 1.25 mL portion of serum kept under nitrogen in sealed glass ampoules.

¹⁾ The sample is to be reconstituted with (1.25 \pm 0.01) mL of distilled water.

		Progesterone in human serum (concentration in the reconstituted material ¹⁾)			
		($\mu\text{g/L}$)		(nmol/L)	
BCR-348R		8.5	\pm	0.4	26.9
ERM-DA347		3.19	\pm	0.07	10.13
				\pm	1.2
				0.21	

Availability: In units of lyophilised material of a 1 mL portion of serum kept under nitrogen in sealed glass ampoules.

¹⁾ The sample is to be reconstituted with (1.0 \pm 0.01) mL of distilled water.

		17 β -Estradiol in human serum (concentration in the reconstituted material) Amount-of-substance concentration (nmol/L)			
BCR-576 ¹⁾		0.114	\pm	0.005	
BCR-577 ²⁾		0.689	\pm	0.032	
BCR-578 ²⁾		1.34	\pm	0.07	

Availability: BCR-576, -577, -578 are lyophilised material of a 5 mL (BCR-576) or 1 mL (BCR-577 and BCR-578) portion of serum kept under nitrogen in sealed glass ampoules.

¹⁾ The sample is to be reconstituted with (5.00 \pm 0.05) mL of distilled water.

²⁾ The sample is to be reconstituted with (1.00 \pm 0.01) mL of distilled water.

3.2.2 CERTIFIED FOR THE TOTAL ELEMENT CONTENT AND OTHER PROPERTIES

	Description	Substance	Metal concentrations in the reconstituted material ¹⁾ ($\mu\text{g/L}$)		
ERM-CE195	Lyophilised bovine blood	Pb Cd ²⁾	416 5.06	\pm \pm	9 0.15
ERM-CE196	Lyophilised bovine blood	Pb Cd ²⁾	772 12.33	\pm \pm	11 0.20
BCR-634	Lyophilised human blood	Pb Cd	46 1.4	\pm \pm	5 0.4
BCR-635	Lyophilised human blood	Pb Cd	210 6.6	\pm \pm	24 0.6
BCR-636	Lyophilised human blood	Pb Cd	$0.52 \cdot 10^3$ 11.6	\pm \pm	$0.05 \cdot 10^3$ 0.6

Availability: In units of lyophilised material equivalent to about 5.75 mL of bovine blood with additives kept under nitrogen in rubber stoppered vials.

BCR-634, BCR-635 and BCR-636 are available in lyophilised form in brown glass vials, containing approximately 0.6 g dry matter equivalent to 3.0 mL of fresh whole blood.

¹⁾ The sample is to be reconstituted with (5.00 \pm 0.01) mL water.

²⁾ Recertified by the JRC.

	Description	Substance	Element concentration in the reconstituted material ¹⁾ (mmol/L)		
BCR-304	Lyophilised human serum	Ca Li Mg	2.201 0.985 1.85	\pm \pm \pm	0.019 0.029 0.03

Availability: In units of lyophilised material equivalent to about 5.3 mL of human serum kept under vacuum in rubber stoppered vials.

¹⁾ The sample is to be reconstituted with (5.00 \pm 0.01) mL bi-distilled water.

	Description	Substance	Metal concentrations ($\mu\text{g/L}$)		
BCR-637	Human serum	Al Se Zn	12.5 81 1110	\pm \pm \pm	3.0 7 220
BCR-638	Human serum	Al Se Zn	55 104 1430	\pm \pm \pm	7 7 210
BCR-639	Human serum	Al Se Zn	194 133 2360	\pm \pm \pm	14 12 140

Availability: supplied in frozen form in white plastic vials containing approximately 4.5 mL serum.

Substance	ERM-DB001 Human hair (mg/kg)		
As	0.044	\pm	0.006
Cd	0.125	\pm	0.007
Cu	33	\pm	4
Hg	0.365	\pm	0.028
Pb	2.14	\pm	0.20
Se	3.24	\pm	0.24
Zn,	209	\pm	12

Availability: supplied in amber glass bottle, provided in aluminium sachet, and contains a minimum amount of 3.5 g of a human hair homogeneous powder

3.2.3 CERTIFIED FOR PROTEIN CONTENT

	Description	Mass concentration in the reconstituted material ¹⁾ (g/L)
BCR-393	Lyophilised Apo A I from human serum	1.06 \pm 0.05

Availability: In units of lyophilised material equivalent to about 1.5 mL of Apolipoprotein solution in sealed glass ampoules under nitrogen.

¹⁾ The sample must be reconstituted with 1.0 mL of phosphate buffer.

	Description	Mass concentration in the reconstituted material ¹⁾ (g/L)
BCR-457	Human Thyroglobulin (Tg)	0.324 \pm 0.018

Availability: In units of lyophilised material in sealed glass ampoules under nitrogen.

¹⁾ The sample must be reconstituted with 1.0 mL of distilled water.

	Description	Protein mass per ampoule ¹⁾ (μg)
BCR-486	Purified alphafoetoprotein (AFP)	100 \pm 9

Availability: BCR-486 is provided in sealed glass ampoules. Each sample is in lyophilised form and it contains purified AFP without additives. The protein mass per ampoule is equivalent to $(100 \pm 9) \mu\text{g}$ when the material is reconstituted with 1.0 mL phosphate buffer according to the specified procedure.

¹⁾ Carbohydrate mass of the molecule is not included.

	Description	Protein mass/ampoule
BCR-613	Prostate specific antigen in the reconstituted material	71 \pm 7 μg

Availability: Lyophilised PSA in sealed glass ampoules kept under argon gas.

	Description	HbA _{1c} /Hb _T in reconstituted material (%)
BCR-405 (RM)	Glycated haemoglobin (HbA _{1c}) in human haemolysate	(6.29 ± 0.18)

Value in brackets is not certified.

Availability: Sealed glass ampoules of lyophilised material equivalent to about 0.5 mL of a solution of haemolysate of human erythrocytes kept under carbonmonoxide.

1) Sample to be reconstituted with 1 mL of deionised water and diluted with appropriate 61pprox.61ing reagent, taking into account that the total haemoglobin (Fe₄) concentration is about 0.23 mmol/L (15 g/L).

ERM-DA470k Human Serum Proteins			
Description	Mass concentration ¹⁾ (g/L)	Description	Mass concentration ¹⁾ (g/L)
α_2 macroglobulin (A2M)	1.43 ± 0.06	Haptoglobin (HPT)	0.889 ± 0.021
α_1 acid glycoprotein (AAG)	0.617 ± 0.013	Immunoglobulin A (IgA)	1.80 ± 0.05
α_1 antitrypsin (AAT)	1.12 ± 0.03	Immunoglobulin G (IgG)	9.17 ± 0.18
Albumin (ALB)	37.2 ± 1.2	Immunoglobulin M (IgM)	0.723 ± 0.027
β -2-microglobulin (B2M)	0.00217 ± 0.00007	Transferrin (TRF)	2.36 ± 0.08
Complement 3c (C3c)	1.00 ± 0.04	Transthyretin (TTR)	0.220 ± 0.018
Complement 4 (C4)	0.162 ± 0.007		

Availability: Glass bottle containing lyophilised materials equivalent to about 1 mL of serum with additives kept under nitrogen.

1) Sample to be reconstituted with (1.00 ± 0.01) g water.

	Description	Mass concentration (mg/L)
ERM-DA471/IFCC	Cystatin C	5.48 ± 0.15

Availability: Glass vial containing lyophilised human serum spiked with cystatin C.

	Description	Mass concentration (mg/L)
ERM-DA474/IFCC	C-reactive protein (CRP)	41.2 ± 2.5

Availability: Glass ampoule containing at least 1 mL processed human serum spiked with CRP.

	Description	Mass concentration (mg/L)
ERM-DA476/IFCC	anti-MPO IgG	84 ± 9
ERM-DA483/IFCC	IgG PR3 ANCA	270 ± 29

Availability: Glass vial containing lyophilised material equivalent to 1 mL of serum with additives kept under nitrogen.

	Description	Mass concentration (μ g/L)
ERM-DA480/IFCC	Abeta 42 in cerebrospinal fluid (CSF) (low level)	0.45 ± 0.07
ERM-DA481/IFCC	Abeta 42 in cerebrospinal fluid (CSF) (medium level)	0.72 ± 0.11
ERM-DA482/IFCC	Abeta 42 in cerebrospinal fluid (CSF) (high level)	1.22 ± 0.18

Availability: Microvial containing at least 0.5 mL frozen liquid.

	Description	Absorbance at 540 nm and 10.00 mm pathlength	Mass concentration (mg/L)	Substance concentration (μ mol/L)
BCR-522	Haemiglobincyanide (HiCN) in bovine blood lysate	0.5457 ± 0.0009	800.3 ± 1.3	49.61 ± 0.08

Availability: Bovine blood lysate in sealed brown glass ampoules (10 mL) equivalent to about 800.3 mg/L of haemiglobincyanide.

	Description	Amount-of-substance concentration of creatinine (μ mol/L)
BCR-573	Creatinine in human serum	68.7 ± 1.4
BCR-574	Creatinine in human serum	105.0 ± 1.3
BCR-575	Creatinine in human serum	404.1 ± 7.1

Availability: BCR-573, -574, -575 are the lyophilised form of approximately 1 mL portion of serum, with no additives. The mass of the lyophilised material contained in the ampoule is about 0.09 g.

BCR-573i (RM) Set of creatinine interfering substances

Availability: Consists of three vials with lyophilised solutions
 - 0.025 mg calcium dobesilate / 1.2 mg cefoxitin;
 - 0.044 mg sodium pyruvate;
 - 0.108 mg bilirubin ditaurate.

	Description	Amount-of-substance fraction (mmol/mol)			
IRMM/IFCC-467	Haemoglobin isolated from whole blood	HbA0/(HbA1c + HbA0)		> 976	

Availability: Provided in vials containing approximately 39 mg of a deep frozen buffered solution.

	Description	Amount-of-substance fraction (mmol/mol)				
ERM-AD500/IFCC	Haemoglobin in buffer	HbA1c/(HbA1c + HbA0)	blank	0.0	±	-0.0; +0.4
			Level 1	28.6	±	0.9
			Level 2	57.8	±	1.3
			Level 3	86.7	±	2.2
			Level 4	118.8	±	2.6
			Level 5	153	±	5

Availability: Provided in vials containing about 1mg of a deep frozen buffered solution.

3.2.4 CERTIFIED FOR CATALYTIC ACTIVITY

	Description	Catalytic concentration in reconstituted material Certified value		
		U/L		µkat/L
BCR-647	Human adenosine deaminase (ADA1), from human erythrocytes ²⁾			2.55 ± 0.09
BCR-693	Human pancreatic lipase from pancreatic juice ⁴⁾			28.9 ± 1.2
BCR-694	Human pancreatic lipase (recombinant) ⁴⁾			17.4 ± 1.0
ERM-AD452/IFCC	γ-Glutamyltransferase partially purified, from pig kidney ³⁾	114.1	± 2.4	1.90 ± 0.04
ERM-AD453k/IFCC	Human lactate dehydrogenase isoenzyme 1 (LD1) ³⁾	330	± 7	5.50 ± 0.12
ERM-AD454k/IFCC	Alanine aminotransferase (ALT) ³⁾	103.8	± 2.6	1.73 ± 0.05
ERM-AD455k/IFCC	Creatine kinase isoenzyme MM (CK-MM) ³⁾	314	± 6	5.23 ± 0.10
IRMM/IFCC-456	Human pancreatic α-Amylase ³⁾			9.1 ± 0.3
ERM-AD457/IFCC	Aspartate Transaminase (AST)	104.6	± 2.7	1.74 ± 0.05

Availability: Sealed glass ampoules of lyophilised material equivalent to about 1 mL of a solution of enzyme stabilized by incorporation in serum albumin matrix of bovine (ERM-AD452/IFCC and ERM-AD457/IFCC) origin kept under dry nitrogen. BCR-647 has been stabilised by incorporation in a matrix of 50 mmol/L Tris/HCl buffer 9pH=7.4) and human serum albumin (30 g/L). IRMM/IFCC-456 is provided in sealed ampoules filled with dry nitrogen. Samples are in lyophilised form and equivalent to about 1 mL of a solution of purified enzyme.

ERM-AD453k/IFCC, ERM-AD454k/IFCC and ERM-AD455k are available in glass vials containing lyophilised powder from 1 mL of buffer solution.

BCR-693 and BCR-694 are provided in ampoules of lyophilised material equivalent to about 1 mL of stabilised enzyme.

¹⁾ According to IFCC recommended method at 30 °C.

²⁾ According to method specified in report.

³⁾ According to IFCC recommended method at 37 °C.

⁴⁾ According to method described in certification report at 37 °C.

3.2.5 CERTIFIED FOR DNA SEQUENCE AND MASS CONCENTRATION

CRMs for monitoring leukaemia (ERM-AD623)

SET OF PLASMID SOLUTIONS		
	Number of specific DNA fragments per plasmid	
	Certified value	Uncertainty
<i>BCR-ABL</i> b3a2 transcript	1	negligible
<i>BCR</i> transcript	1	negligible
<i>GUSB</i> transcript	1	negligible
Copy number concentration of the plasmid		
	Certified value [cp/µL]	Uncertainty [cp/µL]
ERM-AD623a	1.08×10^6	0.13×10^6
ERM-AD623b	1.08×10^5	0.11×10^5
ERM-AD623c	1.03×10^4	0.10×10^4
ERM-AD623d	1.02×10^3	0.09×10^3
ERM-AD623e	1.04×10^2	0.10×10^2
ERM-AD623f	10.0	1.5

Availability: ERM-AD623 is a set of six plasmid solutions (a-f). Each of six vials contains approximately 600 µL of plasmid solution.

CRMS for quantifying mass concentration of Lambda DNA

ERM-AD442K Lambda DNA in a solution		
	DNA mass concentration	
	Certified value [ng/µL]	Uncertainty [ng/µL]
Lambda DNA	57.5	1.1
Copy number concentration		
	Indicative value [cp/µL]	Uncertainty [cp/µL]
Lambda DNA	(1.20×10^9)	(0.17×10^9)

Values in brackets are not certified.

3.2.6 OTHERS

	Description	Parameters of the calibration line		
ERM-AD149	Lyophilised rabbit thromboplastin	Slope	1.257	± 0.013
		Intercept	- 0.242	± 0.019

Availability: ERM-AD149 in sealed glass ampoules containing the lyophilised form of a 0.5 mL aliquot of the extract of rabbit brain tissue, without calcium ion added.

	BCR-665 Asbestos fibres in lung tissue (Number of fibres of more than 1 µm in length in million per g dry tissue)	BCR-666 Asbestos fibres in lung tissue (Number of fibres of more than 1 µm in length in million per g dry tissue)
Amosite + crocidolite Anthophyllite	49 ± 16 1.8 ± 0.9	2.3 ± 0.9 5.1 ± 1.5

Availability: Sealed vials with 100 mg of lung tissue.

	IRMM-435 Pharmaceutical glass containers Alkali leaching and release			
Volume of titration solution 0.01 mol/L HCl per 50 mL of leachate	0.38	±	0.04	mL
Sodium release per volume of leachate	1.41	±	0.14	mg/L
Release of Na ₂ O per volume of leachate	1.91	±	0.19	mg/L

Availability: Each unit of IRMM-435 consists of 20 vials of 18.9 mL brimful capacity, made of a semi-durable type of glass, which screw caps.

4 MATERIALS CERTIFIED FOR PHYSICAL PROPERTIES

4.1 CERTIFIED FOR THERMAL PROPERTIES

Resin bonded glass fibre board IRMM-440

The certified thermal conductivity between - 10 °C and + 50 °C is given by

$$\lambda [\text{W}/(\text{m.K})] = 2.93949 \cdot 10^{-2} + \frac{T}{\text{°C}} \cdot 1.060 \cdot 10^{-4} + \frac{T^2}{(\text{°C})^2} \cdot 2.047 \cdot 10^{-7}$$

The uncertainty of the certified thermal conductivity is ± 0.000 28 W/(m.K) at the 95 % confidence level over the range [- 10 °C / + 50 °C].

An indicative value for the thermal conductivity between -170°C and -10°C is given by

$$\lambda [\text{W}/(\text{m.K})] = 2.95 \cdot 10^{-2} + \frac{T}{\text{°C}} \cdot 1.08 \cdot 10^{-4} + \frac{T^2}{(\text{°C})^2} \cdot 2 \cdot 10^{-8}$$

The indicative uncertainty of thermal conductivity is ± 5% at the 95 % confidence level over the range [- 170 °C / - 10 °C].

These equations are valid for a sample of the reference material within the density range [64 kg/m³ – 78 kg/m³].

IRMM-440 is available in the following dimensions: (300 x 300 x 35) mm, (500 x 500 x 35) mm, (600 x 600 x 35) mm, (1000 x 1000 x 35) mm.

Glass-ceramic BCR-724

The certified thermal diffusivity between 298 K and 1025 K is given by

$$\alpha [\text{m}^2/\text{s} \cdot 10^{-6}] = 4.406 - 1.351 \cdot 10^{-2} \cdot T + 2.133 \cdot 10^{-5} \cdot T^2 - 1.541 \cdot 10^{-8} \cdot T^3 + 4.147 \cdot 10^{-12} \cdot T^4$$

The uncertainty of the certified thermal diffusivity is ± 6.1 [%] at the 95 % confidence level over the range from 298 K to 1025 K.

The certified thermal conductivity between 298 K and 1025 K is given by

$$\lambda [\text{W}/(\text{m} \cdot \text{K})] = 2.332 + 515.1 / T$$

The uncertainty of the certified thermal conductivity is ± 6.5 [%] at the 95 % confidence level over the range from 298 K to 1025 K.

Availability: Glass-ceramic cylinders in different shapes (BCR-724A: diameter = 13.0 mm, height > 18 mm; BCR-724B: diameter = 13.9 mm, height > 21 mm; BCR-724D: diameter = 26.9 mm, height > 22 mm).

	Description	Cold filter plugging point CFPP Temperature (°C)			Cloud point CP (°C)		
ERM-FC395k	Gasoil	- 7.9	\pm	1.6	- 7.2	\pm	3.0
ERM-EF002	Biodiesel (B100 Rapeseed)	- 15.2	\pm	1.3	- 4.5	\pm	1.0
ERM-EF004	Diesel (B7)	- 27.9	\pm	2.7	- 6.8	\pm	0.4

Values in brackets are not certified.

Availability: ERM-FC395k consists of a set two amber glass ampoules, each containing 27 mL of gas oil.

ERM-EF002 consists of a set two amber glass ampoules, each containing 27 mL of biodiesel

ERM-EF004 consists of a set two amber glass ampoules, each containing 27 mL of diesel (B7)

Substance	ERM-EF411 Hard coal	ERM-EF412 Brown coal	ERM-EF413 Furnace coke
Gross calorific value (GCV) (MJ/kg)	29.0 \pm 0.4	26.02 \pm 0.22	29.5 \pm 0.4
Net calorific value (NCV) (MJ/kg)	28.0 \pm 0.4	24.98 \pm 0.25	29.4 \pm 0.5
Volatile matter (g/100 g)	38.1 \pm 1.0	50.1 \pm 0.7	
Ash (g/100 g)	8.3 \pm 0.7	4.11 \pm 0.23	
C (g/100 g)	71.4 \pm 1.0	66.2 \pm 0.7	87.8 \pm 1.9
Ca (g/kg)		9.8 \pm 0.4	2.92 \pm 0.22
Cd (mg/kg)		(0.012 \pm 0.004)	
Cl (mg/kg)	99 \pm 19		
Co (mg/kg)	(3.5 \pm 0.8)		(350 \pm 130)
Cu (mg/kg)		(0.68 \pm 0.22)	
H (g/100 g)	4.80 \pm 0.14	4.88 \pm 0.15	
Hg (mg/kg)	(0.079 \pm 0.015)	0.071 \pm 0.011	
K (mg/kg)		229 \pm 18	
Mg (g/kg)		(3.73 \pm 0.16)	
Mn (mg/kg)		48.6 \pm 1.9	
N (g/100 g)	1.43 \pm 0.10	0.74 \pm 0.06	1.10 \pm 0.07
Na (g/kg)		2.20 \pm 0.12	0.64 \pm 0.07
Pb (mg/kg)		(0.25 \pm 0.05)	(8.41 \pm 1.6)
S (g/100 g)	0.598 \pm 0.017	0.360 \pm 0.023	0.58 \pm 0.12
Sb (mg/kg)	(1.5 \pm 0.4)	(0.024 \pm 0.004)	
Se (mg/kg)	5.1 \pm 1.0	0.96 \pm 0.14	1.33 \pm 0.26
Tl (mg/kg)	(0.24 \pm 0.07)		
V (mg/kg)	(22 \pm 7)	0.57 \pm 0.04	
Zn (mg/kg)	(13 \pm 4)	(0.99 \pm 0.18)	16.0 \pm 2.5

Values in brackets are not certified.

Availability: ERM-EF411, ERM-EF412 and ERM-EF413 are available in units of about 50 g in aluminium-laminated sachets.

4.2 CERTIFIED FOR MECHANICAL PROPERTIES

Shear testing of powders BCR-116

The flow of powders or granulated materials under the force of gravity affects the design and operation of silos used for their bulk storage. The European Federation of Chemical Engineering (EFCE) therefore developed a test method, based on the Jenike Shear Cell, to determine the shear strength of powders under different compaction and loading conditions. The complexity of this method is such that errors due to poor technique can easily arise. A reference material has therefore been produced with which laboratories can verify both their equipment and experimental technique.

Essentially the EFCE method consists of using a known load to compact a powder sample into a cylindrical Jenike Cell composed of two metal rings one upon the other. Having reached critical compaction of the powder, the steady state force necessary to displace the upper ring horizontally with respect to the lower one is determined with the compaction load still applied. Having established steady state shear the normal load on the powder is then reduced and the horizontal force necessary to continue to shear the powder is determined. It is this variation of the shear force as a function of the reduced normal load for a given compaction load which characterizes the powder.

The reference material consists of 3 kg of limestone powder packed in a polyethylene jar. It is accompanied by a certificate giving shear stress as a function of normal applied stress for four different powder compaction stresses.

Creep BCR-425

Creep is the progressive deformation of a material under load. Metallic materials are usually tested at elevated temperatures for periods of 1 000 to 100 000 hours by surrounding them with a suitable constant temperature furnace. The need to operate under such conditions gives rise to problems of alignment, strain measurement, temperature measurement etc. which can result in considerable differences in results between laboratories.

BCR-425 was therefore developed to allow laboratories to validate their creep testing rigs and procedures as a whole within a reasonable time (500 to 600 hours) using a material whose properties are sensitive to test conditions.

The test piece is delivered in the form of a bar of 14 mm diameter and 500 mm length which must be machined by the laboratory to the required shape and size.

At a test temperature of 600 °C, using an applied stress of 160 Mpa the certified properties are as follows:

Creep rate at 400 hours of	(72	±	5)	10^{-6} h^{-1}
Time to a creep strain of 2 %	(278	±	16)	h
Time to a creep strain of 4 %	(557	±	30)	h

Nimonic 75 for ambient temperature tensile properties BCR-661

Property	Certified value
0.2 % proof stress $R_{p0.2}$	(300 ± 8) Mpa
0.5 % proof stress $R_{p0.5}$	(318 ± 7) Mpa
Ultimate tensile strength R_m	(750 ± 14) Mpa
Elongation to fracture A	(40.9 ± 0.9) %
Reduction in area Z	(60 ± 4) %

Availability: BCR-661B is supplied as 1 bar of 500 mm long, sufficient for the manufacture of three test-pieces of about 150 mm long.

Scratch testing BCR-692

Failure event	Critical load	
	Certified value (N)	Uncertainty (N)
Forward chevron cracks at the borders of the scratch track. <i>(Lc₁ shall be taken at the closest end of the event to the scratch track start).</i>	(Lc ₁) 13.6	1.8
Forward chevron cracks at the borders of the scratch track, with local interfacial spallation or with gross interfacial spallation. <i>(Lc₂ shall be taken at the failure event that occurs first and at the closest end of the event to the scratch track start).</i>	(Lc ₂) 17.0	2.1
Gross interfacial shell-shaped spallation. <i>(Lc₃ shall be taken at the first point where the substrate can be seen at the centre of the track in a crescent that goes completely through the track).</i>	(Lc ₃) 28	2.9

Availability: The reference samples are (30x30x5) mm steel coupons coated with a diamond-like carbon coating (DLC) applied by plasma-assisted chemical vapour deposition. The coupons are distributed in a reusable plastic box containing desiccant.

Impact toughness Charpy specimens

Impact toughness is the ability of a material to resist fracture under the effect of shock loading. It is determined by means of a conventional test defined in ISO and CEN Standards.

The test is sensitive to many sources of errors and therefore already 40 years ago, ASTM specification required testing machines to be verified periodically by reference specimens. The recent CEN standard (EN 10045-2: 1993) specifies that impact toughness machines should be certified by using the BCR certified reference Charpy specimens or other specimens traceable to the latter.

The reference specimens described here are supplied by groups of five. Their use must be done in accordance with the instructions given in the certification report, in particular with regard to degreasing. The fracture energy is certified for the CEN striker (knife of 2 mm radius).

Measurements generally need to be performed at 20°C, with the exception of ERM-FA013 where one batch has certified values at 0°C and 20°C to avoid jamming. We will deliver by default a batch certified by 20°C unless the order explicitly mentions the value at 0°C.

For each energy level there are several batches of slightly different values. The following table gives the nominal values, the certificate gives the actual values.

ERM-FA013 – Low Energy 20°C	20-30 J
ERM-FA013 – Low Energy 0°C	20-30 J
ERM-FA015	80 J
ERM-FA016	120 J
ERM-FA415	150J

4.3 CERTIFIED FOR MORPHOLOGICAL PROPERTIES

Particle size distributions					
	Form of Quartz	Certified Property	Size Range (μm)		Unit Size (g)
BCR-066	Powder	Stokes' diameter	0.35	-	3.50
BCR-067	Powder	Stokes' diameter	2.40	-	32.00
BCR-068	Sand	Volume diameter	160.0	-	630.0
BCR-069	Powder	Stokes' diameter	14.0	-	90.0
BCR-070	Powder	Stokes' diameter	1.20	-	20.00

Particle size distributions					
	Form of Quartz	Certified Property	Size Range (μm)		Unit Size (g)
BCR-130	Powder	Volume diameter	50	-	220
BCR-131	Powder	Volume diameter	480	-	1800
BCR-132	Gravel	Volume diameter	1400	-	5000

Particle size distributions					
Corundum	Certified Property			Size Range (μm)	Unit Size (g)
ERM-FD066	Volume-weighted equivalent diameter by laser diffraction, Mie theory, wet dispersion		1.44	-	7.45
	Number-weighted equivalent diameter by scanning electron microscopy (SEM)		1.07	-	5.1
ERM-FD069	Volume-weighted equivalent diameter by laser diffraction, Fraunhofer approximation		13.9	-	79.8
	Volume-weighted equivalent diameter by laser diffraction, Mie theory		15.0	-	82
	Number-weighted area-equivalent diameter by optical microscopy		12.4	-	46

	Description	Specific Surface Area ($\text{m}^2 \cdot \text{g}^{-1}$)	Unit Size (g)
BCR-169	Alpha alumina	0.104	60
BCR-170	Alpha alumina	1.05	60
BCR-171	Alumina	2.95	50
BCR-172	Quartz	2.56	10
BCR-173	Titanium dioxide	8.23	46
BCR-175	Tungsten	0.18	200

Mullite ($3\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2$) BCR-301 (RM)

High crystallinity.

Vitreous phase 0.03 g/g. No other phase detected.

Impurities in g/kg:

Fe_2O_3	< 2	Na_2O	< 1
CaO	< 1.2	K_2O	< 0.5
MgO	< 0.5	TiO_2	< 0.5

Mullite BCR-301 (RM)			
Reflection	Lattice spacing (nm)	Relative intensity	
[110]	0.538 2		0.50
[210]	0.339 0		1
[220]	0.269 5		0.40
[121]	0.220 6		0.59
[331]	0.152 4		0.36

	Description	Micropore volume (cm^3 / g^1)	Median micropore width (nm)
BCR-704	Faujasite type zeolite	0.205	± 0.019
BCR-705	Linde type A zeolite	0.181	± 0.020

Availability: Glass bottle containing 10 g of pellets.

Reference material for depth profiling by ion beam sputtering BCR-261T

In order to achieve the accuracy required when measuring compositional depth profiles using ion beam sputtering in association with Auger Electron Spectroscopy, a reference material of accurately known thickness on a stable substrate is required. BCR-261T is a tantalum pentoxide on tantalum foil reference material existing in two nominal thicknesses of 30 and 100 nm.

	Nominal thickness (nm)	Certified values		
		10^{21} oxygen atoms/m ²	oxide thickness ratio	
BCR-261T	(30) (100)	1.72 ± 0.07 5.40 ± 0.12	0.321 ± 0.013	

Values in brackets are not certified.

Availability: Four rectangular foils of 5 x 10 mm of each oxide thickness.

Colloidal Silica in water ERM-FD100	Equivalent spherical diameter	
	Certified value (nm)	Uncertainty (nm)
Intensity-weighted harmonic mean diameter (DLS)	19.0	0.6
Intensity-based modal Stokes diameter (CLS)	20.1	1.3
Number-based modal diameter (TEM/SEM)	19.4	1.3
Intensity-weighted mean diameter (SAXS)	21.8	0.7
Equivalent spherical diameter, volume-weighted mean (SAXS)	(20.4)	(1.6)
Zeta Potential	(- 43.0 mV)	(22 mV)

Values in brackets are not certified.

Availability: ERM-FD100 is available in 10 mL pre-scored amber glass ampoules containing approximately 9 mL of suspension.

Colloidal Silica in aqueous solution ERM-FD304	Equivalent spherical diameter	
	Certified value (nm)	Uncertainty (nm)
Scattering intensity-weighted harmonic mean diameter (DLS)	42.1	0.6
Extinction intensity-based modal Stokes (CLS)	33.0	3.0
Number-based modal diameter (TEM/SEM)	(27.8)	(1.5)

Values in brackets are not certified.

Availability: ERM-FD304 is available in 10 mL pre-scored amber glass ampoules containing approximately 9 mL of suspension.

Mixture of Silica Nanoparticles in aqueous solution: ERM-FD102	Equivalent diameter			
	Size class A		Size class B	
	Certified value (nm)	Uncertainty (nm)	Certified value (nm)	Uncertainty (nm)
Scattering intensity-weighted arithmetic mean hydromatic diameter (DLS)	17.8	1.5	88.5	2.2
Extinction intensity-weighted modal Stokes'diameter (CLS)	23.9	2.0	88	7
Number-weighted modal area-equivalent diameter (TEM and SEM)	18.2	1.6	84.0	2.1
Number-weighted median area-equivalent diameter (TEM and SEM)	18.3	1.7	83.3	2.3
Number-weighted mean hydrodynamic diameter (PTA)	-	-	(82)	(4)
Number-weighted modal maximum particle height (ATM)	(16.9)	(1.8)	(80)	(6)

Values in brackets are not certified. More indicative & information values are given on the certificate

Availability: ERM-FD102 is available in 10 mL pre-scored amber glass ampoules containing approximately 9 mL of suspension.

Mixture of Silica Nanoparticles in aqueous solution: ERM-FD101b	Size distribution parameter: Weighting / Averaging	Certified value [nm]	Uncertainty [nm]
Hydrodynamic diameter from DLS (cumulants method)	Scattered light intensity-weighted / harmonic mean	89.5	2.3
Hydrodynamic diameter from DLS (distribution calculation algorithms)	Scattered light intensity-weighted / mean (arithmetic, harmonic, geometric) and modal	93	4
Hydrodynamic diameter from PTA	Number-weighted / modal	82	4
	Number-weighted / arithmetic mean	87	4
	Number-weighted / median	82	4
Stokes diameter from CLS ¹ (turbidimetry)	Light extinction-weighted / modal	87	8
Area-equivalent diameter from EM	Number-weighted / modal	83.7	2.2
	Number-weighted / median	83.5	2.2
Mean particle diameter from SAXS (model fitting)	Scattered X-ray intensity-weighted / modal	82.5	1.8
	Volume-weighted / modal	81.7	1.8
	Number-weighted / modal	80.9	1.7
Mean particle diameter from SAXS (Guinier approximation)	(Volume)-weighted / mean	(87)	(6)

Values in brackets are not certified. More material information is given on the certificate

Availability: ERM-FD101b is available in 10 mL pre-scored amber glass ampoules containing approximately 9 mL of suspension.

5 MATERIALS RELATED TO INDUSTRIAL APPLICATIONS

5.1 CERTIFIED FOR COMPOSITION

Certified Parameter	IRMM-441 n-Heptane (g/kg)	IRMM-442 Isooctane (g/kg)
Isooctane, purity by difference		999.85 ± 0.05
n-Heptane, purity by difference	999.85 ± 0.05	
<u>Impurities</u>		
Total organics (other than isoctane)		0.11 ± 0.04
Total organics (other than n-Heptane)	0.12 ± 0.05	
Isooctane	0.07 ± 0.02	
n-Heptane		0.02 ± 0.02
Water	0.03 ± 0.02	0.04 ± 0.02

Availability: IRMM-441 and -442 are supplied in ampoules of 100 mL.

ERM-EF001 – Biodiesel certified for selected parameters specified in EN 14214	Unit	Certified values		
Ester content	% (m/m)	98.9	±	1.7
Linolenic acid methyl ester content	% (m/m)	8.82	±	0.16
Monoglyceride content	% (m/m)	0.65	±	0.04
Diglyceride content	% (m/m)	0.136	±	0.025
Triglyceride content	% (m/m)	<0.1		
Total glycerol content	% (m/m)	0.187	±	0.009
Water content	% (m/m)	0.0205	±	0.0024
Density (at 15 °C)	kg/m ³	883.20	±	0.04
Viscosity (at 40 °C)	mm ² /s	4.465	±	0.005
Oxidation stability (at 110 °C)	h	9.8	±	0.5
Iodine value	g iodine/100 g	112	±	4
Flash point	°C	181	±	14

Availability: Vials containing about 27 mL of biodiesel.

ERM-EF003 - selected parameters diesel (B7)	Certified value ⁸⁾	Uncertainty ⁹⁾	Unit
Fatty acid methyl ester content ¹⁾	6.88	0.17	% (V/V) ⁶⁾
Mono-aromatic hydrocarbon content ²⁾	18.8	0.7	% (m/m) ⁷⁾
-aromatic hydrocarbon content ²⁾	1.84	0.19	% (m/m) ⁷⁾
Polycyclic aromatic hydrocarbon content ²⁾	2.01	0.25	% (m/m) ⁷⁾
Total aromatic hydrocarbon content ²⁾	20.8	0.9	% (m/m) ⁷⁾
Density (at 15.0 °C) ³⁾	837.23	0.07	kg/m ³
Kinematic viscosity (at 40.0 °C) ⁴⁾	2.892	0.012	mm ² /s
Lubricity ⁵⁾	220	60	µm

1) As defined by EN 14078:2014; 2) As defined by EN 12916:2016; 3) As defined by EN ISO 12185:1996; 4) As defined by EN ISO 3104:1996

5) As defined by EN 12156-1:2016.

6) As called in EN 14078:2014, which is equivalent to 10⁻² mL/mL; 7) As called in EN 12916:2006, which is equivalent to 10⁻² g/g.

8) Certified values are values that fulfil the highest standards of accuracy. The given values represent the unweighted mean value of the means of accepted sets of data, each set being obtained in a different laboratory using one of the above mentioned methods for determination. The certified values and their uncertainties are traceable to the International System of Units (SI).

9) The uncertainty of the certified value is the expanded uncertainty with a coverage factor *k* = 2 corresponding to a level of confidence of about 95 % estimated in accordance with ISO/IEC Guide 98-3, Guide to the Expression of Uncertainty in Measurement (GUM:1995), ISO, 2008.

Availability: Vials containing about 27 mL of diesel (B7).

	Description	Cold filter plugging point CFPP Temperature (°C)	Cloud point CP (°C)
ERM-EF002	Biodiesel (B100 Rapeseed)	- 15.2 ± 1.3	- 4.5 ± 1.0

Availability: set two ERM-EF002 consists of a amber glass ampoules, each containing 27 mL of biodiesel.

	Description	Cold filter plugging point CFPP Temperature (°C)	Cloud point CP (°C)
ERM-EF004	Diesel (B7)	- 27.9 ± 2.7	- 6.8 ± 0.4

Availability: ERM-EF004 consists of a set two amber glass each containing 27 mL of diesel (B7)

	Description	Substance	Certified values			
BCR-032	Moroccan Phosphate rock	CaO	518	±	4	g/kg
		P ₂ O ₅	329.8	±	1.7	g/kg
		CO ₂	51.0	±	0.8	g/kg
		F	40.4	±	0.6	g/kg
		SiO ₂	20.9	±	1.2	g/kg
		SO ₃	18.4	±	0.8	g/kg
		Al ₂ O ₃	5.5	±	0.6	g/kg
		MgO	4.0	±	0.1	g/kg
		Fe ₂ O ₃	2.3	±	0.1	g/kg
		As	(9.5	±	0.5	mg/kg)
		B	(22.6	±	2.2	mg/kg)
		Cd	(20.8	±	0.7	mg/kg)
		Cr	(257	±	16	mg/kg)
		Co	(0.59	±	0.06	mg/kg)
		Cu	(33.7	±	1.4	mg/kg)
		Hg	(0.055	±	0.011	mg/kg)
		Mn	(18.8	±	1.3	mg/kg)
		Ni	(34.6	±	1.9	mg/kg)
		Ti	(171	±	10	mg/kg)
		V	(153	±	7	mg/kg)
		Zn	(253	±	6	mg/kg)

Values in brackets are not certified.

Availability: Units of about 100 g in the form of fine powder.

	Description	Substance	Certified values (g/kg)		
BCR-010	Tin ore	Sn	765.9	±	1.2

Availability: This CRM is contained in brown glass bottles. The approximate quantity per unit is 225 g.

	Description	Substance	Certified values (g/kg)		
BCR-033	Super-phosphate	P ₂ O ₅	193.4	±	1.2
		SO ₄	428.0	±	4.1
		CaO	314.8	±	3.1
		SiO ₂	29.2	±	1.2
		F	16.5	±	0.5
		Al ₂ O ₃	11.0	±	0.8
		Fe ₂ O ₃	4.0	±	0.2
		MgO	2.1	±	0.2
		K	502.5	±	1.1
		Cl	478.0	±	0.9
BCR-113	Potassium Chloride	Na	15.3	±	0.2
		Ca	1.03	±	0.04
		Mg	0.24	±	0.01
		water soluble K	501.3	±	0.7
		K	418.0	±	0.9
		SO ₄	533	±	2
BCR-114	Potassium Sulphate	Cl	18.5	±	0.1
		Na	11.0	±	0.1
		Ca	9.4	±	0.2
		Mg	0.74	±	0.01
		water soluble K	417.6	±	0.8
		K	418.0	±	0.9
BCR-178	Calcium Ammonium Nitrate	SO ₄	533	±	2
		Cl	18.5	±	0.1
		Na	11.0	±	0.1
		Ca	9.4	±	0.2
		Mg	0.74	±	0.01
BCR-179	Urea	water soluble K	417.6	±	0.8
		NH ₄ – N	130.44	±	0.32
		NO ₃ - N	130.15	±	0.57
		total – N	260.19	±	0.54
		Ca	88.82	±	0.27
		total – N	465.4	±	0.8
		Uric – n	460.9	±	0.9
		Biuret	10.37	±	0.11

Availability: Units of about 100 g in the form of fine powder.

Substance	BCR-126A			BCR-126B		
	Lead crystal glass (cg/g)			Lead crystal glass (cg/g)		
SiO ₂	57.80	±	0.11	57.87	±	0.18
PbO	23.98	±	0.06	24.09	±	0.13
K ₂ O	9.99	±	0.07	9.98	±	0.13
Al ₂ O ₃	0.126	±	0.013	0.137	±	0.027
Fe ₂ O ₃	0.005 5	±	0.001 2	0.006 0	±	0.001 6
Sb ₂ O ₃	0.291	±	0.012	0.291	±	0.019
BaO	1.053	±	0.030	1.03	±	0.05
CaO	1.033	±	0.030	1.01	±	0.05
MgO	0.512	±	0.013	0.513	±	0.023
ZnO	1.01	±	0.04	1.00	±	0.04
Na ₂ O	3.57	±	0.07	3.59	±	0.11
Li ₂ O	0.494	±	0.016	0.487	±	0.015
Density at 20 °C	2.990 5	±	0.001 6 g/cm ³	2.994 7	±	0.002 6 g/cm ³
Refractive index n _D ^{20 °C} at 589 nm	1.559 67	±	0.000 22	1.560 04	±	0.000 18

Availability: In the form of square plates (100 × 100 mm) and 10 mm thickness.

5.2 CERTIFIED FOR TRACE ELEMENT CONTENT

	Material	Certified value	Form	Unit
IRMM-521	Ni	< 0.1 mg Co kg ⁻¹	B: 0.5 mm wire R: 0.1 mm foil	100 cm ² (1.8 g) 75 cm ² (6.7 g)
IRMM-522	Cu	< 0.05 mg Co kg ⁻¹ 0.95 ± 0.04 mg Ag kg ⁻¹	A: 0.1 mm foil B: 1.0 mm foil C: 0.5 mm wire D: 1.0 mm wire	100 cm ² (8.9 g) 20 cm ² (17.8 g) 1 m (1.8 g) 1 m (7.0 g)
IRMM-523	Al	< 0.1 mg Na kg ⁻¹	A: 0.1 mm foil B: 1.0 mm foil C: 1.0 mm wire	100 cm ² (2.7 g) 20 cm ² (5.4 g) 1 m (2.1 g)
IRMM-524	Fe	< 0.05 mg Co kg ⁻¹ < 0.1 mg Mn kg ⁻¹	A: 0.1 mm foil B: 0.5 mm wire	100 cm ² (7.9 g) 1 m (1.6 g)
IRMM-525	Nb	19.6 ± 1.8 mg Ta kg ⁻¹	A: 0.02 mm foil B: 0.1 mm foil C: 0.5 mm wire	20 cm ² (0.3 g) 20 cm ² (1.7 g) 1 m (1.7 g)
IRMM-526	Nb	0.30 ± 0.09 mg Ta kg ⁻¹	A: 0.02 mm foil B: 0.1 mm foil C: 0.5 mm wire	20 cm ² (0.3 g) 20 cm ² (1.7 g) 1 m (1.7 g)
IRMM-529	Rh	< 5 g Pt kg ⁻¹ 26.0 ± 0.6 g Ir kg ⁻¹	0.05 mm foil	20 cm ² (1.2 g)
IRMM-531	Ti	< 0.1 mg Sc kg ⁻¹	A: 0.1 mm foil B: 0.5 mm foil C: 0.5 mm wire	100 cm ² (4.5 g) 20 cm ² (4.5 g) 1 m (1 g)
IRMM-527R	Al – 0.1% Co	1.001 ± 0.024 g Co kg ⁻¹	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm ² (2.7 g) 1 m (0.5 g) 1 m (2.1 g)
IRMM-528R	Al – 1.0% Co	10.02 ± 0.23 g Co kg ⁻¹	A: 0.1 mm foil C: 1.0 mm wire	100 cm ² (2.7 g) 1 m (2.1 g)
IRMM-530R	Al – 0.1% Au	1.003 ± 0.012 g Au kg ⁻¹	A: 0.1 mm foil C: 1.0 mm wire	100 cm ² (2.7 g) 1 m (2.1 g)
ERM-EB530	Al – 0.1% Au	1.005 ± 0.007 g Au kg ⁻¹	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	50 cm ² (1.4 g) 1 m (0.5 g) 1 m (2.1 g)
IRMM-532	Al – 0.01% Co	0.100 0 ± 0.002 5 g Co kg ⁻¹	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm ² (2.7 g) 1 m (0.5 g) 1 m (2.1 g)
IRMM-533	Al – 0.1% Ag	0.996 ± 0.017 g Ag kg ⁻¹	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm ² (2.7 g) 1 m (0.5 g) 1 m (2.1 g)
IRMM-534	Al – 2.0% Sc	19.95 ± 0.20 g Sc kg ⁻¹	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm ² (2.7 g) 1 m (0.5 g) 1 m (2.1 g)

	Description	Substance	Certified value (mg/kg)	Form, dimensions ¹⁾ and availability
BCR-017A	Copper	P	6.85 ± 0.29	
BCR-017B		S	10.4 ± 0.6	A: Ø 42 mm, h 30 mm
BCR-022A	Copper (electrolytic tough pitch)	O	138 ± 7	B: chips (bottle with 50 g) Ø 26 mm, h 9 mm
BCR-022B	Copper (electrolytic tough pitch)	O	138 ± 7	Ø 9 mm, h 50 mm
BCR-054R	Copper (low oxygen)	O	0.47 ± 0.07	Ø 7 mm, h 50 mm
BCR-058	Copper (continuous cast)	O	390 ± 24	Ø 7 mm, h 50 mm
BCR-024B	Titanium	N	117 ± 13	
BCR-024C		O	608 ± 23	B: 25 cubes of 0.4 g
BCR-059A	Titanium alloy Ti6Al4V	O	1750 ± 70	C: 25 cubes of 0.2 g
BCR-059B		N	172 ± 27	A: Ø 26 mm, h 9 mm
BCR-318	Titanium	H	12.2 ± 0.6	B: 25 cubes of 0.2 g Ø 7 mm, h 1 mm (bottle with 74pprox.. 100 discs)
BCR-275	Zirconium alloy Zircaloy-4	O	1670 ± 50	Ø 13 mm, h 1 mm
		N	39.0 ± 1.7	(bottle with 10 discs)
		C	113 ± 4	
BCR-276	Zirconium alloy Zircaloy-4	O	1540 ± 80	Ø 4.5 mm, h 2 mm
		N	41 ± 9	(bottle with 74pprox.. 100 discs)
		C	108 ± 11	
BCR-102	Tungsten carbide powder	O	185 ± 4	Bottles containing 2 – 3 g powder, sealed under argon in an aluminium container

1) Ø = diameter, h = height

Substance	BCR-286 Electrolytically refined lead (mg/kg)	BCR-287 Thermally refined lead (mg/kg)		BCR-288 Lead with added impurities (mg/kg)
Ag		15.20 ± 0.21		
As		67.3 ± 1.1		55.7 ± 1.6
Bi	21.5 ± 0.5	0.356 ± 0.024	215.8 ± 2.4	
Cd		0.98 ± 0.05	33.3 ± 0.9	
Cu		0.040 ± 0.015	19.3 ± 0.4	
Sb	0.099 ± 0.021			< 0.2
Se				32.8 ± 1.3
Te				2.26 ± 0.08
Tl	2.47 ± 0.07	0.73 ± 0.04	8.2 ± 0.4	
Zn	< 0.1	< 0.1		

Availability: CRMs are available as follows: BCR-286A and -287A: blocks of 60 × 60 × 12 mm, BCR-286B, -287B and -288B: chips in bottles containing about 160 g.

Substance	BCR-321 Unalloyed zinc (mg/kg)	ERM-EB322 Unalloyed zinc (mg/kg)	ERM-EB323 Unalloyed zinc (mg/kg)	ERM-EB324 Unalloyed zinc (mg/kg)	ERM-EB325 Unalloyed zinc (mg/kg)	BCR-326 Unalloyed zinc (mg/kg)	BCR-327 Unalloyed zinc (mg/kg)
Al	< 0.7						
Cd	(0.23 ± 0.03)	15.08 ± 0.30	6.51 ± 0.21	48.6 ± 1.1	94.7 ± 2.5	203.0 ± 2.0	301.4 ± 2.3
Cu	(0.97 ± 0.05)	5.89 ± 0.15	18.9 ± 0.4	9.87 ± 0.18	47.5 ± 2.0	104.8 ± 2.7	(0.56 ± 0.11)
Fe	(2.22 ± 0.14)	19.1 ± 0.8	11.3 ± 0.7	58.5 ± 1.6	56.1 ± 3.3	264.8 ± 2.1	144.0 ± 1.3
In	< 0.2						
Pb	4.85 ± 0.20	15.0 ± 0.5	48.6 ± 0.9	26.1 ± 0.5	142 ± 9	307.0 ± 1.6	409.4 ± 2.3
Sn	< 0.5	5.6 ± 0.6	18.7 ± 0.7	9.8 ± 0.5	46.1 ± 2.0		
Tl	0.78 ± 0.10	5.28 ± 0.30	10.8 ± 0.5	19.9 ± 0.5	36.8 ± 1.2		

Values in brackets are not certified.

Availability: Discs of 80 mm diameter and 20 mm thickness (BCR-321, -326, -327) and 60 mm diameter, 30 mm thickness, respectively (ERM-EB322, EB323, EB324 and EB325).

Substance	BCR-351 ZnAl4 (mg/kg)	BCR-352 ZnAl4 (mg/kg)	BCR-353 ZnAl4 (mg/kg)	BCR-354 ZnAl4 (mg/kg)	BCR-355 ZnAl4 (mg/kg)
Al	[43.55 ± 0.11] × 10 ³	[41.50 ± 0.10] × 10 ³	[39.5 ± 0.4] × 10 ³	[37.27 ± 0.16] × 10 ³	[34.43 ± 0.13] × 10 ³
Cd	(0.21 ± 0.03)	2.88 ± 0.12	10.44 ± 0.16	29.7 ± 0.4	58.1 ± 0.4
Cu	12.13 ± 0.15	31.26 ± 0.29	100.0 ± 0.8	312.3 ± 2.5	1035 ± 6
In	< 0.2	3.02 ± 0.28	2.55 ± 0.23	9.8 ± 0.9	24.6 ± 1.4
Mg	131.0 ± 0.9	283.0 ± 1.8	452.5 ± 2.4	602 ± 5	786 ± 6
Ni	(1.9 ± 0.6)	6.74 ± 0.16		83.1 ± 2.9	268 ± 8
Pb	4.50 ± 0.20	(6.4 ± 1.6)	24.4 ± 1.3	30.8 ± 1.2	56.9 ± 1.9
Sn	< 1	3.0 ± 0.7	5.6 ± 0.6	14.1 ± 1.1	29.1 ± 2.0
Tl	0.74 ± 0.06	3.2 ± 0.4	3.95 ± 0.22	11.01 ± 0.20	23.25 ± 0.28

Values in brackets () are not certified.

Availability: Discs of 80 mm diameter and 20 mm thickness.

	BCR-356 ZnAl4Cu1 (mg/kg)	BCR-357 ZnAl4Cu1 (mg/kg)	BCR-360 ZnAl4Cu1 (mg/kg)	BCR-361 ZnAl4Cu1 (mg/kg)
Al	[44.34 ± 0.11] × 10 ³	[42.27 ± 0.11] × 10 ³	[34.27 ± 0.12] × 10 ³	[40.68 ± 0.19] × 10 ³
Cd	0.73 ± 0.09	2.83 ± 0.10	59.5 ± 0.6	(0.80 ± 0.17)
Cu	[3.944 ± 0.022] × 10 ³	[5.849 ± 0.021] × 10 ³	[12.34 ± 0.05] × 10 ³	[7.98 ± 0.04] × 10 ³
Fe	31.5 ± 0.6	25.7 ± 1.2		10.34 ± 0.26
In	< 0.2	3.30 ± 0.14	29.8 ± 0.6	(< 0.2)
Mg	132.3 ± 1.8	273 ± 4	705 ± 5	
Ni	3.43 ± 0.19	9.82 ± 0.25	267 ± 8	
Pb	9.87 ± 0.23	13.8 ± 0.6	73.9 ± 1.4	5.31 ± 0.20
Sn	(0.32 ± 0.16)	3.51 ± 0.14	33.0 ± 0.8	46.3 ± 0.9
Tl	0.79 ± 0.05	2.76 ± 0.05	25.9 ± 0.7	37.4 ± 0.5

Values in brackets () are not certified.

Availability: Discs of 80 mm diameter and 20 mm thickness.

Substance	BCR-089 TiAl6V4 (mg/kg)	BCR-090 Titanium with added impurities (mg/kg)
Al	59700 ± 400	
C	38 ± 10	
B		28.2 ± 1.4
Co		501 ± 14
Cr	122 ± 6	533 ± 11
Cu	10.3 ± 1.2	513 ± 9
Fe	515 ± 16	563 ± 16
H	31 ± 5	
Hf	0.126 ± 0.011	
Mn	4.2 ± 0.6	314 ± 10
Mo	15.2 ± 1.8	488 ± 11
N	212 ± 33	
Nb		(492 ± 26)
Ni	106 ± 7	667 ± 7
O	1660 ± 60	
Sb	1.94 ± 0.12	
Sn	10.4 ± 1.7	(710 ± 50)
Ta	0.30 ± 0.09	
V	39760 ± 290	
W	1.6 ± 0.4	(500 ± 40)
Zr	2.8 ± 0.6	(436 ± 13)

Values in brackets are not certified.

Availability: BCR-089: Cylinder of 40 mm Ø and 20 mm height. BCR-090A: Cylinder of 40 mm Ø and 20 mm height. BCR-090B: Cubes of about 0.2 g in bottles containing approximately 25 g.

Substance	BCR-098 Zircaloy-4				
Cr	906	µg/g	±	9	µg/g
Fe	2143	mg/g	±	0.020	mg/g
Hf	77.6	µg/g	±	3.0	µg/g
Sn	14.60	mg/g	±	0.09	mg/g

Availability: Bottles containing about 10 g of chips.

Substance	BCR-074 Electrolytic copper (OFHC) (mg/kg)	ERM-EB074 Electrolytic copper (mg/kg)	ERM-EB075 Electrolytic copper with added impurities (mg/kg)
Ag	12.8 ± 0.7	1.03 ± 0.07	10.8 ± 0.6
Al	0.78 ± 0.14	1.23 ± 0.08	2.3 ± 0.4
As		0.52 ± 0.06	3.18 ± 0.10
Au		0.31 ± 0.06	1.46 ± 0.14
Be		0.51 ± 0.04	1.08 ± 0.24
Bi	(0.10 ± 0.03)	0.40 ± 0.04	1.79 ± 0.11
Cd	< 0.02	0.83 ± 0.06	2.69 ± 0.09
Co	< 0.05	0.37 ± 0.04	2.64 ± 0.08
Cr	< 0.1	5.8 ± 0.8	1.40 ± 0.07
Fe	1.14 ± 0.06	(< 0.1)	9.3 ± 0.4
Hg		0.49 ± 0.07	(< 0.35)
In		2.03 ± 0.27	1.83 ± 0.10
Mg		0.93 ± 0.07	7.0 ± 0.7
Mn	1.27 ± 0.05	0.61 ± 0.08	1.35 ± 0.07
Ni	1.04 ± 0.11	1.53 ± 0.25	2.18 ± 0.16
P	0.97 ± 0.05	2.7 ± 0.4	2.59 ± 0.30
Pb		(3.3 ± 1.0)	4.8 ± 0.9
S	0.576 ± 0.030	0.57 ± 0.04	25 ± 4
Sb	0.37 ± 0.04	0.55 ± 0.07	2.93 ± 0.14
Se		(1.5 ± 0.4)	1.69 ± 0.10
Si		0.50 ± 0.06	2.6 ± 0.4
Sn	< 0.07	0.97 ± 0.18	2.13 ± 0.11
Te	(0.21 ± 0.08)	(< 0.25)	1.78 ± 0.12
Ti		2.2 ± 0.4	3.2 ± 0.5
W		(8.8 ± 1.7)	(< 0.1)
Zn	0.46 ± 0.07	6.51 ± 0.29	(20 ± 5)
Zr			

Values in brackets are not certified.

Availability: CRMs are available as follows:

BCR-074A: Cylinder of 40 mm Ø, 30 mm height

ERM-EB074A and ERM-EB075A: Disc of 39 mm Ø, 30 mm height

ERM-EB074B and ERM-EB075B: Cylinder of 8 mm Ø, 100 mm length

ERM-EB074C and ERM-EB075C: 50 g Chips of approximately 250 mg, in amber glass bottle

	Substance	Quaternary bronze (g/kg)	Brass (g/kg)	Arsenic-Copper (g/kg)	Lead-bronze (g/kg)	Tin-bronze (g/kg)
BCR-691	As	1.94 ± 0.10	0.99 ± 0.10	46.0 ± 2.7	2.85 ± 0.22	1.94 ± 0.20
	Pb	79 ± 7	3.9 ± 0.4	1.75 ± 0.14	92 ± 17	2.04 ± 0.18
	Sn	71.6 ± 2.1	20.6 ± 0.7	2.02 ± 0.29	101 ± 8	70 ± 6
	Zn	60.2 ± 2.2	148 ± 5	0.55 ± 0.05	1.48 ± 0.24	1.57 ± 0.25

Availability: Set of five discs (one of each composition) of 35 mm Ø and 2 mm thickness, packed in a box.

	Description	Certified S content (g/kg)
BCR-331	Steam Coal	4.99 ± 0.10
BCR-332	High Volatile Industrial Coal	9.61 ± 0.17
BCR-336	High Volatile Steam Coal	32.90 ± 0.26

Availability: These CRMs are available in units of about 20 g in ampoules with argon atmosphere.

Substance	BCR-460 Total Fluorine in coal powder (mg/kg)		
Cl F	(59 225)	± ±	18) 6

Values in brackets are not certified.

Availability: BCR-460 in glass bottles containing about 40 g.

Substance	BCR-461 Fluorine in clay (mg/kg)		
F	568	±	60

Availability: The samples are provided in units of 30 g in glass bottles.

	Description	S content (g/kg)		
ERM-EF672	Gasoil	0.203	±	0.006
ERM-EF671	Gasoil	0.452	±	0.009
ERM-EF104	Gasoil	1.019	±	0.019
BCR-105	Gasoil	3.63	±	0.10
BCR-106	Gasoil	5.02	±	0.08
BCR-107	Gasoil	10.40	±	0.15
ERM-EF211	Petrol	0.048 8	±	0.001 7

Availability: The materials are available in dark glass ampoules sealed under nitrogen. ERM-EF104, -671 and -672 contain 8 mL, BCR-105, -106 and -107 contain 25 g. ERM-EF211 is available in clear borosilicate glass ampoules and contains 19 mL.

	Description	Solvent Yellow 124 (SY124) content (mg/kg)		
ERM-EF317	Gasoil	0.141	±	0.018

Availability: ERM-EF317 is available in dark glass ampoules sealed under nitrogen, containing 20 mL.

Certified Parameter	IRMM-441 n-Heptane (%)	IRMM-442 Isooctane (%)
n-Heptane, purity by difference	99.985 ± 0.005	99.985 ± 0.005
Isooctane, purity by difference		
<u>Impurities</u>		
Total organics (other than isoctane)		0.011 ± 0.004
Total organics (other than n-Heptane)	0.012 ± 0.005	
Isooctane	0.007 ± 0.002	
n-Heptane		0.002 ± 0.002
Water	0.003 ± 0.002	0.004 ± 0.002
Lead	< 0.5 µg/L	< 1 µg/L

Availability: IRMM-441 and -442 are supplied in ampoules of 100 mL.

	Description	Substance	Certified values (g/kg)		
BCR-109	Zinc ore (blende)	Pb	7.38	±	0.03
		Fe	145.1	±	0.6
		Cu	9.46	±	0.08
		Cd	4.61	±	0.09
		Mg	0.20	±	0.01
		F	0.081	±	0.004
		Hg	0.00 96	±	0.000 12
BCR-110	Zinc ore (blende)	Pb	97.8	±	0.4
		Fe	5.46	±	0.10
		Cu	16.28	±	0.12
		Cd	10.51	±	0.07
		Mg	1.36	±	0.04
		F	0.055	±	0.003
		Hg	0.148 4	±	0.002 5

Availability: These RMs are contained in brown glass bottles. The approximate quantity per unit is 200 g for BCR-109 and 75 g for BCR-110.

	Description	Substance	Certified values (g/kg)		
BCR-032	Moroccan Phosphate rock	CaO	517.6	±	3.2
		P ₂ O ₅	329.8	±	1.7
		CO ₂	51.0	±	0.8
		F	40.4	±	0.6
		SiO ₂	20.9	±	1.2
		SO ₃	18.4	±	0.8
		Al ₂ O ₃	5.5	±	0.6
		MgO	4.0	±	0.1
		Fe ₂ O ₃	2.3	±	0.1
		As	9.5 × 10 ⁻³	±	0.5 × 10 ⁻³
		B	22.6 × 10 ⁻³	±	2.2 × 10 ⁻³
		Cd	20.8 × 10 ⁻³	±	0.7 × 10 ⁻³
		Cr	257 × 10 ⁻³	±	16 × 10 ⁻³
		Co	0.59 × 10 ⁻³	±	0.06 × 10 ⁻³
		Cu	33.7 × 10 ⁻³	±	1.4 × 10 ⁻³
		Hg	55 × 10 ⁻⁶	±	11 × 10 ⁻⁶
		Mn	18.8 × 10 ⁻³	±	1.3 × 10 ⁻³
		Ni	34.6 × 10 ⁻³	±	1.9 × 10 ⁻³
		Ti	171 × 10 ⁻³	±	10 × 10 ⁻³
		V	153 × 10 ⁻³	±	7 × 10 ⁻³
		Zn	253 × 10 ⁻³	±	6 × 10 ⁻³

Availability: Units of about 100 g in the form of fine powder.

Substance	BCR-664 Glass (mg/kg)		
As	5.9	±	0.4
Ba	29.1	±	0.7
Cd	5.7	±	0.4
Cl	68	±	8
Co	2.77	±	0.21
Cr	2.65	±	0.13
Pb	53.1	±	2.6
Sb	24.3	±	1.0
Se	8.6	±	0.5

Availability: Glass plate of (50 x 50 x 7) mm.

	Material	Uranium mass fraction (mg/kg)		
IRMM-540R	Uranium-doped oxide glass	15.0	±	0.9
IRMM-541	Uranium-doped oxide glass	49.4	±	2.7

Availability: Glass disc of 15 mm diameter and 2 mm thickness, polished on both sides.

Substance	ERM-EC590 Polyethylene (LDPE) g/kg			ERM-EC591 Polypropylene (PP) g/kg			
	Br	2,4,4'-TriBDE (BDE-28)	0.213	±	0.09	2.08	±
2,2',4,4'-TetraBDE (BDE-47)	0.23	±	0.04	0.0025	±	0.0004	
2,2',4,4',5-PentaBDE (BDE-99)	0.302	±	0.030	0.245	±	0.023	
2,2',4,4',6-PentaBDE (BDE-100)	0.063	±	0.005	0.32	±	0.04	
2,2',4,4',5,5-HexaBDE (BDE-153)	0.047	±	0.006	0.066	±	0.007	
2,2',4,4',5,6-HexaBDE (BDE-154)	0.0257	±	0.0026	0.044	±	0.006	
2,2',3,4,4',5,6-HeptaBDE (BDE-183)	0.132	±	0.012	0.026	±	0.004	
2,2',3,3',4,4',6,6-OctaBDE + 2,2',3,4,4',5,6,6-OctaBDE (BDE-197+204)	0.076	±	0.010	0.087	±	0.008	
DecaBDE (BDE-209)	0.65	±	0.10	0.052	±	0.009	
DecaBB (BB-209)	0.63	±	0.10	0.78	±	0.09	
Sb	(0.756	±	0.025)	0.74	±	0.08	
				(0.713	±	0.022)	

Values in brackets are not certified.

Availability: Brown glass bottle with 20 g granulate.

Substance	ERM-EC680m Polyethylene (low level) mg/kg			ERM-EC681m Polyethylene (high level)			
As	4.7	±	0.4	17.0	±	1.2	mg/kg
Br	181	±	9	1.43	±	0.08	g/kg
Cd	20.8	±	0.9	146	±	5	mg/kg
Cl	(84	±	11)	0.38	±	0.06	g/kg
Cr	9.6	±	0.5	45.1	±	1.9	mg/kg
Hg	2.56	±	0.16	9.9	±	0.8	mg/kg
Pb	11.3	±	0.4	69.7	±	2.5	mg/kg
S	86	±	9	0.64	±	0.10	g/kg
Sb	9.6	±	0.7	86	±	7	mg/kg
Sn	20.7	±	1.6	99	±	6	mg/kg
Zn	194	±	12	1.17	±	0.04	g/kg

Values in brackets are not certified

Availability: Brown glass bottle with 100 g granulate.

Cd in polyethylene (VDA 001-004)

A set of four certified reference materials for Cd in polyethylene (40.9 mg/kg, 75.9 mg/kg, 197.9 mg/kg and 407 mg/kg) has been certified by the JRC on behalf of VDA (Verband der Automobilindustrie e.V., Frankfurt). Information can be obtained from JRC, Geel (B).

5.3 OTHERS

Antimony implanted in silicon ERM-EG001 (BAM-L001)

Certified quantity	Certified value ($10^{16} \cdot \text{cm}^{-2}$)
Areal density of Sb atoms	4.81 ± 0.06
Isotope amount ratio $n(^{121}\text{Sb}) / n(^{123}\text{Sb})$	1.435 ± 0.006

Availability: The sample is a 10 mm x 10 mm silicon chip with a thermally grown surface oxide layer and Sb ions implanted with an energy of 400 keV. The certified value for the areal density of Sb atoms is valid for fractions of the chip surface down to 0.15 mm^2 in size.

Cementite Grains in Carburised Pure Iron (IRMM-471)

	Certified value (g/kg)
Carbon mass fraction in cementite grains	66.9 ± 2.7

Availability: 4-5 mm long rod with 5 mm diameter.

6 MATERIALS RELATED TO ISOTOPIC MEASUREMENTS

6.1 CERTIFIED FOR ISOTOPE ABUNDANCE RATIO (AMOUNT RATIO)

BCR-123 Ethanol				
Parameter	Ethanol H		Ethanol M	Ethanol L
(D/H) _I	109.65×10^{-6}	$\pm 0.20 \times 10^{-6}$	101.69×10^{-6}	$\pm 0.17 \times 10^{-6}$
(D/H) _{II}	119.76×10^{-6}	$\pm 0.25 \times 10^{-6}$	130.94×10^{-6}	$\pm 0.21 \times 10^{-6}$
R	2.184	± 0.005	2.575	± 0.006

Availability: Units of 3 sealed NMR tubes containing respectively H-, M-, and L-ethanols, to which the tetramethylurea internal standard and the C₆F₆ lock substance are added. 10 mm (BCR-123A) or 15 mm (BCR-123B) O.D. NRM tubes can be supplied.

Parameter	Unit	BCR-656 (96% ethanol)	BCR-657 (Sugar)	BCR-660 (Ethanol in water)
(D/H) _I by ² H-NMR	ppm	102.84 \pm 0.20		102.90 \pm 0.16
(D/H) _{II} by ² H-NMR	ppm	132.07 \pm 0.30		131.95 \pm 0.23
R by ² H-NMR		2.570 \pm 0.005		2.567 \pm 0.005
$\delta^{13}\text{C}_{\text{VPDB}}$ by IRMS	‰	-26.91 \pm 0.07	-10.76 \pm 0.04	-26.72 \pm 0.09
(D/H) _w of water (IRMS)	ppm			148.68 \pm 0.14
Alcoholic grade t _D	w/w %	(94)		11.96 \pm 0.06 ¹⁾

1) in v/v %

Value in brackets is not certified.

Availability: BCR-656: Units of 25 mL of 96 % vol. neutral ethanol from wine in glass bottle;

BCR-657: Units of 80pprox.. 1 g of dry glucose in a sealed amber vial;

BCR-660: Units of 450 mL of aqueous ethanol solution in glass bottle.

Code	Description	Isotope amount content	Amount ratios					Unit size
IRMM-	0.5 M HNO ₃	$n(^{68}\text{Zn})/n(^{67}\text{Zn})$ $(^{\text{64}}\text{Zn})\cdot\text{g}^{-1}$	$n(^{66}\text{Zn})/n(^{64}\text{Zn})$	$n(^{67}\text{Zn})/n(^{64}\text{Zn})$	$n(^{68}\text{Zn})/n(^{64}\text{Zn})$	$n(^{70}\text{Zn})/n(^{64}\text{Zn})$	5 mL	
			1.070 00 (47)	0.004 679 7(58)	0.021 337 4 (98)	0.022 830 9 (89)	0.000 067 57 (32)	
007/1		148.261 (49) $\cdot 10^{-9}$ mol	1.033 83 (45)	0.005 275 8 (57)	0.043 039 (20)	0.044 495 (18)	0.000 083 00 (31)	
007/2		142.842 (47) $\cdot 10^{-9}$ mol	1.012 45 (45)	0.007 057 4 (57)	0.107 896 (50)	0.109 239 (45)	0.000 129 11 (32)	
007/3		125.44 (25) $\cdot 10^{-9}$ mol	1.005 44 (45)	0.009 954 0 (60)	0.213 339 (99)	0.214 499 (88)	0.000 204 07 (44)	
007/4		107.096 (40) $\cdot 10^{-9}$ mol	1.001 62 (45)	0.016 608 8 (80)	0.455 60 (21)	0.456 34 (19)	0.000 376 29 (90)	
007/5		79.518 (32) $\cdot 10^{-9}$ mol	0.999 61 (45)	0.035 323 (17)	1.136 83 (52)	1.136 39 (46)	0.000 860 6 (23)	
007/6		45.821 (19) $\cdot 10^{-9}$ mol						

Code	Description	Amount ratios	Unit size
		$n(^{41}\text{Ca})/n(^{40}\text{Ca})$	
ERM-AE701/1 ERM-AE701/2 ERM-AE701/3 ERM-AE701/4 ERM-AE701/5 ERM-AE701/6 ERM-AE701/7 ERM-AE701/8	0.6 M HNO_3 solution	1.011 4(68) · 10^{-6} 1.023 5(69) · 10^{-7} 1.018 1(69) · 10^{-8} 1.047 9(71) · 10^{-9} 1.052 0(71) · 10^{-10} 1.091 3(74) · 10^{-11} 1.054 9(72) · 10^{-12} 1.052 4(71) · 10^{-13}	25 mL

Code	Description	Isotope amount fraction (-100)						Amount ratios			Unit size		
IRMM-009	0.2 M HNO_3 solution	^{24}Mg	^{25}Mg	^{26}Mg				$n(^{25}\text{Mg})/n(^{24}\text{Mg})$	$n(^{26}\text{Mg})/n(^{24}\text{Mg})$		4 mL		
		78.992(25)	10.003(9)	11.005(19)				0.126 63(13)	0.139 32(26)				
IRMM-010	Pt metal	^{190}Pt	^{192}Pt	^{194}Pt	^{195}Pt	^{196}Pt	^{198}Pt	$n(^{190}\text{Pt})/n(^{195}\text{Pt})$	$n(^{192}\text{Pt})/n(^{195}\text{Pt})$	$n(^{194}\text{Pt})/n(^{195}\text{Pt})$	30 mg (wire)		
		0.011 7(11)	0.782(17)	32.86(27)	33.78(16)	25.21(23)	7.356(82)	0.000 347(34)	0.023 15(48)	0.973(11)			
IRMM-011	H_3BO_3 solid	^{10}B	^{11}B					$n(^{10}\text{B})/n(^{11}\text{B})$			1 g		
		19.824 (20)	80.176(20)					0.247 26(32)					
IRMM-012	1 M HCl solution	^{50}Cr	^{52}Cr	^{53}Cr	^{54}Cr			$n(^{50}\text{Cr})/n(^{52}\text{Cr})$	$n(^{53}\text{Cr})/n(^{52}\text{Cr})$	$n(^{54}\text{Cr})/n(^{52}\text{Cr})$	5 mL		
		4.345(9)	83.789(2)	9.501(11)	2.365(5)			0.051 86(10)	0.113 39(15)	0.028 22(06)			
IRMM-016	Li_2CO_3 solid	^6Li	^7Li					$n(^6\text{Li})/n(^7\text{Li})$			1 g		
		7.588 9(75)	92.411 1(75)					0.082 121(87)					
IRMM-018°	SiO_2 solid	^{28}Si	^{29}Si	^{30}Si				$n(^{29}\text{Si})/n(^{28}\text{Si})$	$n(^{30}\text{Si})/n(^{28}\text{Si})$		5 g		
		92.220 36(49)	4.687 30(36)	3.092 34(37)				0.050 827 2(40)	0.033 532 0(42)				

6.2 CERTIFIED FOR ISOTOPE AMOUNT CONTENT

Code	Description	Isotope amount content	Isotope enrichment	Unit size
IRMM-610	H ₃ BO ₃ aqueous solution	3.683 11 (88) μmol ¹⁰ B·g ⁻¹	$n(^{10}\text{B})/n(^{11}\text{B}) = 18.80$ (2)	5 mL
IRMM-611	H ₃ BO ₃ aqueous solution	4.025 (40) μmol ¹¹ B·g ⁻¹	$n(^{10}\text{B})/n(^{11}\text{B}) = 0.247$ 26 (32)	5 mL
IRMM-615	0.5 M HCl solution	3.850 (14) μmol ⁶ Li·g ⁻¹	$n(^6\text{Li})/n(^7\text{Li}) = 21.897$ (44)	5 mL
IRMM-618	0.5 M HNO ₃ solution	112.13 (17) μmol ⁸⁷ Rb·kg ⁻¹	$n(^{85}\text{Rb})/n(^{87}\text{Rb}) = 0.20$ 498 (24)	5 mL
IRMM-619	0.5 M HNO ₃ solution	85.00 (24) μmol ⁸⁵ Rb·kg ⁻¹	$n(^{85}\text{Rb})/n(^{87}\text{Rb}) = 2.593$ 0 (20)	5 mL
IRMM-620	4.5 M HCl solution	173.35 (16) μmol ⁵⁷ Fe·kg ⁻¹	$n(^{54}\text{Fe})/n(^{57}\text{Fe}) = < 0.0001$ $n(^{56}\text{Fe})/n(^{57}\text{Fe}) = 0.025$ 39 (31) $n(^{58}\text{Fe})/n(^{57}\text{Fe}) = 0.025$ 16 (18)	5 mL
IRMM-621	1 M HNO ₃ solution	97.35 (15) μmol ¹¹¹ Cd·kg ⁻¹	$n(^{106}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{108}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{110}\text{Cd})/n(^{111}\text{Cd}) = 0.004$ 44 (42) $n(^{112}\text{Cd})/n(^{111}\text{Cd}) = 0.021$ 74 (10) $n(^{113}\text{Cd})/n(^{111}\text{Cd}) = 0.005$ 818 (56) $n(^{114}\text{Cd})/n(^{111}\text{Cd}) = 0.010$ 875 (88) $n(^{116}\text{Cd})/n(^{111}\text{Cd}) = 0.001$ 629 (44)	4 mL
IRMM-622	1 M HNO ₃ solution	9.739 (18) μmol ¹¹¹ Cd·kg ⁻¹	$n(^{106}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{108}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{110}\text{Cd})/n(^{111}\text{Cd}) = 0.004$ 44 (42) $n(^{112}\text{Cd})/n(^{111}\text{Cd}) = 0.021$ 74 (10) $n(^{113}\text{Cd})/n(^{111}\text{Cd}) = 0.005$ 818 (56) $n(^{114}\text{Cd})/n(^{111}\text{Cd}) = 0.010$ 875 (88) $n(^{116}\text{Cd})/n(^{111}\text{Cd}) = 0.001$ 629 (44)	4 mL
IRMM-624	1 M HCl solution	174.84 (42) μmol ⁵⁰ Cr·kg ⁻¹	$n(^{52}\text{Cr})/n(^{50}\text{Cr}) = 0.066$ 41 (50) $n(^{53}\text{Cr})/n(^{50}\text{Cr}) = 0.000$ 323 (64) $n(^{54}\text{Cr})/n(^{50}\text{Cr}) = 0.000$ 11 (11)	5 mL
IRMM-625	1 M HCl solution	144.233 (90) μmol ⁵² Cr·kg ⁻¹	$n(^{50}\text{Cr})/n(^{52}\text{Cr}) = 0.051$ 85 (20) $n(^{53}\text{Cr})/n(^{52}\text{Cr}) = 0.113$ 33 (38) $n(^{54}\text{Cr})/n(^{52}\text{Cr}) = 0.028$ 35 (34)	5 mL
IRMM-632	1 M HNO ₃ solution	0.096 84 (41) μmol ⁶⁵ Cu·g ⁻¹	$n(^{63}\text{Cu})/n(^{65}\text{Cu}) = 0.002$ 892 1 (92)	5 mL
ERM-AE633	1 M HNO ₃ solution	5.998 (36) μmol ⁶³ Cu·g ⁻¹	$n(^{65}\text{Cu})/n(^{63}\text{Cu}) = 0.445$ 63 (42)	4 – 5 mL
IRMM-634	1.8 M HCl solution	163.61 (38) μmol ⁵⁶ Fe·kg ⁻¹	$n(^{54}\text{Fe})/n(^{56}\text{Fe}) = 0.063$ 70 (27) $n(^{57}\text{Fe})/n(^{56}\text{Fe}) = 0.023$ 096 (72) $n(^{58}\text{Fe})/n(^{56}\text{Fe}) = 0.003$ 071 (29)	5 mL
ERM-AE637	0.2 M HNO ₃ solution	0.791 37(30) μmol ²⁴ Mg·g ⁻¹	$n(^{26}\text{Mg})/n(^{24}\text{Mg}) = 0.139$ 68 (32) $n(^{25}\text{Mg})/n(^{24}\text{Mg}) = 0.126$ 86 (18)	5 mL
ERM-AE638	0.1 M HNO ₃ solution	0.857 4 (34) μmol ²⁶ Mg·g ⁻¹	$n(^{24}\text{Mg})/n(^{26}\text{Mg}) = 0.003$ 104 (26) $n(^{25}\text{Mg})/n(^{26}\text{Mg}) = 0.001$ 084 (11)	5 mL
ERM-AE639	0.5 M HCl solution	11.891 (50) μmol ²⁰² Hg·g ⁻¹	$n(^{196}\text{Hg})/n(^{202}\text{Hg}) = 0.004$ 972 (46) $n(^{198}\text{Hg})/n(^{202}\text{Hg}) = 0.330$ 6 (21) $n(^{199}\text{Hg})/n(^{202}\text{Hg}) = 0.561$ 9 (28) $n(^{200}\text{Hg})/n(^{202}\text{Hg}) = 0.770$ 5 (28) $n(^{201}\text{Hg})/n(^{202}\text{Hg}) = 0.441$ 26 (88) $n(^{204}\text{Hg})/n(^{202}\text{Hg}) = 0.230$ 27 (75)	5 mL

ERM-AE640	0.5 M HCl solution	14.71 (11) nmol $^{202}\text{Hg}\cdot\text{g}^{-1}$	$n(^{196}\text{Hg})/n(^{202}\text{Hg}) = 0.000\ 018\ 09\ (38)$ $n(^{198}\text{Hg})/n(^{202}\text{Hg}) = 0.000\ 623\ (11)$ $n(^{199}\text{Hg})/n(^{202}\text{Hg}) = 0.001\ 603\ (16)$ $n(^{200}\text{Hg})/n(^{202}\text{Hg}) = 0.005\ 499\ (34)$ $n(^{201}\text{Hg})/n(^{202}\text{Hg}) = 0.013\ 351\ (52)$ $n(^{204}\text{Hg})/n(^{202}\text{Hg}) = 0.002\ 595\ (21)$	5 mL
ERM-AE641	Cl in water	18.959 (15) $\mu\text{mol } ^{35}\text{Cl}\cdot\text{g}^{-1}$	$n(^{37}\text{Cl})/n(^{35}\text{Cl}) = 0.319\ 77\ (83)$	4 – 5 mL
ERM-AE642	Cl in water	4.375 (26) $\mu\text{mol } ^{37}\text{Cl}\cdot\text{g}^{-1}$	$n(^{35}\text{Cl})/n(^{37}\text{Cl}) = 0.019\ 14\ (48)$	4 – 5 mL
IRMM-643	2.8 M HNO ₃ solution	334.33 (84) $\mu\text{mol } ^{32}\text{S}\cdot\text{kg}^{-1}$	$n(^{33}\text{S})/n(^{32}\text{S}) = 0.007\ 877\ 6\ (58)$ $n(^{34}\text{S})/n(^{32}\text{S}) = 0.044\ 149\ 3\ (78)$ $n(^{36}\text{S})/n(^{32}\text{S}) = 0.000\ 153\ 40\ (94)$	5 mL
IRMM-644	3.2 M HNO ₃ solution	326.28 (80) $\mu\text{mol } ^{32}\text{S}\cdot\text{kg}^{-1}$	$n(^{33}\text{S})/n(^{32}\text{S}) = 0.007\ 969\ 8\ (70)$ $n(^{34}\text{S})/n(^{32}\text{S}) = 0.045\ 162\ 2\ (82)$ $n(^{36}\text{S})/n(^{32}\text{S}) = 0.000\ 170\ 0\ (58)$	5 mL
IRMM-645	2.8 M HNO ₃ solution	371.96 (57) $\mu\text{mol } ^{32}\text{S}\cdot\text{kg}^{-1}$	$n(^{33}\text{S})/n(^{32}\text{S}) = 0.007\ 747\ 6\ (38)$ $n(^{34}\text{S})/n(^{32}\text{S}) = 0.042\ 747\ 3\ (62)$ $n(^{36}\text{S})/n(^{32}\text{S}) = 0.000\ 145\ 1\ (42)$	5 mL
IRMM-646	2.8 M HNO ₃ solution	4586 (27) $\mu\text{mol } ^{34}\text{S}\cdot\text{kg}^{-1}$	$n(^{32}\text{S})/n(^{34}\text{S}) = 0.038\ 314\ 9\ (31)$ $n(^{33}\text{S})/n(^{34}\text{S}) = 0.000\ 470\ 88\ (15)$ $n(^{36}\text{S})/n(^{34}\text{S}) = 0.000\ 018\ 1\ (37)$	5 mL
ERM-AE647	1 M HNO ₃ solution	134.974 (73) $\mu\text{mol } ^{63}\text{Cu}\cdot\text{g}^{-1}$	$n(^{65}\text{Cu})/n(^{63}\text{Cu}) = 0.445\ 60\ (74)$	4 mL
ERM-AE649	1 M HNO ₃ solution	0.836 88 (27) $\mu\text{mol } ^{205}\text{Tl}\cdot\text{g}^{-1}$	$n(^{203}\text{Tl})/n(^{205}\text{Tl}) = 0.418\ 91\ (18)$	4 – 5 mL
IRMM-651	0.5 M HNO ₃ solution	0.077 506 (30) $\mu\text{mol } ^{64}\text{Zn}\cdot\text{g}^{-1}$	$n(^{66}\text{Zn})/n(^{64}\text{Zn}) = 0.557\ 17\ (30)$ $n(^{67}\text{Zn})/n(^{64}\text{Zn}) = 0.080\ 702\ (34)$ $n(^{68}\text{Zn})/n(^{64}\text{Zn}) = 0.366\ 27\ (12)$ $n(^{70}\text{Zn})/n(^{64}\text{Zn}) = 0.011\ 981\ (22)$	5 mL
IRMM-652	0.5 M HNO ₃ solution	0.156 000 (50) $\mu\text{mol } ^{64}\text{Zn}\cdot\text{g}^{-1}$	$n(^{66}\text{Zn})/n(^{64}\text{Zn}) = 0.004\ 107\ 3\ (59)$ $n(^{67}\text{Zn})/n(^{64}\text{Zn}) = 0.000\ 499\ 87\ (96)$ $n(^{68}\text{Zn})/n(^{64}\text{Zn}) = 0.002\ 029\ 5\ (23)$ $n(^{70}\text{Zn})/n(^{64}\text{Zn}) = 0.000\ 052\ 76\ (34)$	5 mL
IRMM-653	0.5 M HNO ₃ solution	0.138 014 (60) $\mu\text{mol } ^{67}\text{Zn}\cdot\text{g}^{-1}$	$n(^{64}\text{Zn})/n(^{67}\text{Zn}) = 0.013\ 191\ 5\ (81)$ $n(^{66}\text{Zn})/n(^{67}\text{Zn}) = 0.024\ 551\ 6\ (70)$ $n(^{68}\text{Zn})/n(^{67}\text{Zn}) = 0.051\ 086\ (36)$ $n(^{70}\text{Zn})/n(^{67}\text{Zn}) = 0.000\ 527\ 8\ (18)$	5 mL
IRMM-654	0.5 M HNO ₃ solution	0.146 098 (48) $\mu\text{mol } ^{68}\text{Zn}\cdot\text{g}^{-1}$	$n(^{64}\text{Zn})/n(^{68}\text{Zn}) = 0.00\ 489\ 4\ (38)$ $n(^{66}\text{Zn})/n(^{68}\text{Zn}) = 0.003\ 186\ 8\ (46)$ $n(^{68}\text{Zn})/n(^{68}\text{Zn}) = 0.001\ 411\ 3\ (17)$ $n(^{70}\text{Zn})/n(^{68}\text{Zn}) = 0.000\ 194\ 98\ (78)$	5 mL
ERM-AE671	$\text{CH}_3^{202}\text{HgCl}$ in 2 % ethanol/water solution	0.015 1 (7) $\mu\text{mol } \text{CH}_3^{202}\text{HgCl}\cdot\text{g}^{-1}$	$n(^{196}\text{Hg})/n(^{202}\text{Hg}) = 0.000\ 018$ $n(^{198}\text{Hg})/n(^{202}\text{Hg}) = 0.000\ 62$ $n(^{199}\text{Hg})/n(^{202}\text{Hg}) = 0.001\ 60$ $n(^{200}\text{Hg})/n(^{202}\text{Hg}) = 0.005\ 50$ $n(^{201}\text{Hg})/n(^{202}\text{Hg}) = 0.013\ 4$ $n(^{204}\text{Hg})/n(^{202}\text{Hg}) = 0.002\ 60$	5 g
IRMM-3702	1 M HNO ₃ solution	1.512 (30) $\mu\text{mol } ^{64}\text{Zn}\cdot\text{g}^{-1}$	$n(^{66}\text{Zn})/n(^{64}\text{Zn}) = 0.563\ 97\ (30)$ $n(^{67}\text{Zn})/n(^{64}\text{Zn}) = 0.082\ 166\ (35)$ $n(^{68}\text{Zn})/n(^{64}\text{Zn}) = 0.375\ 19\ (16)$ $n(^{70}\text{Zn})/n(^{64}\text{Zn}) = 0.012\ 418\ (23)$	3 mL

INDEX

NUMERICAL LIST

ERM / CRM	DESIGNATION	PAGE NO.
ERM-AC057	AFLATOXIN B1 IN ACETONITRILE	30
ERM-AC058	AFLATOXIN B2 IN ACETONITRILE	30
ERM-AC059	AFLATOXIN G1 IN ACETONITRILE	30
ERM-AC060	AFLATOXIN G2 IN ACETONITRILE	27
ERM-AC213	PAHs IN ACETONITRILE / TOLUENE	12
ERM-AC626	ARSENObETAINe IN WATER	30
ERM-AC699	ZEARALENONE IN ACETONITRILE	30
ERM-AD149	THROMBOPLASTIN RABBIT (prothrombin time)	64
ERM-AD413	PLASMID DNA FRAGMENTS OF MON 810 MAIZE	32
ERM-AD415	PLASMID DNA FRAGMENTS OF NK603 MAIZE	33
ERM-AD425	PLASMID DNA FRAGMENTS OF 356043 SOYBEAN	36
ERM-AD427	PLASMID DNA FRAGMENTS OF 98140 MAIZE	37
ERM-AD442k	LAMBDA DNA	63
ERM-AD452/IFCC	γ -GLUTAMYLTRANSFERASE (catalytic concentration)	62
ERM-AD453k/IFCC	LACTATE DEHYDROGENASE ISOENZYME 1 (LD1)	62
ERM-AD454k/IFCC	ALANINE AMINOTRANSFERASE (ALT)	62
ERM-AD455k/IFCC	CREATINE KINASE ISOENZYME MM (CK-MM)	62
ERM-AD457/IFCC	ASPARTATE TRANSAMINASE (AST)	62
ERM-AD482	RUMINANT pDNA CALIBRANT	56
ERM-AD483	PORCINE pDNA CALIBRANT	56
ERM-AD500/IFCC	HAEMOGLOBIN IN BUFFER	62
ERM-AD623	BCR-ABL pDNA CALIBRANT	63
ERM-AE633	COPPER (natural) spike, nitrate solution	82
ERM-AE637	MAGNESIUM (natural) spike, nitrate solution	82
ERM-AE638	MAGNESIUM-26 spike, nitrate solution	82
ERM-AE639	MERCURY (natural) spike, chloride solution	82
ERM-AE640	MERCURY-202 spike, chloride solution	83
ERM-AE641	CHLORIDE (natural) spike, chloride solution	83
ERM-AE642	CHLORIDE-37 spike, chloride solution	83
ERM-AE647	COPPER-63, nitrate solution	83
ERM-AE649	THALLIUM (natural) spike, nitrate solution	83
ERM-AE671	METHYLMERCURY IN 2 % ETHANOL/WATER SOLUTION	83
ERM-AE701	CALCIUM-41 isotopic, nitrate solution (set of 8 units)	81
ERM-BB124	PORK MUSCLE	55
ERM-BB130	PORK MUSCLE	54
ERM-BB184	BOVINE MUSCLE (trace elements)	45
ERM-BB186	PIG KIDNEY (trace elements)	45
ERM-BB350	FISH OIL	41
ERM-BB384	PORK MUSCLE	46, 50
ERM-BB386	BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol) (blank)	54
ERM-BB389	BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol)	54
ERM-BB422	FISH MUSCLE (trace elements)	45
ERM-BB430	PORK FAT (pesticides)	42
ERM-BB444	NATURAL PORK FAT (blank)	41
ERM-BB445	SPIKED PORK FAT (very low level)	41
ERM-BB446	SPIKED PORK FAT (low level)	41

ERM-BB492	MILK POWDER (oxytetracycline)	55
ERM-BB493	MILK POWDER (oxytetracycline) (blank)	55
ERM-BC190	RAPESEED (colza) (S, total glucosinolate, medium level)	43
ERM-BC366	RAPESEED (colza) (S, total glucosinolate, low level)	43
ERM-BC367	RAPESEED (colza) (S, total glucosinolate, high level)	43
ERM-BC211	RICE (As species)	24
ERM-BC381	RYE FLOUR	46, 50
ERM-BC382	WHEAT FLOUR	46, 50
ERM-BC514	HARICOTS BEANS (dietary fibre)	51
ERM-BC515	CARROT (dietary fibre)	51
ERM-BC516	APPLE (dietary fibre)	51
ERM-BC517	FULL FAT SOYA (dietary fibre)	51
ERM-BC716	MAIZE	44
ERM-BC717	MAIZE	44
ERM-BD150	SKIMMED MILK POWDER (trace elements)	45
ERM-BD151	SKIMMED MILK POWDER (trace elements)	45
ERM-BD273	TOASTED BREAD	57
ERM-BD282	WHOLE MILK POWDER (aflatoxin M1, zero level)	43
ERM-BD283	WHOLE MILK POWDER (aflatoxin M1, low level)	43
ERM-BD284	WHOLE MILK POWDER (aflatoxin M1, high level)	43
ERM-BD518	BRAN BREAKFAST CEREAL (dietary fibre)	51
ERM-BD600	WHOLE MILK POWDER (vitamins)	49
ERM-BE375	COMPOUND FEEDINGSTUFF (very low level)	43
ERM-BE376	COMPOUND FEEDINGSTUFF (high level)	43
ERM-BF410p	GENETICALLY MODIFIED ROUNDUP READY SOYA	31
ERM-BF411	GENETICALLY MODIFIED Bt-176 MAIZE	31
ERM-BF412k	GENETICALLY MODIFIED Bt-11 MAIZE	32
ERM-BF413k	GENETICALLY MODIFIED MON 810 MAIZE	32
ERM-BF414	GENETICALLY MODIFIED GA21 MAIZE	33
ERM-BF415	GENETICALLY MODIFIED NK603 MAIZE	33
ERM-BF416	GENETICALLY MODIFIED MON 863 MAIZE	33
ERM-BF417	GENETICALLY MODIFIED MON 863 x MON 810 MAIZE	34
ERM-BF418	GENETICALLY MODIFIED 1507 MAIZE	34
ERM-BF419	GENETICALLY MODIFIED H7-1 SUGAR BEET	34
ERM-BF420	GENETICALLY MODIFIED 3272 MAIZE	34
ERM-BF421	GENETICALLY MODIFIED EH92-527-1 POTATO	35
ERM-BF422	GENETICALLY MODIFIED 281-24-236 X 3006-210-23 COTTON SEED	35
ERM-BF423	GENETICALLY MODIFIED MIR604 MAIZE	35
ERM-BF424	GENETICALLY MODIFIED 59122 MAIZE	35
ERM-BF425	GENETICALLY MODIFIED SOYA 356043	36
ERM-BF426	GENETICALLY MODIFIED SOYA 305423	36
ERM-BF427	GENETICALLY MODIFIED 98140 MAIZE	36
ERM-BF428	GENETICALLY MODIFIED GHB119 COTTON	37
ERM-BF429	GENETICALLY MODIFIED T304-40 COTTON	37
ERM-BF430	GENETICALLE MODIFIED AM04-1020 POTATO	37
ERM-BF431	GENETICALLE MODIFIED AV43-6-G7 POTATO	38
ERM-BF432	GENETICALLY MODIFIED SOYA DAS-68416-4	38
ERM-BF433	GENETICALLY MODIFIED DAS-40278-9 MAIZE	38
ERM-BF434	GENETICALLY MODIFIED 73496 RAPESEED	38
ERM-BF435	GENETICALLY MODIFIED PH05-026-0048 POTATO	39
ERM-BF436	GENETICALLY MODIFIED DAS-44406-6 SOYA	39
ERM-BF437	GENETICALLY MODIFIED DAS-81419-2 SOYA	39
ERM-BF438	GENETICALLY MODIFIED VCO-Ø1981-5 MAIZE	39
ERM-BF439	GENETICALLY MODIFIED DP-ØØ4114-3 MAIZE	40

ERM-BF440	GENETICALLY MODIFIED DAS-81910-7 COTTON	40
ERM-CA100	SURFACE WATER	28
ERM-CA400	SEAWATER (Hg)	21
ERM-CA408	SIMULATED RAINWATER (major components)	20
ERM-CA615	GROUND WATER	21
ERM-CA616	GROUND WATER	20
ERM-CA713	WASTE WATER	21
ERM-CC141	LOAM SOIL	13
ERM-CC144	SEWAGE SLUDGE	13
ERM-CC580	ESTUARINE SEDIMENT (Hg, methylmercury)	14, 23
ERM-CC690	CALCAREOUS SOIL	13
ERM-CD200	BLADDERWRACK (<i>Fucus vesiculosus</i>) (trace elements)	17
ERM-CD281	RYE GRASS	16
ERM-CE100	FISH TISSUE	28
ERM-CE195	BOVINE BLOOD (Pb, Cd)	59
ERM-CE196	BOVINE BLOOD (Pb, Cd)	59
ERM-CE278k	MUSSEL TISSUE (elements)	18
ERM-CE464	TUNA FISH (total and methylmercury)	19, 24
ERM-CE477	MUSSEL TISSUE (butyltins)	24
ERM-CZ100	FINE DUST (PM ₁₀ -like) (PAHs)	24
ERM-CZ120	FINE DUST (PM ₁₀ -like) (elements)	15
ERM-DA192	HUMAN SERUM (cortisol unspiked)	59
ERM-DA193	HUMAN SERUM (cortisol spiked)	59
ERM-DA347	HUMAN SERUM (progesterone)	59
ERM-DA451/IFCC	CORTISOL REFERENCE SERUM PANEL	58
ERM-DA470k/IFCC	HUMAN SERUM (proteins)	61
ERM-DA471/IFCC	HUMAN SERUM (cystatin C)	61
ERM-DA474/IFCC	HUMAN SERUM (CRP)	61
ERM-DA476/IFCC	IgG ANTI-MPO IN HUMAN SERUM	61
ERM-DA480/IFCC	ABETA 42 IN CEREBROSPINAL FLUID (CSF) (low level)	61
ERM-DA481/IFCC	ABETA 42 IN CEREBROSPINAL FLUID (CSF) (medium level)	61
ERM-DA482/IFCC	ABETA 42 IN CEREBROSPINAL FLUID (CSF) (high level)	61
ERM-DA483/IFCC	IgG PR3 ANCA IN HUMAN SERUM	61
ERM-DB001	HUMAN HAIR (trace elements)	60
ERM-EB074A-B-C	ELECTROLYTIC COPPER (trace elements)	76
ERM-EB075A-B-C	ELECTROLYTIC COPPER WITH ADDED IMPURITIES (trace elements)	76
ERM-EB322	UNALLOYED ZINC (trace elements)	74
ERM-EB323	UNALLOYED ZINC (trace elements)	74
ERM-EB324	UNALLOYED ZINC (trace elements)	74
ERM-EB325	UNALLOYED ZINC (trace elements)	74
ERM-EB530A	Al-0.1 % Au (0.1 mm foil)	73
ERM-EB530B	Al-0.1 % Au (0.5 mm wire)	73
ERM-EB530C	Al-0.1 % Au (1.0 mm wire)	73
ERM-EC590	POLYETHYLENE (LDPE)	78
ERM-EC591	POLYPROPYLENE (PP)	78
ERM-EC680m	POLYETHYLENE (low level)	79
ERM-EC681m	POLYETHYLENE (high level)	79
ERM-EF001	BIODIESEL	71
ERM-EF002	BIODIESEL (B100 Rapeseed)	65, 71
ERM-EF003	DIESEL (B7)	71
ERM-EF004	DIESEL (B7)	65, 71
ERM-EF104	GAS OIL (0.1019 % S)	77
ERM-EF211	PETROL	77

ERM-EF317	GAS OIL (Solvent Yellow 124)	77
ERM-EF411	HARD COAL	65
ERM-EF412	BROWN COAL	65
ERM-EF413	FURNACE COKE	65
ERM-EF671	GAS OIL (0.0452 % S)	77
ERM-EF672	GAS OIL (0.0203 % S)	77
ERM-EG001	ANTIMONY IMPLANTED IN SILICON	79
ERM-FA013	CHARPY SPECIMENS Low Energy 20°C (impact toughness)	67
ERM-FA013	CHARPY SPECIMENS Low Energy 0°C (impact toughness)	67
ERM-FA015	CHARPY SPECIMENS 80 J (impact toughness)	67
ERM-FA016	CHARPY SPECIMENS 120 J (impact toughness)	67
ERM-FA415	CHARPY SPECIMENS 150 J (impact toughness)	67
ERM-FC395k	GAS OIL (CFPP and CP)	65
ERM-FD066	CORUNDUM	68
ERM-FD069	CORUNDUM	68
ERM-FD100	COLLOIDAL SILICA (particle size)	69
ERM-FD101b	COLLOIDAL SILICA IN AQUEOUS SOLUTION (nanoparticles)	69
ERM-FD102	COLLOIDAL SILICA IN AQUEOUS SOLUTION (particle size)	69
ERM-FD304	COLLOIDAL SILICA (particle size)	69
BCR-010	TIN ORE CONCENTRATE (Sn)	72
BCR-017A, B	COPPER (S, P)	74
BCR-022A, B	ELECTROLYTIC TOUGH PITCH COPPER (O)	74
BCR-024B, C	TITANIUM (O, N)	74
BCR-032	MOROCCAN PHOSPHATE ROCK (trace elements)	72, 78
BCR-033	SUPERPHOSPHATE (various parameters)	72
BCR-038	FLY ASH FROM PULVERISED COAL (trace elements)	15
BCR-046	BENZO[b]CHRYSENE (purity)	4
BCR-047	BENZO[b]FLUORANTHENE (purity)	4
BCR-048R	BENZO[k]FLUORANTHENE (purity)	4
BCR-049	BENZO[jj]FLUORANTHENE (purity)	4
BCR-050	BENZO[e]PYRENE (purity)	4
BCR-052	BENZO[ghi]PERYLENE (purity)	4
BCR-054R	COPPER (O)	74
BCR-058	CONTINUOUS CAST COPPER (O)	74
BCR-059A, B	Ti 6Al 4V ALLOY (O)	74
BCR-066	QUARTZ (particle size 0.35 – 3.50 microns)	68
BCR-067	QUARTZ (particle size 2.40 – 32.00 microns)	68
BCR-068	QUARTZ (particle size 160 – 630 microns)	68
BCR-069	QUARTZ (particle size 14 – 90 microns)	68
BCR-070	QUARTZ (1.20 – 20.00 microns)	68
BCR-074A	ELECTROLYTIC COPPER (trace elements)	76
BCR-077R	1-METHYLCHRYSENE (purity)	4
BCR-078R	2-METHYLCHRYSENE (purity)	4
BCR-079R	3-METHYLCHRYSENE (purity)	4
BCR-080R	4-METHYLCHRYSENE (purity)	4
BCR-081R	5-METHYLCHRYSENE (purity)	4
BCR-089	TiAl6V4 (Al, V)	75
BCR-090A, B	TITANIUM (impurities)	75
BCR-091	ANTHANTHRENE (purity)	4
BCR-092	10-AZABENZO[a]PYRENE (purity)	4
BCR-093R	1-METHYLBENZ[a]ANTHRACENE (purity)	4
BCR-094	DIBENZ[a,c]ANTHRACENE (purity)	4
BCR-095	DIBENZ[a,j]ANTHRACENE (purity)	4

BCR-096	DIBENZO[a,l]PYRENE (88mylas)	4
BCR-097	BENZO[a]FLUORANTHENE (purity)	4
BCR-098	ZIRCALOY-4 (trace element impurities)	75
BCR-102	TUNGSTEN CARBIDE POWDER (O)	74
BCR-105	GAS OIL (0.363 % S)	77
BCR-106	GAS OIL (0.502 % S)	77
BCR-107	GAS OIL (1.040 % S)	77
BCR-109	ZINC ORE CONCENTRATE (trace elements)	77
BCR-110	ZINC ORE CONCENTRATE (trace elements)	77
BCR-113	POTASSIUM CHLORIDE FERTILIZER (elemental composition)	72
BCR-114	POTASSIUM SULPHATE FERTILIZER (elemental composition)	72
BCR-115	ANIMAL FEED (Organochlorine pesticides)	42
BCR-116	LIMESTONE POWDERS (for shear testing)	66
BCR-121	WHOLEMEAL FLOUR (vitamins)	49
BCR-122	MARGARINE (vitamins)	49
BCR-123A, B	3 REFERENCE ETHANOLS (H, M, L) (for SNIF-NMR)	29, 80
BCR-126A/B	LEAD GLASS (composition/refractive index)	73
BCR-129	HAY POWDER (elements)	16
BCR-130	QUARTZ (particle size 50 – 220 microns)	68
BCR-131	QUARTZ (particle size 480 – 1800 microns)	68
BCR-132	QUARTZ (particle size 1400 – 5000 microns)	68
BCR-133	DIBENZO[a,e]PYRENE (88mylas)	4
BCR-134	BENZO[c]PHENANTHRENE (purity)	4
BCR-136R	BENZO[b]NAPHTHO[2,3-d]THIOPHENE (purity)	4
BCR-137R	BENZO[b]NAPHTHO[1,2-d]THIOPHENE (purity)	4
BCR-138	DIBENZO[a,h]ANTHRACENE (purity)	4
BCR-139	BENZO[ghi]FLUORANTHENE (purity)	4
BCR-140	BENZO[c]CHRYSENE (purity)	4
BCR-142R	LIGHT SANDY SOIL (trace elements)	13
BCR-143R	SEWAGE SLUDGE AMENDED SOIL (trace elements)	13
BCR-145R	SEWAGE SLUDGE (mixed origin) (trace elements)	14
BCR-146R	SEWAGE SLUDGE (industrial origin) (trace elements)	14
BCR-152	DIBENZ[a,l]ACRIDINE (88mylas)	4
BCR-153R	DIBENZ[a,h]ACRIDINE (purity)	4
BCR-154	DIBENZ[a,j]ACRIDINE (purity)	4
BCR-155	DIBENZ[a,c]ACRIDINE (88mylas)	4
BCR-156R	DIBENZ[c,h]ACRIDINE (purity)	4
BCR-157	BENZ[a]ACRIDINE (purity)	4
BCR-158	BENZ[c]ACRIDINE (purity)	4
BCR-159	DIBENZO[a,h]PYRENE (88mylas)	4
BCR-160R	FLUORANTHENE (purity)	4
BCR-162R	SOYA-MAIZE OIL BLEND (fatty acid profile)	47
BCR-163	BEEF-PORK FAT BLEND (fatty acid profile)	47
BCR-165	LATEX SPHERES (particle diameter 2 microns)	58
BCR-166	LATEX SPHERES (particle diameter 4.8 microns)	58
BCR-167	LATEX SPHERES (particle diameter 9.6 microns)	58
BCR-168	PICENE (purity)	4
BCR-169	ALPHA ALUMINA (0.10 m ² /g) (nitrogen BET specific surface area)	68
BCR-170	ALPHA ALUMINA (1.05 m ² /g) (nitrogen BET specific surface area)	68
BCR-171	ALUMINA (2.95 m ² /g) (nitrogen BET specific surface area)	68
BCR-172	QUARTZ (2.50 m ² /g) (nitrogen BET specific surface area)	68
BCR-173	TITANIA (8.23 m ² /g) (nitrogen BET specific surface area)	68
BCR-175	TUNGSTEN (0.18 m ² /g) (nitrogen BET specific surface area)	68
BCR-176R	FLY ASH (trace elements)	15

BCR-177R	PYRENE (purity)	4
BCR-178	CALCIUM AMMONIUM NITRATE FERTILIZER (composition)	72
BCR-179	UREA FERTILIZER (composition)	72
BCR-185R	BOVINE LIVER (trace elements)	45
BCR-187	NATURAL MILK POWDER (pesticides)	42
BCR-188	SPIKED MILK POWDER (pesticides)	42
BCR-191	BROWN BREAD (trace elements)	45
BCR-261T	TANTALUM PENTOXIDE ON TANTALUM FOIL	69
BCR-262R	DEFATTED PEANUT MEAL (aflatoxin B1, blank)	43
BCR-263R	DEFATTED PEANUT MEAL (aflatoxin B1, medium level)	43
BCR-264	DEFATTED PEANUT MEAL (aflatoxin B1, high level)	43
BCR-265	DIBENZO[a,e]FLUORANTHENE (89mylas)	4
BCR-266	7H-DIBENZO (c,g) CARBAZOLE (purity)	4
BCR-267	INDENO[1,2,3-cd]FLUORANTHENE (purity)	4
BCR-269	CHRYSENE (purity)	4
BCR-270	TRIPHENYLENE (purity)	4
BCR-271	BENZ[a]ANTHRACENE (purity)	4
BCR-272	CORONENE (purity)	4
BCR-273	SINGLE CELL PROTEIN (major elements)	46
BCR-274	SINGLE CELL PROTEIN (trace elements)	46
BCR-275	ZIRCALOY (C, N, O)	74
BCR-276	ZIRCALOY (C, N, O)	74
BCR-277R	ESTUARINE SEDIMENT (trace elements)	14
BCR-280R	LAKE SEDIMENT (trace elements)	14
BCR-286A, B	ELECTROLYTICALLY REFINED LEAD (trace elements)	74
BCR-287A, B	THERMALLY REFINED LEAD (trace elements)	74
BCR-288B	LEAD WITH ADDED IMPURITIES (trace elements)	74
BCR-289	2,4'-DICHLOROBIPHENYL (IUPAC No. 8) (purity)	5
BCR-290	2,3,3'-TRICHLOROBIPHENYL (IUPAC No. 20) (purity)	5
BCR-291	2,4,4'-TRICHLOROBIPHENYL (IUPAC No. 28) (purity)	5
BCR-293	2,2',5,5'-TETRACHLOROBIPHENYL (IUPAC No. 52) (purity)	5
BCR-296	2,2',3,4,4',5'-HEXACHLOROBIPHENYL (IUPAC No. 138) (purity)	5
BCR-297	2,2',4,4',5,5'-HEXACHLOROBIPHENYL (IUPAC No. 153) (purity)	5
BCR-298	2,2',3,4,4',5,5'-HEPTACHLOROBIPHENYL (IUPAC No. 180) (purity)	5
BCR-301 (RM)	MULLITE (lattice spacing, other parameters)	68
BCR-304	HUMAN SERUM (Ca, Mg, Li)	59
BCR-305	1-NITROPYRENE (purity)	5
BCR-306	1-NITRONAPHTALENE (purity)	5
BCR-307	2-NITRONAPHTALENE (purity)	5
BCR-308	9-NITROANTHRACENE (purity)	5
BCR-309	6-NITROCHRYSENE (purity)	5
BCR-310	3-NITROFLUORANTHENE (purity)	5
BCR-311	6-NITROBENZO[a]PYRENE (purity)	5
BCR-312	2-NITRO-7-METHOXYNAPHTHO[2,1-b]FURAN (purity)	5
BCR-318	TITANIUM (H)	74
BCR-320R	CHANNEL SEDIMENT (trace elements)	14
BCR-321	UNALLOYED ZINC (trace elements)	74
BCR-326	UNALLOYED ZINC (disc) (trace elements)	74
BCR-327	UNALLOYED ZINC (disc) (trace elements)	74
BCR-331	LOW VOLATILE STEAM COAL (S)	76
BCR-332	HIGH VOLATILE INDUSTRIAL COAL (S)	76
BCR-336	HIGH VOLATILE STEAM COAL (S)	76
BCR-337	DIBENZO[b,d]FURAN (89mylas)	5
BCR-339	BENZO[c,d]PYREN-6-ONE (purity)	5

BCR-340	BENZO[b]NAPHTHO (1,2-d) FURAN (purity)	5
BCR-341	BENZO[b]NAPHTHO (2,1-d) FURAN (purity)	5
BCR-342	BENZO[a]FLUORENONE (purity)	5
BCR-348R	HUMAN SERUM (high progesterone)	59
BCR-349	COD LIVER OIL (PCBs)	41
BCR-351	ZnAl4 (trace elements)	75
BCR-352	ZnAl4 (trace elements)	75
BCR-353	ZnAl4 (trace elements)	75
BCR-354	ZnAl4 (trace elements)	75
BCR-355	ZnAl4 (trace elements)	75
BCR-356	ZnAl4Cu1 (trace elements)	75
BCR-357	ZnAl4Cu1 (trace elements)	75
BCR-360	ZnAl4Cu1 (trace elements)	75
BCR-361	ZnAl4Cu1 (trace elements)	75
BCR-365	PCB STANDARD SOLUTION	5
BCR-375	COMPOUND FEED (aflatoxin B1 blank)	43
BCR-377	MAIZE FLOUR (deoxynivalenol blank)	43
BCR-383	HARICOTS VERTS (major nutrients)	46, 50
BCR-385R	PEANUT BUTTER (aflatoxins low level)	43
BCR-393	HUMAN APOLIPOPROTEIN A I (mass concentration)	60
BCR-396	WHEAT FLOUR (deoxynivalenol blank)	43
3BCR-401R	PEANUT BUTTER (aflatoxins very low level)	43
BCR-402	WHITE CLOVER (trace elements)	16
BCR-405 (RM)	HUMAN HAEMOLYSATE (glycated haemoglobin (HbA1c)	61
BCR-412	BOVINE MUSCLE (diethylstilbestrol blank)	54
BCR-414	PLANKTON (trace elements)	17
BCR-420	WASTE MINERAL OIL (low PCB level)	27
BCR-423 (RM)	AFLATOXIN M1 STANDARD SOLUTION	29
BCR-425	NIMONIC 75 FOR CREEP TESTING	66
BCR-431	BRUSSELS SPROUT (vitamins)	49
BCR-444	PORCINE MUSCLE (chloramphenicol blank)	54
BCR-446	RAPESEED (oil, moisture, volatiles)	50
BCR-447	RAPESEED (oil, moisture, volatiles)	50
BCR-449	WASTE MINERAL OIL (high PCB level)	27
BCR-450	NATURAL MILK POWDER (PCBs)	41
BCR-457	HUMAN THYROGLOBULIN (Tg) (mass concentration)	60
BCR-459	COCONUT OIL (PAH blank)	40
BCR-460	COAL (F)	76
BCR-461	CLAY (F)	76
BCR-462	COASTAL SEDIMENT (butyltins)	23
BCR-465	RICE FLOUR (90mylase, low level)	49
BCR-466	RICE FLOUR (90mylase, medium level)	49
BCR-467	RICE FLOUR (90mylase, high level)	49
BCR-471	WHEAT (ochratoxin A, blank)	44
BCR-474-5	BOVINE LIVER (trenbolone blank and positive)	54
BCR-479	FRESH WATER (nitrate, low level)	20
BCR-480	FRESH WATER (nitrate, high level)	20
BCR-481	INDUSTRIAL SOIL (PCBs)	25
BCR-482	LICHEN (trace elements)	17
BCR-483	SEWAGE SLUDGE AMENDED SOIL (trace elements)	22
BCR-484	SEWAGE SLUDGE AMENDED (terra rossa) SOIL (trace elements)	22
BCR-485	MIXED VEGETABLES (vitamins)	49
BCR-486	PURIFIED HUMAN ALFAFOETOPROTEIN (protein mass)	60
BCR-487	PIG LIVER (vitamins)	49

BCR-490	FLY ASH (PCDDs and PCDFs)	26
BCR-502	BOVINE URINE (clenbuterol and salbutamol)	54
BCR-503	BOVINE URINE (clenbuterol and salbutamol)	54
BCR-504	BOVINE URINE (clenbuterol and salbutamol)	54
BCR-505	ESTUARINE WATER (trace elements)	19
BCR-519	ANHYDROUS BUTTER FAT (triglycerides)	48
BCR-522	BOVINE BLOOD LYSATE (haemoglobincyanide)	61
BCR-524	INDUSTRIAL SOIL (PAHs)	25
BCR-528	BACILLUS CEREUS (number of colony forming particles)	51
BCR-529	INDUSTRIAL SANDY SOIL (PCDDs, PCDFs)	25
BCR-530	INDUSTRIAL CLAY SOIL (PCDDs, PCDFs)	25
BCR-535	FRESHWATER HARBOUR SEDIMENT (PAHs)	25
BCR-536	FRESHWATER HARBOUR SEDIMENT (PCBs)	25
BCR-537	PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film A))	51
BCR-539	PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film C))	51
BCR-543	MUSSEL (dc-saxitoxin)	44
BCR-545	WELDING DUST LOADED ON FILTER (Cr VI, Cr)	24
BCR-546	FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	57
BCR-547	ACETALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	57
BCR-548	ACROLEIN-2,4-DINITROPHENYLHYDRAZONE (purity)	57
BCR-549	ACETONE-2,4-DINITROPHENYLHYDRAZONE (purity)	57
BCR-550	GLUTARALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	57
BCR-551	2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE	57
BCR-552	2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE (blank)	57
BCR-553	FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter	57
BCR-554	FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter (blank)	57
BCR-555	CHLORINATED HYDROCARBONS ON TENAX	58
BCR-573	HUMAN SERUM (low creatinine)	61
BCR-573i	CREATININE (interfering substances)	62
BCR-574	HUMAN SERUM (medium creatinine)	61
BCR-575	HUMAN SERUM (high creatinine)	61
BCR-576	HUMAN SERUM (17 β -ESTRADIOL, low level)	59
BCR-577	HUMAN SERUM (17 β -ESTRADIOL, medium level)	59
BCR-578	HUMAN SERUM (17 β -ESTRADIOL, high level)	59
BCR-579	COASTAL SEAWATER (Hg)	19
BCR-596	AQUATIC PLANT (Cr)	16
BCR-598	COD LIVER OIL (Organochlorine pesticides)	41
BCR-599	EWES'/GOATS' CURD (for adulteration with cows' milk) 0 and 1 % cows' milk	56
BCR-607	MILK POWDER (PCDDs, PCDFs)	42
BCR-611	GROUND WATER (Br, high level)	20
BCR-612	GROUND WATER (Br, low level)	20
BCR-613	PROSTATE SPECIFIC ANTIGEN (protein mass)	60
BCR-614	POLYCHLORODIBENZO-P-DIOXINS (PCDD) AND POLYCHLORODIBENZOFURANS (PCDFs)	6
BCR-615	FLY ASH (LOW LEVEL) (PCDDs and PCDFs)	26
BCR-627	TUNA FISH TISSUE (As species)	24
BCR-632	BUTTER FAT	48
BCR-633	ANHYDROUS BUTTER FAT (tracers)	47
BCR-634	HUMAN BLOOD (Pb, Cd)	59
BCR-635	HUMAN BLOOD (Pb, Cd)	59
BCR-636	HUMAN BLOOD (Pb, Cd)	59
BCR-637	HUMAN SERUM (Al, Se, Zn)	60
BCR-638	HUMAN SERUM (Al, Se, Zn)	60
BCR-639	HUMAN SERUM (Al, Se, Zn)	60
BCR-644	ARTIFICIAL FOODSTUFF (major nutrients)	49

BCR-645	ARTIFICIAL FOODSTUFF (major nutrients)	49
BCR-646	FRESHWATER SEDIMENT (butyltin and phenyltin compounds)	23
BCR-647	HUMAN ADENOSINE DEAMINASE (ADA 1)	62
BCR-648	BOVINE LIVER (CLENBUTEROL BLANK)	54
BCR-649	BOVINE LIVER (CLENBUTEROL POSITIVE)	54
BCR-651	BEER (EtOH, low level)	49
BCR-652	BEER (EtOH, very low level)	49
BCR-653	WINE (EtOH, low level)	49
BCR-656	ISOTOPE RATIOS IN ABSOLUTE ALCOHOL	29, 80
BCR-657	ISOTOPE RATIOS IN GLUCOSE	29, 80
BCR-660	ISOTOPE RATIOS IN ALCOHOLIC SOLUTION	29, 80
BCR-661	NIMONIC 75 FOR TENSILE PROPERTIES	66
BCR-663	SAXITOXIN IN ACETIC ACID	30
BCR-664	GLASS (trace elements)	78
BCR-665	LUNG TISSUE (asbestos fibres)	64
BCR-666	LUNG TISSUE (asbestos fibres)	64
BCR-667	ESTUARINE SEDIMENT	13
BCR-668	MUSSEL TISSUE	19
BCR-670	LEMNA MINOR (aquatic plant)	18
BCR-673	BOVINE EYE (CLENBUTEROL BLANK)	54
BCR-674	BOVINE EYE (CLENBUTEROL POSITIVE)	54
BCR-677	SEWAGE SLUDGE (PCDDs and PCDFs)	26
BCR-679	TRACE ELEMENTS IN WHITE CABBAGE	46
BCR-682	MUSSEL TISSUE	27
BCR-683	BEECH WOOD	27
BCR-684	RIVER SEDIMENT (extractable phosphorous)	22
BCR-685	SKIM MILK POWDER	50
BCR-691	COPPER ALLOYS	76
BCR-692	SCRATCH TESTING	67
BCR-693	HUMAN PANCREATIC LIPASE (from pancreatic juice)	62
BCR-694	HUMAN PANCREATIC LIPASE (recombinant)	62
BCR-695	PIG LIVER (CTC free)	55
BCR-696	PIG LIVER (CTC incurred)	55
BCR-697	PIG MUSCLE (CTC free)	55
BCR-700	ORGANIC-RICH SOIL (extractable elements)	22
BCR-701	LAKE SEDIMENT (trace elements)	23
BCR-704	FAUJASITE TYPE ZEOLITE (micropore volume and width)	68
BCR-705	LINDE TYPE A ZEOLITE (micropore volume and width)	68
BCR-706	PIG KIDNEY (CTC free)	55
BCR-707	PIG KIDNEY (CTC incurred)	55
BCR-708	DAIRY FEED (nutritional properties)	50
BCR-709	PIG FEED (nutritional properties)	50
BCR-718	HERRING (PCBs)	27
BCR-723	ROAD DUST (trace elements)	15
BCR-724A-D	GLASS-CERAMIC	65
BCR-725	SALMON TISSUE	55
IRMM/IFCC-456	ALPHA-AMYLASE (catalytic concentration)	62
IRMM/IFCC-467	HAEMOGLOBIN HbA0	62
IRMM-007/1-6	Zinc-64, nitrate solution	80
IRMM-009	Mg, isotopic, nitrate solution	81
IRMM-010	PLATINUM, isotopic, metal	81
IRMM-011	BORIC ACID, isotopic, solid	81
IRMM-012	CHROMIUM, isotopic, chloride solution	81

IRMM-016	LITHIUM CARBONATE, isotopic, solid	81
IRMM-018a	SILICON DIOXIDE, isotopic, solid	81
IRMM-311	Genomic DNA of <i>Bacillus licheniformis</i> DSM 5749	51
IRMM-312	Genomic DNA of <i>Bacillus subtilis</i> DSM 5750	52
IRMM-313	Genomic DNA of <i>Campylobacter coli</i> (CNET068) and <i>Campylobacter jejuni</i> (CNET112)	52
IRMM-315	4-DEOXYNIVALENOL in acetonitrile	31
IRMM-316	NIVALENOL in acetonitrile	31
IRMM-351	ESCHERICHIA COLI 0157 (NCTC 12900)	52
IRMM-352	SALMONELLA ENTERITIDIS (NCTC 12694)	52
IRMM-354	CANDIDA ALBICANS (NCPP 3179)	53
IRMM-355	ENTEROCOCCUS FAECALIS (CIP 106877)	53
IRMM-359	CHEESE (SEA)	44
IRMM-3702	ZINC-64 spike, nitrate solution	83
IRMM-426	WILD BERRIES	57
IRMM-427	PIKE-PERCH (PFASs in fish tissue)	27
IRMM-428	WATER (PFASs in water)	28
IRMM-435	PHARMACEUTICAL GLASS	64
IRMM-440	RESIN-BONDED FIBRE BOARD (thermal conductivity)	64
IRMM-441	n-HEPTANE (purity)	71, 74
IRMM-442	ISOOCTANE (purity)	71, 74
IRMM-443-1	EUROSOIL (adsorption coefficients and pH)	28
IRMM-443-2	EUROSOIL (adsorption coefficients and pH)	28
IRMM-443-3	EUROSOIL (adsorption coefficients and pH)	28
IRMM-443-4	EUROSOIL (adsorption coefficients and pH)	28
IRMM-443-5	EUROSOIL (adsorption coefficients and pH)	28
IRMM-443-7	EUROSOIL (adsorption coefficients and pH)	28
IRMM-447	Genomic DNA of <i>Listeria monocytogenes</i>	53
IRMM-448	Genomic DNA of <i>Campylobacter jejuni</i>	53
IRMM-449	Genomic DNA of <i>Escherichia coli</i>	53
IRMM-468	THYROXINE (T4)	58
IRMM-469	3,3',5' TRIIODOTHYRONINE (T3)	58
IRMM-471	CEMENTITE GRAINS IN CARBURISED PURE IRON	79
IRMM-481	Peanut Test Material Kit	56
IRMM-521	Ni	73
IRMM-522	Cu	73
IRMM-523	Al	73
IRMM-524	Fe	73
IRMM-525	Nb	73
IRMM-526	Nb	73
IRMM-527R	Al-0.1 % Co	73
IRMM-528R	Al-1.0 % Co	73
IRMM-529	Rh	73
IRMM-530R	Al-0.1 % Au	73
IRMM-531	Ti	73
IRMM-532	Al-0.01 % Co	73
IRMM-533	Al-0.1 % Ag	73
IRMM-534	Al-2.0 % Sc	73
IRMM-540R	OXIDE GLASS (15 ppm U)	78
IRMM-541	OXIDE GLASS (50 ppm U)	78
IRMM-610	BORON-10 spike, aqueous solution	82
IRMM-611	BORON (natural) spike, aqueous solution	82
IRMM-615	LITHIUM-6 spike, chloride solution	82
IRMM-618	RUBIDIUM-87 spike, nitrate solution	82
IRMM-619	RUBIDIUM (natural) spike, nitrate solution	82

IRMM-620	IRON-57 spike, chloride solution	82
IRMM-621	CADMIUM-111 spike, nitrate solution	82
IRMM-622	CADMIUM-111 spike, nitrate solution	82
IRMM-624	CHROMIUM-50 spike, chloride solution	82
IRMM-625	CHROMIUM (natural) spike, chloride solution	82
IRMM-632	COPPER-65 spike, nitrate solution	82
IRMM-634	IRON (natural) spike, chloride solution	82
IRMM-643	SULPHUR-32 spike, nitrate solution	83
IRMM-644	SULPHUR-32 spike, nitrate solution	83
IRMM-645	SULPHUR-32 spike, nitrate solution	83
IRMM-646	SULPHUR-34 spike, nitrate solution	83
IRMM-651	ZINC-64 spike, nitrate solution	83
IRMM-652	ZINC-64 spike, nitrate solution	83
IRMM-653	ZINC-67 spike, nitrate solution	83
IRMM-654	ZINC-68 spike, nitrate solution	83
IRMM-801	COCOA BUTTER	47
IRMM-804	RICE FLOUR	45
STA-003m	TETRAMETHYLUREA	29
VDA 001-004	POLYETHYLENE (40, 75, 200, 400 mg/kg Cd)	79

ALPHABETICAL LIST

DESIGNATION	ERM / CRM	PAGE NO.
γ -GLUTAMYLTRANSFERASE (catalytic concentration)	ERM-AD452/IFCC	62
10-AZABENZO[a]PYRENE (purity)	BCR-092	4
1-METHYLBENZ[a]ANTHRACENE (purity)	BCR-093R	4
1-METHYLCHRYSENE (purity)	BCR-077R	4
1-NITRONAPHTALENE (purity)	BCR-306	5
1-NITROPYRENE (purity)	BCR-305	5
2,2',3,4,4',5,5'-HEPTACHLOROBIPHENYL (IUPAC No. 180) (purity)	BCR-298	5
2,2',3,4,4',5-HEXACHLOROBIPHENYL (IUPAC No. 138) (purity)	BCR-296	5
2,2',4,4',5,5'-HEXACHLOROBIPHENYL (IUPAC No. 153) (purity)	BCR-297	5
2,2',5,5'-TETRACHLOROBIPHENYL (IUPAC No. 52) (purity)	BCR-293	5
2,3,3'-TRICHLOROBIPHENYL (IUPAC No. 20) (purity)	BCR-290	5
2,4,4'-TRICHLOROBIPHENYL (IUPAC No. 28) (purity)	BCR-291	5
2,4'-DICHLOROBIPHENYL (IUPAC No. 8) (purity)	BCR-289	5
2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE	BCR-551	57
2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE (blank)	BCR-552	57
2-METHYLCHRYSENE (purity)	BCR-078R	4
2-NITRO-7-METHOXYNAPHTHO[2,1-b]FURAN (purity)	BCR-312	5
2-NITRONAPHTALENE (purity)	BCR-307	5
3 REFERENCE ETHANOLS (H, M, L) (for SNIF-NMR)	BCR-123A, B	29, 80
3,3',5 TRIIODOTHYRONINE (T3)	IRMM-469	58
3-METHYLCHRYSENE (purity)	BCR-079R	4
3-NITROFLUORANTHENE (purity)	BCR-310	5
4-DEOXYNIVALENOL in acetonitrile	IRMM-315	31
4-METHYLCHRYSENE (purity)	BCR-080R	4
5-METHYLCHRYSENE (purity)	BCR-081R	4
6-NITROBENZO[a]PYRENE (purity)	BCR-311	5
6-NITROCHRYSENE (purity)	BCR-309	5
7H-DIBENZO (c,g) CARBAZOLE (purity)	BCR-266	4
9-NITROANTHRACENE (purity)	BCR-308	5
ABETA 42 IN CEREBROSPINAL FLUID (CSF) (low level)	ERM-DA480/IFCC	61
ABETA 42 IN CEREBROSPINAL FLUID (CSF) (medium level)	ERM-DA481/IFCC	61
ABETA 42 IN CEREBROSPINAL FLUID (CSF) (high level)	ERM-DA482/IFCC	61
ACETALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-547	57
ACETONE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-549	57
ACROLEIN-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-548	57
AFLATOXIN B1 IN ACETONITRILE	ERM-AC057	30
AFLATOXIN B2 IN ACETONITRILE	ERM-AC058	30
AFLATOXIN G1 IN ACETONITRILE	ERM-AC059	30
AFLATOXIN G2 IN ACETONITRILE	ERM-AC060	27
AFLATOXIN M1 STANDARD SOLUTION	BCR-423 (RM)	29
Al	IRMM-523	73
Al-0.01 % Co	IRMM-532	73
Al-0.1 % Ag	IRMM-533	73
Al-0.1 % Au	IRMM-530R	73
Al-0.1 % Au (0.1 mm foil)	ERM-EB530A	73
Al-0.1 % Au (0.5 mm wire)	ERM-EB530B	73
Al-0.1 % Au (1.0 mm wire)	ERM-EB530C	73
Al-0.1 % Co	IRMM-527R	73
Al-1.0 % Co	IRMM-528R	73

AI-2.0 % Sc	IRMM-534	73
ALANINE AMINOTRANSFERASE (ALT)	ERM-AD454k/IFCC	62
ALPHA ALUMINA (0.10 m ² /g) (nitrogen BET specific surface area)	BCR-169	68
ALPHA ALUMINA (1.05 m ² /g) (nitrogen BET specific surface area)	BCR-170	68
ALPHA-AMYLASE (catalytic concentration)	IRMM/IFCC-456	62
ALUMINA (2.95 m ² /g) (nitrogen BET specific surface area)	BCR-171	68
ANHYDROUS BUTTER FAT (tracers)	BCR-633	47
ANHYDROUS BUTTER FAT (triglycerides)	BCR-519	48
ANIMAL FEED (Organochlorine pesticides)	BCR-115	42
ANTHANTHRENE (purity)	BCR-091	4
ANTIMONY IMPLANTED IN SILICON	ERM-EG001	79
APPLE (dietary fibre)	ERM-BC516	51
AQUATIC PLANT (Cr)	BCR-596	16
ARSENOBETAINE IN WATER	ERM-AC626	30
ARTIFICIAL FOODSTUFF (major nutrients)	BCR-644	49
ARTIFICIAL FOODSTUFF (major nutrients)	BCR-645	49
ASPARTATE TRANSAMINASE (AST)	ERM-AD457/IFCC	62
BACILLUS CEREUS (number of colony forming particles)	BCR-528	51
BCR-ABL pDNA CALIBRANT	ERM-AD623	63
BEECH WOOD	BCR-683	27
BEEF-PORK FAT BLEND (fatty acid profile)	BCR-163	47
BEER (EtOH, low level)	BCR-651	49
BEER (EtOH, very low level)	BCR-652	49
BENZ[a]ACRIDINE (purity)	BCR-157	4
BENZ[a]ANTHRACENE (purity)	BCR-271	4
BENZ[c]ACRIDINE (purity)	BCR-158	4
BENZO[a]FLUORANTHENE (purity)	BCR-097	4
BENZO[a]FLUORENONE (purity)	BCR-342	5
BENZO[b]CHRYSENE (purity)	BCR-046	4
BENZO[b]FLUORANTHENE (purity)	BCR-047	4
BENZO[b]NAPHTHO (1,2-d) FURAN (purity)	BCR-340	5
BENZO[b]NAPHTHO (2,1-d) FURAN (purity)	BCR-341	5
BENZO[b]NAPHTHO[1,2-d]THIOPHENE (purity)	BCR-137R	4
BENZO[b]NAPHTHO[2,3-d]THIOPHENE (purity)	BCR-136R	4
BENZO[c,d]PYREN-6-ONE (purity)	BCR-339	5
BENZO[c]CHRYSENE (purity)	BCR-140	4
BENZO[c]PHENANTHRENE (purity)	BCR-134	4
BENZO[e]PYRENE (purity)	BCR-050	4
BENZO[ghi]FLUORANTHENE (purity)	BCR-139	4
BENZO[ghi]PERYLENE (purity)	BCR-052	4
BENZO[jj]FLUORANTHENE (purity)	BCR-049	4
BENZO[k]FLUORANTHENE (purity)	BCR-048R	4
BIODIESEL	ERM-EF001	71
BIODIESEL (B100 Rapeseed)	ERM-EF002	65, 71
BLADDERWRACK (<i>Fucus vesiculosus</i>) (trace elements)	ERM-CD200	17
BORIC ACID, isotopic, solid	IRMM-011	81
BORON (natural) spike, aqueous solution	IRMM-611	82
BORON-10 spike, aqueous solution	IRMM-610	82
BOVINE BLOOD (Pb, Cd)	ERM-CE195	59
BOVINE BLOOD (Pb, Cd)	ERM-CE196	59
BOVINE BLOOD LYSATE (haemiglobincyanide)	BCR-522	61
BOVINE EYE (CLENBUTEROL BLANK)	BCR-673	54
BOVINE EYE (CLENBUTEROL POSITIVE)	BCR-674	54
BOVINE LIVER (CLENBUTEROL BLANK)	BCR-648	54

BOVINE LIVER (CLENBUTEROL POSITIVE)	BCR-649	54
BOVINE LIVER (trace elements)	BCR-185R	45
BOVINE LIVER (trenbolone blank and positive)	BCR-474-5	54
BOVINE MUSCLE (diethylstilbestrol blank)	BCR-412	54
BOVINE MUSCLE (trace elements)	ERM-BB184	45
BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol)	ERM-BB389	54
BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol) (blank)	ERM-BB386	54
BOVINE URINE (clenbuterol and salbutamol)	BCR-502	54
BOVINE URINE (clenbuterol and salbutamol)	BCR-503	54
BOVINE URINE (clenbuterol and salbutamol)	BCR-504	54
BRAN BREAKFAST CEREAL (dietary fibre)	ERM-BD518	51
BROWN BREAD (trace elements)	BCR-191	45
BROWN COAL	ERM-EF412	65
BRUSSELS SPROUT (vitamins)	BCR-431	49
BUTTER FAT	BCR-632	48
CADMIUM-111 spike, nitrate solution	IRMM-621	82
CADMIUM-111 spike, nitrate solution	IRMM-622	82
CALCAREOUS SOIL	ERM-CC690	13
CALCIUM AMMONIUM NITRATE FERTILIZER (composition)	BCR-178	72
CALCIUM-41 isotopic, nitrate solution (set of 8 units)	ERM-AE701	81
CANDIDA ALBICANS (NCPF 3179)	IRMM-354	53
CARROT (dietary fibre)	ERM-BC515	51
CEMENTITE GRAINS IN CARBURISED PURE IRON	IRMM-471	79
CHANNEL SEDIMENT (trace elements)	BCR-320R	14
CHARPY SPECIMENS 120 J (impact toughness)	ERM-FA016	67
CHARPY SPECIMENS 150 J (impact toughness)	ERM-FA415	67
CHARPY SPECIMENS 80 J (impact toughness)	ERM-FA015	67
CHARPY SPECIMENS Low Energy 0°C (impact toughness)	ERM-FA013	67
CHARPY SPECIMENS Low Energy 20°C (impact toughness)	ERM-FA013	67
CHEESE (SEA)	IRMM-359	44
CHLORIDE (natural) spike, chloride solution	ERM-AE641	83
CHLORIDE-37 spike, chloride solution	ERM-AE642	83
CHLORINATED HYDROCARBONS ON TENAX	BCR-555	58
CHROMIUM (natural) spike, chloride solution	IRMM-625	82
CHROMIUM, isotopic, chloride solution	IRMM-012	81
CHROMIUM-50 spike, chloride solution	IRMM-624	82
CHRYSENE (purity)	BCR-269	4
CLAY (F)	BCR-461	76
COAL (F)	BCR-460	76
COASTAL SEAWATER (Hg)	BCR-579	19
COASTAL SEDIMENT (butyltins)	BCR-462	23
COCOA BUTTER	IRMM-801	47
COCONUT OIL (PAH blank)	BCR-459	4
COD LIVER OIL (Organochlorine pesticides)	BCR-598	41
COD LIVER OIL (PCBs)	BCR-349	41
COLLOIDAL SILICA (particle size)	ERM-FD100	69
COLLOIDAL SILICA (particle size)	ERM-FD304	69
COLLOIDAL SILICA IN AQUEOUS SOLUTION (nanoparticles)	ERM-FD101b	69
COLLOIDAL SILICA IN AQUEOUS SOLUTION (particle size)	ERM-FD102	69
COMPOUND FEED (aflatoxin B1 blank)	BCR-375	43
COMPOUND FEEDINGSTUFF (high level)	ERM-BE376	43
COMPOUND FEEDINGSTUFF (very low level)	ERM-BE375	43
CONTINUOUS CAST COPPER (O)	BCR-058	74
COPPER (natural) spike, nitrate solution	ERM-AE633	82

COPPER (O)	BCR-054R	74
COPPER (S, P)	BCR-017A, B	74
COPPER ALLOYS	BCR-691	76
COPPER-63, nitrate solution	ERM-AE647	83
COPPER-65 spike, nitrate solution	IRMM-632	82
CORONENE (purity)	BCR-272	4
CORTISOL REFERENCE SERUM PANEL	ERM-DA451/IFCC	58
CORUNDUM	ERM-FD066	68
CORUNDUM	ERM-FD069	68
CREATINE KINASE ISOENZYME MM (CK-MM)	ERM-AD455k/IFCC	62
CREATININE (interfering substances)	BCR-573i	62
Cu	IRMM-522	73
DAIRY FEED (nutritional properties)	BCR-708	50
DEFATTED PEANUT MEAL (aflatoxin B1, blank)	BCR-262R	43
DEFATTED PEANUT MEAL (aflatoxin B1, high level)	BCR-264	43
DEFATTED PEANUT MEAL (aflatoxin B1, medium level)	BCR-263R	43
DIBENZ[a,c]ACRIDINE (98mylas)	BCR-155	4
DIBENZ[a,h]ACRIDINE (purity)	BCR-153R	4
DIBENZ[a,l]ACRIDINE (98mylas)	BCR-152	4
DIBENZ[a,j]ACRIDINE (purity)	BCR-154	4
DIBENZ[a,c]ANTHRACENE (purity)	BCR-094	4
DIBENZ[a,j]ANTHRACENE (purity)	BCR-095	4
DIBENZ[c,h]ACRIDINE (purity)	BCR-156R	4
DIBENZO[a,e]FLUORANTHENE (98mylas)	BCR-265	4
DIBENZO[a,h]PYRENE (98mylas)	BCR-159	4
DIBENZO[a,e]PYRENE (98mylas)	BCR-133	4
DIBENZO[a,h]ANTHRACENE (purity)	BCR-138	4
DIBENZO[a,l]PYRENE (98mylas)	BCR-096	4
DIBENZO[b,d]FURAN (98mylas)	BCR-337	5
ELECTROLYTIC COPPER WITH ADDED IMPURITIES (trace elements)	ERM-EB075A-B-C	76
ELECTROLYTIC COPPER (trace elements)	ERM-EB074A-B-C	76
ELECTROLYTIC COPPER (trace elements)	BCR-074A	76
ELECTROLYTIC TOUGH PITCH COPPER (O)	BCR-022A, B	74
ELECTROLYTICALLY REFINED LEAD (trace elements)	BCR-286A, B	74
ENTEROCOCCUS FAECALIS (CIP 106877)	IRMM-355	53
ESCHERICHIA COLI 0157 (NCTC 12900)	IRMM-351	52
ESTUARINE SEDIMENT	BCR-667	13
ESTUARINE SEDIMENT (Hg, methylmercury)	ERM-CC580	14, 23
ESTUARINE SEDIMENT (trace elements)	BCR-277R	14
ESTUARINE WATER (trace elements)	BCR-505	19
EUROSOIL (adsorption coefficients and pH)	IRMM-443-1	28
EUROSOIL (adsorption coefficients and pH)	IRMM-443-2	28
EUROSOIL (adsorption coefficients and pH)	IRMM-443-3	28
EUROSOIL (adsorption coefficients and pH)	IRMM-443-4	28
EUROSOIL (adsorption coefficients and pH)	IRMM-443-5	28
EUROSOIL (adsorption coefficients and pH)	IRMM-443-7	28
EWES'/GOATS' CURD (for adulteration with cows' milk) 0 and 1 % cows' milk	BCR-599	56
FAUJASITE TYPE ZEOLITE (mircopore volume and width)	BCR-704	68
Fe	IRMM-524	73
FINE DUST (PM ₁₀ -like) (elements)	ERM-CZ120	15
FINE DUST (PM ₁₀ -like) (PAHs)	ERM-CZ100	24
FISH MUSCLE (trace elements)	ERM-BB422	45
FISH OIL	ERM-BB350	41
FISH TISSUE	ERM-CE100	28

FLUORANTHENE (purity)	BCR-160R	4
FLY ASH (LOW LEVEL) (PCDDs and PCDFs)	BCR-615	26
FLY ASH (PCDDs and PCDFs)	BCR-490	26
FLY ASH (trace elements)	BCR-176R	15
FLY ASH FROM PULVERISED COAL (trace elements)	BCR-038	15
FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-546	57
FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter	BCR-553	57
FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter (blank)	BCR-554	57
FRESH WATER (nitrate, high level)	BCR-480	20
FRESH WATER (nitrate, low level)	BCR-479	20
FRESHWATER HARBOUR SEDIMENT (PAHs)	BCR-535	25
FRESHWATER HARBOUR SEDIMENT (PCBs)	BCR-536	25
FRESHWATER SEDIMENT (butyltin and phenyltin compounds)	BCR-646	23
FULL FAT SOYA (dietary fibre)	ERM-BC517	51
FURNACE COKE	ERM-EF413	65
GAS OIL (0.0203 % S)	ERM-EF672	77
GAS OIL (0.0452 % S)	ERM-EF671	77
GAS OIL (0.1019 % S)	ERM-EF104	77
GAS OIL (0.363 % S)	BCR-105	77
GAS OIL (0.502 % S)	BCR-106	77
GAS OIL (1.040 % S)	BCR-107	77
GAS OIL (CFPP and CP)	ERM-FC395k	65
GAS OIL (Solvent Yellow 124)	ERM-EF317	77
GENETICALLE MODIFIED AM04-1020 POTATO	ERM-BF430	37
GENETICALLE MODIFIED AV43-6-G7 POTATO	ERM-BF431	38
GENETICALLY MODIFIED DP-ØØ4114-3 MAIZE	ERM-BF439	40
GENETICALLY MODIFIED VCO-Ø1981-5 MAIZE	ERM-BF438	39
GENETICALLY MODIFIED 1507 MAIZE	ERM-BF418	34
GENETICALLY MODIFIED 281-24-236 X 3006-210-23 COTTON SEED	ERM-BF422	35
GENETICALLY MODIFIED 3272 MAIZE	ERM-BF420	34
GENETICALLY MODIFIED 59122 MAIZE	ERM-BF424	35
GENETICALLY MODIFIED 73496 RAPESEED	ERM-BF434	38
GENETICALLY MODIFIED 98140 MAIZE	ERM-BF427	36
GENETICALLY MODIFIED Bt-11 MAIZE	ERM-BF412	32
GENETICALLY MODIFIED Bt-176 MAIZE	ERM-BF411	31
GENETICALLY MODIFIED DAS-40278-9 MAIZE	ERM-BF433	38
GENETICALLY MODIFIED DAS-44406-6 SOYA	ERM-BF436	39
GENETICALLY MODIFIED DAS-81419-2 SOYA	ERM-BF437	39
GENETICALLY MODIFIED DAS-81910-7 COTTON	ERM-BF440	40
GENETICALLY MODIFIED EH92-527-1 POTATO	ERM-BF421	35
GENETICALLY MODIFIED GA21 MAIZE	ERM-BF414	33
GENETICALLY MODIFIED GHB119 COTTON	ERM-BF428	37
GENETICALLY MODIFIED H7-1 SUGAR BEET	ERM-BF419	34
GENETICALLY MODIFIED MIR604 MAIZE	ERM-BF423	35
GENETICALLY MODIFIED MON 810 MAIZE	ERM-BF413k	32
GENETICALLY MODIFIED MON 863 MAIZE	ERM-BF416	33
GENETICALLY MODIFIED MON 863 x MON 810 MAIZE	ERM-BF417	34
GENETICALLY MODIFIED NK603 MAIZE	ERM-BF415	33
GENETICALLY MODIFIED PH05-026-0048 POTATO	ERM-BF435	39
GENETICALLY MODIFIED ROUNDUP READY SOYA	ERM-BF410p	31
GENETICALLY MODIFIED SOYA 305423	ERM-BF426	36
GENETICALLY MODIFIED SOYA 356043	ERM-BF425	36
GENETICALLY MODIFIED SOYA DAS-68416-4	ERM-BF432	38
GENETICALLY MODIFIED T304-40 COTTON	ERM-BF429	37

Genomic DNA of <i>Bacillus licheniformis</i> DSM 5749	IRMM-311	51
Genomic DNA of <i>Bacillus subtilis</i> DSM 5750	IRMM-312	52
Genomic DNA of <i>Campylobacter coli</i> (CNET068) and <i>Campylobacter jejuni</i> (CNET112)	IRMM-313	52
Genomic DNA of <i>Campylobacter jejuni</i>	IRMM-448	53
Genomic DNA of <i>Escherichia coli</i>	IRMM-449	53
Genomic DNA of <i>Listeria monocytogenes</i>	IRMM-447	53
GLASS (trace elements)	BCR-664	78
GLASS-CERAMIC	BCR-724A-D	65
GLUTARALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-550	57
GROUND WATER	ERM-CA615	21
GROUND WATER	ERM-CA616	20
GROUND WATER (Br, high level)	BCR-611	20
GROUND WATER (Br, low level)	BCR-612	20
HAEMOGLOBIN HbA0	IRMM/IFCC-467	62
HAEMOGLOBIN IN BUFFER	ERM-AD500/IFCC	62
HARD COAL	ERM-EF411	65
HARICOTS BEANS (dietary fibre)	ERM-BC514	51
HARICOTS VERTS (major nutrients)	BCR-383	46, 50
HAY POWDER (elements)	BCR-129	16
HERRING (PCBs)	BCR-718	27
HIGH VOLATILE INDUSTRIAL COAL (S)	BCR-332	76
HIGH VOLATILE STEAM COAL (S)	BCR-336	76
HUMAN ADENOSINE DEAMINASE (ADA 1)	BCR-647	62
HUMAN APOLIPOPROTEIN A I (mass concentration)	BCR-393	60
HUMAN BLOOD (Pb, Cd)	BCR-634	59
HUMAN BLOOD (Pb, Cd)	BCR-635	59
HUMAN BLOOD (Pb, Cd)	BCR-636	59
HUMAN HAEMOLYSATE (glycated haemoglobin (HbA1c))	BCR-405 (RM)	61
HUMAN HAIR (trace elements)	ERM-DB001	60
HUMAN PANCREATIC LIPASE (from pancreatic juice)	BCR-693	62
HUMAN PANCREATIC LIPASE (recombinant)	BCR-694	62
HUMAN SERUM (17 β -ESTRADIOL, high level)	BCR-578	59
HUMAN SERUM (17 β -ESTRADIOL, low level)	BCR-576	59
HUMAN SERUM (17 β -ESTRADIOL, medium level)	BCR-577	59
HUMAN SERUM (Al, Se, Zn)	BCR-637	60
HUMAN SERUM (Al, Se, Zn)	BCR-638	60
HUMAN SERUM (Al, Se, Zn)	BCR-639	60
HUMAN SERUM (Ca, Mg, Li)	BCR-304	59
HUMAN SERUM (cortisol spiked)	ERM-DA193	59
HUMAN SERUM (cortisol unspiked)	ERM-DA192	59
HUMAN SERUM (CRP)	ERM-DA474/IFCC	61
HUMAN SERUM (cystatin C)	ERM-DA471/IFCC	61
HUMAN SERUM (high creatinine)	BCR-575	61
HUMAN SERUM (high progesterone)	BCR-348R	59
HUMAN SERUM (low creatinine)	BCR-573	61
HUMAN SERUM (medium creatinine)	BCR-574	61
HUMAN SERUM (progesterone)	ERM-DA347	59
HUMAN SERUM (proteins)	ERM-DA470k/IFCC	61
HUMAN THYROGLOBULIN (Tg) (mass concentration)	BCR-457	60
IgG ANTI-MPO IN HUMAN SERUM	ERM-DA476/IFCC	61
IgG PR3 ANCA IN HUMAN SERUM	ERM-DA483/IFCC	61
INDENO[1,2,3-cd]FLUORANTHENE (purity)	BCR-267	4
INDUSTRIAL CLAY SOIL (PCDDs, PCDFs)	BCR-530	25

INDUSTRIAL SANDY SOIL (PCDDs, PCDFs)	BCR-529	25
INDUSTRIAL SOIL (PAHs)	BCR-524	25
INDUSTRIAL SOIL (PCBs)	BCR-481	25
IRON (natural) spike, chloride solution	IRMM-634	82
IRON-57 spike, chloride solution	IRMM-620	82
ISOOCTANE (purity)	IRMM-442	71, 74
ISOTOPE RATIOS IN ABSOLUTE ALCOHOL	BCR-656	29, 80
ISOTOPE RATIOS IN ALCOHOLIC SOLUTION	BCR-660	29, 80
ISOTOPE RATIOS IN GLUCOSE	BCR-657	29, 80
LACTATE DEHYDROGENASE ISOENZYME 1 (LD1)	ERM-AD453k/IFCC	62
LAKE SEDIMENT (trace elements)	BCR-280R	14
LAKE SEDIMENT (trace elements)	BCR-701	23
LAMBDA DNA	ERM-AD442k	63
LATEX SPHERES (particle diameter 2 microns)	BCR-165	58
LATEX SPHERES (particle diameter 4.8 microns)	BCR-166	58
LATEX SPHERES (particle diameter 9.6 microns)	BCR-167	58
LEAD GLASS (composition/refractive index)	BCR-126A/B	73
LEAD WITH ADDED IMPURITIES (trace elements)	BCR-288B	74
LEMNA MINOR (aquatic plant)	BCR-670	18
LICHEN (trace elements)	BCR-482	17
LIGHT SANDY SOIL (trace elements)	BCR-142R	13
LIMESTONE POWDERS (for shear testing)	BCR-116	66
LINDE TYPE A ZEOLITE (mircopore volume and width)	BCR-705	68
LITHIUM CARBONATE, isotopic, solid	IRMM-016	81
LITHIUM-6 spike, chloride solution	IRMM-615	82
LOAM SOIL	ERM-CC141	13
LOW VOLATILE STEAM COAL (S)	BCR-331	76
LUNG TISSUE (asbestos fibres)	BCR-665	64
LUNG TISSUE (asbestos fibres)	BCR-666	64
MAGNESIUM (natural) spike, nitrate solution	ERM-AE637	82
MAGNESIUM-26 spike, nitrate solution	ERM-AE638	82
MAIZE	ERM-BC716	44
MAIZE	ERM-BC717	44
MAIZE FLOUR (deoxynivalenol blank)	BCR-377	43
MARGARINE (vitamins)	BCR-122	49
MERCURY (natural) spike, chloride solution	ERM-AE639	82
MERCURY-202 spike, chloride solution	ERM-AE640	83
METHYLMERCURY IN 2 % ETHANOL/WATER SOLUTION	ERM-AE671	83
Mg, isotopic, nitrate solution	IRMM-009	81
MILK POWDER (oxytetracycline)	ERM-BB492	55
MILK POWDER (oxytetracycline) (blank)	ERM-BB493	55
MILK POWDER (PCDDs, PCDFs)	BCR-607	42
MIXED VEGETABLES (vitamins)	BCR-485	49
MOROCCAN PHOSPHATE ROCK (trace elements)	BCR-032	72, 78
MULLITE (lattice spacing, other parameters)	BCR-301 (RM)	68
MUSSEL (dc-saxitoxin)	BCR-543	44
MUSSEL TISSUE	BCR-668	19
MUSSEL TISSUE	BCR-682	27
MUSSEL TISSUE (butyltins)	ERM-CE477	24
MUSSEL TISSUE (elements)	ERM-CE278k	18
NATURAL MILK POWDER (PCBs)	BCR-450	41
NATURAL MILK POWDER (pesticides)	BCR-187	42
NATURAL PORK FAT (blank)	ERM-BB444	41
Nb	IRMM-525	73

Nb	IRMM-526	73
n-HEPTANE (purity)	IRMM-441	71, 74
Ni	IRMM-521	73
NIMONIC 75 FOR CREEP TESTING	BCR-425	66
NIMONIC 75 FOR TENSILE PROPERTIES	BCR-661	66
NIVALENOL in acetonitrile	IRMM-316	31
ORGANIC-RICH SOIL (extractable elements)	BCR-700	22
OXIDE GLASS (15 ppm U)	IRMM-540R	78
OXIDE GLASS (50 ppm U)	IRMM-541	78
PAHs IN ACETONITRILE / TOLUENE	ERM-AC213	12
PCB STANDARD SOLUTION	BCR-365	5
PEANUT BUTTER (aflatoxins low level)	BCR-385R	43
PEANUT BUTTER (aflatoxins very low level)	3BCR-401R	43
Peanut Test Material Kit	IRMM-481	56
PETROL	ERM-EF211	77
PHARMACEUTICAL GLASS	IRMM-435	64
PICENE (purity)	BCR-168	4
PIG FEED (nutritional properties)	BCR-709	50
PIG KIDNEY (CTC free)	BCR-706	55
PIG KIDNEY (CTC incurred)	BCR-707	55
PIG KIDNEY (trace elements)	ERM-BB186	45
PIG LIVER (CTC free)	BCR-695	55
PIG LIVER (CTC incurred)	BCR-696	55
PIG LIVER (vitamins)	BCR-487	49
PIG MUSCLE (CTC free)	BCR-697	55
PIKE-PERCH (PFASs in fish tissue)	IRMM-427	27
PLANKTON (trace elements)	BCR-414	17
PLASMID DNA FRAGMENTS OF 356043 SOYBEAN	ERM-AD425	36
PLASMID DNA FRAGMENTS OF 98140 MAIZE	ERM-AD427	37
PLASMID DNA FRAGMENTS OF MON 810 MAIZE	ERM-AD413	32
PLASMID DNA FRAGMENTS OF NK603 MAIZE	ERM-AD415	33
PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film A)	BCR-537	51
PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film C)	BCR-539	51
PLATINUM, isotopic, metal	IRMM-010	81
POLYCHLORODIBENZO-P-DIOXINS (PCDD) AND POLYCHLORODIBENZOFURANS (PCDFs)	BCR-614	6
POLYETHYLENE (40, 75, 200, 400 mg/kg Cd)	VDA 001-004	79
POLYETHYLENE (high level)	ERM-EC681m	79
POLYETHYLENE (LDPE)	ERM-EC590	78
POLYETHYLENE (low level)	ERM-EC680m	79
POLYPROPYLENE (PP)	ERM-EC591	78
PORCINE MUSCLE (chloramphenicol blank)	BCR-444	54
PORCINE pDNA CALIBRANT	ERM-AD483	56
PORK FAT (pesticides)	ERM-BB430	42
PORK MUSCLE	ERM-BB124	55
PORK MUSCLE	ERM-BB130	54
PORK MUSCLE	ERM-BB384	46, 50
POTASSIUM CHLORIDE FERTILIZER (elemental composition)	BCR-113	72
POTASSIUM SULPHATE FERTILIZER (elemental composition)	BCR-114	72
PROSTATE SPECIFIC ANTIGEN (protein mass)	BCR-613	60
PURIFIED HUMAN ALFAFOETOPROTEIN (protein mass)	BCR-486	60
PYRENE (purity)	BCR-177R	4
QUARTZ (1.20 – 20.00 microns)	BCR-070	68
QUARTZ (2.50 m ² /g) (nitrogen BET specific surface area)	BCR-172	68

QUARTZ (particle size 0.35 – 3.50 microns)	BCR-066	68
QUARTZ (particle size 14 – 90 microns)	BCR-069	68
QUARTZ (particle size 1400 – 5000 microns)	BCR-132	68
QUARTZ (particle size 160 – 630 microns)	BCR-068	68
QUARTZ (particle size 2.40 – 32.00 microns)	BCR-067	68
QUARTZ (particle size 480 – 1800 microns)	BCR-131	68
QUARTZ (particle size 50 – 220 microns)	BCR-130	68
RAPESEED (colza) (S, total glucosinolate, high level)	ERM-BC367	43
RAPESEED (colza) (S, total glucosinolate, low level)	ERM-BC366	43
RAPESEED (colza) (S, total glucosinolate, medium level)	ERM-BC190	43
RAPESEED (oil, moisture, volatiles)	BCR-446	50
RAPESEED (oil, moisture, volatiles)	BCR-447	50
RESIN-BONDED FIBRE BOARD (thermal conductivity)	IRMM-440	64
Rh	IRMM-529	73
RICE (As species)	ERM-BC211	24
RICE FLOUR	IRMM-804	45
RICE FLOUR (103mylase, low level)	BCR-465	49
RICE FLOUR (103mylase, medium level)	BCR-466	49
RICE FLOUR (103mylase, high level)	BCR-467	49
RIVER SEDIMENT (extractable phosphorous)	BCR-684	22
ROAD DUST (trace elements)	BCR-723	15
RUBIDIUM (natural) spike, nitrate solution	IRMM-619	82
RUBIDIUM-87 spike, nitrate solution	IRMM-618	82
RUMINANT pDNA CALIBRANT	ERM-AD482	56
RYE FLOUR	ERM-BC381	46, 50
RYE GRASS	ERM-CD281	16
SALMON TISSUE	BCR-725	55
SALMONELLA ENTERITIDIS (NCTC 12694)	IRMM-352	52
SAXITOXIN IN ACETIC ACID	BCR-663	30
SCRATCH TESTING	BCR-692	67
SEAWATER (Hg)	ERM-CA400	21
SEWAGE SLUDGE	ERM-CC144	13
SEWAGE SLUDGE (industrial origin) (trace elements)	BCR-146R	14
SEWAGE SLUDGE (mixed origin) (trace elements)	BCR-145R	14
SEWAGE SLUDGE (PCDDs and PCDFs)	BCR-677	26
SEWAGE SLUDGE AMENDED (terra rossa) SOIL (trace elements)	BCR-484	22
SEWAGE SLUDGE AMENDED SOIL (trace elements)	BCR-143R	13
SEWAGE SLUDGE AMENDED SOIL (trace elements)	BCR-483	22
SILICON DIOXIDE, isotopic, solid	IRMM-018a	81
SIMULATED RAINWATER (major components)	ERM-CA408	20
SINGLE CELL PROTEIN (major elements)	BCR-273	46
SINGLE CELL PROTEIN (trace elements)	BCR-274	46
SKIM MILK POWDER	BCR-685	50
SKIMMED MILK POWDER (trace elements)	ERM-BD150	45
SKIMMED MILK POWDER (trace elements)	ERM-BD151	45
SOYA-MAIZE OIL BLEND (fatty acid profile)	BCR-162R	47
SPIKED MILK POWDER (pesticides)	BCR-188	42
SPIKED PORK FAT (low level)	ERM-BB446	41
SPIKED PORK FAT (very low level)	ERM-BB445	41
SULPHUR-32 spike, nitrate solution	IRMM-643	83
SULPHUR-32 spike, nitrate solution	IRMM-644	83
SULPHUR-32 spike, nitrate solution	IRMM-645	83
SULPHUR-34 spike, nitrate solution	IRMM-646	83
SUPERPHOSPHATE (various parameters)	BCR-033	72

SURFACE WATER	ERM-CA100	28
TANTALUM PENTOXIDE ON TANTALUM FOIL	BCR-261T	69
TETRAMETHYLUREA	STA-003m	29
THALLIUM (natural) spike, nitrate solution	ERM-AE649	83
THERMALLY REFINED LEAD (trace elements)	BCR-287A, B	74
THROMBOPLASTIN RABBIT (prothrombin time)	ERM-AD149	64
THYROXINE (T4)	IRMM-468	58
Ti	IRMM-531	73
Ti 6AL 4V ALLOY (O)	BCR-059A, B	74
TiAl6V4 (Al, V)	BCR-089	75
TIN ORE CONCENTRATE (Sn)	BCR-010	72
TITANIA (8.23 m ² /g) (nitrogen BET specific surface area)	BCR-173	68
TITANIUM (H)	BCR-318	74
TITANIUM (impurities)	BCR-090A, B	75
TITANIUM (O, N)	BCR-024B, C	74
TOasted BREAD	ERM-BD273	57
TRACE ELEMENTS IN WHITE CABBAGE	BCR-679	46
TRIPHENYLENE (purity)	BCR-270	4
TUNA FISH (total and methylmercury)	ERM-CE464	19, 24
TUNA FISH TISSUE (As species)	BCR-627	24
TUNGSTEN (0.18 m ² /g) (nitrogen BET specific surface area)	BCR-175	68
TUNGSTEN CARBIDE POWDER (O)	BCR-102	74
UNALLOYED ZINC (disc) (trace elements)	BCR-326	74
UNALLOYED ZINC (disc) (trace elements)	BCR-327	74
UNALLOYED ZINC (trace elements)	ERM-EB322	74
UNALLOYED ZINC (trace elements)	ERM-EB323	74
UNALLOYED ZINC (trace elements)	ERM-EB324	74
UNALLOYED ZINC (trace elements)	ERM-EB325	74
UNALLOYED ZINC (trace elements)	BCR-321	74
UREA FERTILIZER (composition)	BCR-179	72
WASTE MINERAL OIL (high PCB level)	BCR-449	27
WASTE MINERAL OIL (low PCB level)	BCR-420	27
WASTE WATER	ERM-CA713	21
WATER (PFASs in water)	IRMM-428	28
WELDING DUST LOADED ON FILTER (Cr VI, Cr)	BCR-545	24
WHEAT (ochratoxin A, blank)	BCR-471	44
WHEAT FLOUR	ERM-BC382	46, 50
WHEAT FLOUR (deoxynivalenol blank)	BCR-396	43
WHITE CLOVER (trace elements)	BCR-402	16
WHOLE MILK POWDER (aflatoxin M1, high level)	ERM-BD284	43
WHOLE MILK POWDER (aflatoxin M1, low level)	ERM-BD283	43
WHOLE MILK POWDER (aflatoxin M1, zero level)	ERM-BD282	43
WHOLE MILK POWDER (vitamins)	ERM-BD600	49
WHOLEMEAL FLOUR (vitamins)	BCR-121	49
WILD BERRIES	IRMM-426	57
WINE (EtOH, low level)	BCR-653	49
ZEARALENONE IN ACETONITRILE	ERM-AC699	30
ZINC ORE CONCENTRATE (trace elements)	BCR-109	77
ZINC ORE CONCENTRATE (trace elements)	BCR-110	77
ZINC-64 spike, nitrate solution	IRMM-3702	83
ZINC-64 spike, nitrate solution	IRMM-651	83
ZINC-64 spike, nitrate solution	IRMM-652	83
Zinc-64, nitrate solution	IRMM-007/1-6	80
ZINC-67 spike, nitrate solution	IRMM-653	83

ZINC-68 spike, nitrate solution	IRMM-654	83
ZIRCALOY (C, N, O)	BCR-275	74
ZIRCALOY (C, N, O)	BCR-276	74
ZIRCALOY-4 (trace element impurities)	BCR-098	75
ZnAl4 (trace elements)	BCR-351	75
ZnAl4 (trace elements)	BCR-352	75
ZnAl4 (trace elements)	BCR-353	75
ZnAl4 (trace elements)	BCR-354	75
ZnAl4 (trace elements)	BCR-355	75
ZnAl4Cu1 (trace elements)	BCR-356	75
ZnAl4Cu1 (trace elements)	BCR-357	75
ZnAl4Cu1 (trace elements)	BCR-360	75
ZnAl4Cu1 (trace elements)	BCR-361	75